


ENM 3600: Introduction to Data-driven Modeling

Lecture #5: Probability and Statistics primer




Correlation and linear dependence





Covariance vs Mutual Information

$$\text{cov}(X, Y) \quad I(X; Y)$$




@ari_seff

The multivariate Gaussian



$$p(\mathbf{x} \mid \boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-\frac{D}{2}} |\boldsymbol{\Sigma}|^{-\frac{1}{2}} \exp \left(-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right)$$


The multivariate Gaussian



Notation	$\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$
Parameters	$\boldsymbol{\mu} \in \mathbf{R}^k$ — location $\boldsymbol{\Sigma} \in \mathbf{R}^{k \times k}$ — covariance (positive semi-definite matrix)
Support	$\mathbf{x} \in \boldsymbol{\mu} + \text{span}(\boldsymbol{\Sigma}) \subseteq \mathbf{R}^k$
PDF	$\det(2\pi\boldsymbol{\Sigma})^{-\frac{1}{2}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}$, exists only when $\boldsymbol{\Sigma}$ is positive-definite
Mean	$\boldsymbol{\mu}$
Mode	$\boldsymbol{\mu}$
Variance	$\boldsymbol{\Sigma}$


Marginals and conditionals of a Gaussian

$$p(\mathbf{x}, \mathbf{y}) = \mathcal{N} \left(\begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}, \begin{bmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{yx} & \Sigma_{yy} \end{bmatrix} \right)$$



Marginal distribution

$$p(\mathbf{x}) = \int p(\mathbf{x}, \mathbf{y}) d\mathbf{y} = \mathcal{N}(\mathbf{x} | \mu_x, \Sigma_{xx})$$




Conditional distribution

$$p(\mathbf{x} | \mathbf{y}) = \mathcal{N}(\mu_{x|y}, \Sigma_{x|y})$$


$$\mu_{x|y} = \mu_x + \Sigma_{xy} \Sigma_{yy}^{-1} (\mathbf{y} - \mu_y)$$

$$\Sigma_{x|y} = \Sigma_{xx} - \Sigma_{xy} \Sigma_{yy}^{-1} \Sigma_{yx}$$




These are unique properties that make the Gaussian distribution very simple and attractive to compute with! It is essentially our main building block for computing under uncertainty.

Transformations



Maximum likelihood estimation

$$\theta_{\text{MLE}} = \arg \max_{\theta \in \Theta} p(\mathcal{D}|\theta)$$



Bayesian estimation

$$p(\theta|\mathcal{D}) = \frac{p(\mathcal{D}|\theta)p(\theta)}{p(\mathcal{D})}$$

