



# ENM 53 I: Data-driven Modeling and Probabilistic Scientific Computing

## *Lecture #3: Primer on Probability and Statistics*

Paris Perdikaris  
January 28, 2021




# The Gaussian distribution





<b>Notation</b>	$\mathcal{N}(\mu, \sigma^2)$
<b>Parameters</b>	$\mu \in \mathbb{R}$ = mean ( <b>location</b> ) $\sigma^2 > 0$ = variance (squared <b>scale</b> )
<b>Support</b>	$x \in \mathbb{R}$
<b>PDF</b>	$\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$
<b>CDF</b>	$\frac{1}{2} \left[ 1 + \operatorname{erf}\left(\frac{x-\mu}{\sigma\sqrt{2}}\right) \right]$
<b>Quantile</b>	$\mu + \sigma\sqrt{2} \operatorname{erf}^{-1}(2F - 1)$
<b>Mean</b>	$\mu$
<b>Median</b>	$\mu$
<b>Mode</b>	$\mu$
<b>Variance</b>	$\sigma^2$

# Correlation and linear dependence




# The multivariate Gaussian



$$p(\mathbf{x} \mid \boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-\frac{D}{2}} |\boldsymbol{\Sigma}|^{-\frac{1}{2}} \exp \left( -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right)$$

# The multivariate Gaussian




<b>Notation</b>	$\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$
<b>Parameters</b>	$\boldsymbol{\mu} \in \mathbf{R}^k$ — location $\boldsymbol{\Sigma} \in \mathbf{R}^{k \times k}$ — covariance (positive semi-definite matrix)
<b>Support</b>	$\mathbf{x} \in \boldsymbol{\mu} + \text{span}(\boldsymbol{\Sigma}) \subseteq \mathbf{R}^k$
<b>PDF</b>	$\det(2\pi\boldsymbol{\Sigma})^{-\frac{1}{2}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}$ , exists only when $\boldsymbol{\Sigma}$ is positive-definite
<b>Mean</b>	$\boldsymbol{\mu}$
<b>Mode</b>	$\boldsymbol{\mu}$
<b>Variance</b>	$\boldsymbol{\Sigma}$




# Marginals and conditionals of a Gaussian

$$p(\mathbf{x}, \mathbf{y}) = \mathcal{N} \left( \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}, \begin{bmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{yx} & \Sigma_{yy} \end{bmatrix} \right)$$



*Marginal distribution*

$$p(\mathbf{x}) = \int p(\mathbf{x}, \mathbf{y}) d\mathbf{y} = \mathcal{N}(\mathbf{x} \mid \mu_x, \Sigma_{xx})$$




*Conditional distribution*

$$p(\mathbf{x} \mid \mathbf{y}) = \mathcal{N}(\mu_{x \mid y}, \Sigma_{x \mid y})$$


$$\mu_{x \mid y} = \mu_x + \Sigma_{xy} \Sigma_{yy}^{-1} (\mathbf{y} - \mu_y)$$

$$\Sigma_{x \mid y} = \Sigma_{xx} - \Sigma_{xy} \Sigma_{yy}^{-1} \Sigma_{yx}$$



These are unique properties that make the Gaussian distribution very simple and attractive to compute with! It is essentially our main building block for computing under uncertainty.

# Transformations



# Maximum likelihood estimation

$$\theta_{\text{MLE}} = \arg \max_{\theta \in \Theta} p(\mathcal{D}|\theta)$$