


Practice Problems

1. Give the average value of $f(x) = \cos(2x)$ on the interval $\left[-\frac{\pi}{2}, \frac{\pi}{4}\right]$.
2. The graph of $f(x)$ is shown below.



- a. Give the area of the region bounded between the graph of $f(x)$ and the x -axis on the interval $[-2, 3]$.
 - b. $\int_{-2}^3 f(x) dx =$
3. Give the average value of $f(x) = x^2 - 2x + 4$ on the interval $[-1, 2]$, and verify the conclusion of the mean value theorem for integrals for this function on this interval.
 4. Sketch the region bounded between the graphs of $f(x) = 3 - x^2$ and $g(x) = 2x$. Then find the area of the region.
 5. Find the **area** bounded by the graph of $f(x) = x^3 - x^2$ and the x -axis on the interval $[0, 2]$.
 6. Sketch the region bounded by the curves $x + y = 3$ and $x = y^2 + y$. Then give a formula for the area of the region involving integral(s) in x . Repeat the process with integral(s) in y . Finally, find the area of the region.

7. Sketch the region bounded between $f(x) = 2x + 3$ and $g(x) = x^2$. Rotate this region around the y -axis to generate a solid, and then find the volume of the solid.
8. Sketch the region **in the first quadrant** bounded between $f(x) = 2x + 3$ and $g(x) = x^2$. Rotate this region around the y -axis to generate a solid, and then find the volume of the solid.
9. Revolve the region bounded by the line $y = 4$ and the graph of $f(x) = x^2$ about the x -axis to generate a solid. Find the volume.
10. The region bounded between the graphs of $f(x) = x^3 - x^2$ and $g(x) = 2x$ on the interval $[0, 2]$ is rotated around the y -axis to generate a solid. Find the volume.
11. Sketch the region in the first quadrant bounded between the graphs of $f(x) = x + 3$ and $g(x) = (x + 1)^2$. Rotate this region around the y -axis to generate a solid.
 - a. Give a formula involving integral(s) in y for the volume generated.
 - b. Give a formula involving integral(s) in x for the volume generated.
 - c. Give the volume of the solid.
12. Repeat the previous problem, assuming that the region is rotated around the x -axis to generate a volume.
13. The base of a solid is given by the bounded between $f(x) = 2x + 3$ and $g(x) = x^2$. Cross sections taken perpendicular to the x -axis are squares. What is the volume of the solid.
14. The intersection of a solid with the xy plane is the region bounded between $f(x) = 2x + 3$ and $g(x) = x^2$. Cross section taken perpendicular to the x -axis are circles whose diagonals are contained in the xy plane. What is the volume of the solid.