TI Year 4 Metrics with Methodology

Updated: 2/19/2020

| Measurement Stewards Key |                  |                     |               |
|--------------------------|------------------|---------------------|---------------|
| CMS Core Set PEDS Adult  | CMS<br>ScoreCard | Statewide<br>(STCs) | NCQA<br>HEDIS |
| •                        |                  |                     |               |

| Pediatric PCP   |   |  |  |
|---|---|--|--|
| Chosen Performance Measure  | Measure Description   |  |  |
| Well child visits in third, fourth, fifth and sixth years of life | Percentage of children ages 3 to 6 who had one or more well-child visits with a primary care practitioner (PCP) during the measurement year.  |  |  |
| Adolescent well-care visits                                       | Percentage of adolescents ages 12 to 21 who had at least one comprehensive well-care visit with a primary care practitioner (PCP) or an obstetric/gynecologic (OB/GYN) practitioner during the measurement year.  |  |  |
| Well-child visits in the first 15 months of life                  | Percentage of children who turned 15 months old during the measurement year and who had the following number of well-child visits with a primary care practitioner (PCP) during their first 15 months of life: No well-child visitsFour well-child visitsFive well-child visitsSix or more well-child visitsThree well-child visits |  |  |

| Pediatric BH   |  |                         |
|--|--|-------------------------|
| Chosen Performance Measure   | Measure Description  | Measurement<br>Stewards |
| Pediatric follow-up after hosp. for mental illness<br>ages 6-17 (30 - Day) | Percentage of discharges for children ages 6 to 17 who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the child received follow-up within 30 days after discharge | •••                     |
| Pediatric follow-up after hosp. for mental illness<br>ages 6-17 (7 - Day)  | Percentage of discharges for children ages 6 to 17 who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the child received follow-up within 7 days after discharge  | •••                     |
| Metabolic monitoring for children and adolescents on antipsychotics        | Assesses the percentage of children and adolescents with ongoing antipsychotic medication use who had metabolic testing during the year.   | •                       |

| Adult PCP & BH   |   |     |  |
|--|---|-----|--|
| Chosen Performance Measure   | Measure Description   |     |  |
| Follow up after hospitalization for Mental Illness: 18 and older (30 - Day)                                  | Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 30 days after discharge | ••• |  |
| Follow up after hospitalization for Mental Illness: 18 and older (7 - Day)                                   | Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 7 days after discharge  | ••• |  |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are using antipsychotic medications | Percentage of beneficiaries ages 18 to 64 with schizophrenia, schizoaffective disorder, or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.   |     |  |

| Justice  |   |                         |
|--|---|-------------------------|
| Chosen Performance Measure   | Measure Description   | Measurement<br>Stewards |
| Initiation of Alcohol and Other Drug Abuse or Dependence<br>Treatment (14 - Day)                             | Percentage of beneficiaries age 18 and older with a new episode of alcohol or other drug (AOD) abuse or dependence who received initiation of AOD Treatment. Percentage of beneficiaries who initiated treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization, telehealth, or medication treatment within 14 days of the diagnosis |                         |
| Engagement of Alcohol and Other Drug Abuse or Dependence Treatment (34 - Day)                                | Percentage of beneficiaries age 18 and older with a new episode of alcohol or other drug (AOD) abuse or dependence who received continued engagement of AOD Treatment. Percentage of beneficiaries who initiated treatment and who had two or more additional AOD services or medication treatment within 34 days of the initiation visit   | •••                     |
| Follow up after hospitalization for Mental Illness:<br>18 and older (30 - Day)                               | Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 30 days after discharge   | •••                     |
| Follow up after hospitalization for Mental Illness:<br>18 and older (7 - Day)                                | Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 7 days after discharge  | •••                     |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are using antipsychotic medications | Percentage of beneficiaries ages 18 to 64 with schizophrenia, schizoaffective disorder, or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.   | ••                      |

| Hospital   |   |                         |
|--|---|-------------------------|
| Chosen Performance Measure   | Measure Description   | Measurement<br>Stewards |
| Follow up after hospitalization for Mental Illness:<br>18 and older (30 - Day) | Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitionerPercentage of discharges for which the beneficiary received follow-up within 30 days after discharge | •••                     |
| Follow up after hospitalization for Mental Illness:<br>18 and older (7 - Day)  | Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitionerPercentage of discharges for which the beneficiary received follow-up within 7 days after discharge  | •••                     |