



Tropical Journal of Natural Product Research

Available online at <https://www.tjnpr.org>

Original Research Article



Phytochemical Profiling, GC-MS Analysis, and Antioxidant Activity of Polar Solvent Extracts of *Conoclinium coelestinum* Leaves

Othuke B. Odeghe^{1*}, Divine A. Onobrudu¹, Osuvwe C. Orororo¹, Success Ikoya¹, Jacinta E. Apitikori-Owumi², Egoamaka O. Egbune³, Bartholomew C. Nwogweze⁴, Evelyn T. Ojugbeli¹, Oluwaseun E. Agboola⁵, Ejiro P. Awhin¹, Lisa I. Ekakitie¹, Michael O. Ovir⁶, Ifeanyi H. Ofoke⁷, Cyril C. Dunkwu¹, Kingsley C. Enyi¹, Vanessa A. Derhie¹, Precious O. Isaac¹, Joseph C. Mordi¹ and Adebayo O. Adegoke⁸

¹Department of Medical Biochemistry, Faculty of Basic Medical Sciences, Delta state University, P.M.B. 1, Abraka, Nigeria

²Department of Pharmacognosy and Traditional Medicine, Faculty of Pharmacy, Delta state University, P.M.B. 1, Abraka, Nigeria

³Department of Biochemistry, Faculty of Science, Delta state University, P.M.B. 1, Abraka, Nigeria

⁴Department of Physiology, Faculty of Basic Medical Sciences, Delta state University, P.M.B. 1, Abraka, Nigeria

⁵Department of Biochemistry, Faculty of Science, Afe Babalola University, Ado-Ekiti, Ekiti State, Nigeria

⁶Department of Biochemistry, Faculty of Science, Michael and Cecilia Ibru University, Delta State, Nigeria

⁷Department of Biochemistry, Faculty of Science, Madonna University, Elele, Rivers State, Nigeria

⁸Department of Medical Laboratory Science, Faculty of Health Sciences, Madonna University, Elele, Rivers State, Nigeria

ARTICLE INFO

Article history:

Received 21 February 2025

Revised 18 May 2025

Accepted 05 June 2025

Published online 01 July 2025

ABSTRACT

The importance of medicinal plants as source of potent drugs for therapeutic purposes is currently of global interest to researchers. Phytochemicals are compounds from plants used for prevention and treatment of health disorders. Antioxidants are molecules that inhibit free radicals. This research aimed to investigate the phytochemical profile, and antioxidant activity of polar solvent extracts of *Conoclinium coelestinum* leaves. The pulverized leaves was extracted with methanol, ethanol, and water using Soxhlet apparatus. Phytochemical screening of the extracts was done using chemical methods and by gas chromatography-mass spectrometry (GC-MS). Antioxidant activity was evaluated using DPPH, FRAP, ABTS, lipid peroxidation, H₂O₂, nitric oxide, and superoxide radical scavenging assays. Phytochemical screening revealed the presence of alkaloids, glycosides, saponins, flavonoids, terpenoids, and steroids in all three solvent extracts. GC-MS analysis identified twenty-seven (27) bioactive compounds, with hexadecanoic acid ethyl ester (8.89%), 2(3H)-furanone, dihydro-5-pentyl (6.52%), and butanal oxime (5.18%) as the major compounds. Of all the three extracts, the methanol extract significantly ($p < 0.05$) demonstrated exceptional antioxidant activity, achieving 94.15% DPPH inhibition at 400 µg/mL ($IC_{50} = 95.65$ µg/mL) and FRAP activity of 81.12% ($IC_{50} = 100.90$ µg/mL). The methanol extract also demonstrated significant ($P < 0.05$) lipid peroxidation inhibitory activity, H₂O₂, superoxide, and nitric oxide scavenging activities, with percentage inhibition of 64.07%, 76.12%, 90.12%, and 60.07%, respectively at 400 µg/mL, highlighting its capacity to neutralize reactive oxygen species and protect against oxidative damage. These findings affirm *C. coelestinum* as a potential source of natural antioxidants, supported by its rich bioactive compounds profile.

Keywords: *Conoclinium coelestinum*, Gas Chromatography-Mass Spectrometry, Phytochemicals, Antioxidant.

Introduction

Phytochemicals are chemical substances produced by plants, and they have the ability to combat fungal, bacterial, and viral infections in plants. Some foods like beans, fruits and vegetables possess phytochemicals that can help prevent the onset of diseases.^{1,2} They possess antioxidant properties, and aid in eliminating harmful free

radicals.^{3,4} Antioxidants are substances that can remove free radicals from body cells and prevents radical-induced oxidative damage. They are present in numerous plant-derived foods such as spinach, arugula, kale, etc. A variety of medicinal plants have been thoroughly assessed for their antioxidant properties lately.⁵ Antioxidants derived from medicinal and aromatic plants are used to create natural antioxidant formulations for food, health, and cosmetic purposes.⁶ Phytochemical screening is a crucial technique used experimentally to assess the antioxidant compounds present in plants.⁷ *Conoclinium coelestinum*, belonging to the Asteraceae family, is popularly referred to as "Blue Mistflower" and is a herbaceous flowering plant found in North America. *Conoclinium coelestinum* grows in damp soils and along the edges of woods. It acts as a beacon for butterflies, transforming gardens into centers of activity. In conventional medicine, its blossoms and roots are used to treat and manage conditions like fever and abdominal pain.⁸ The plant can be found in the eastern, central, and Midwest regions of the United States, where it has been traditionally utilized for the management of asthma, treatment of fever, cough, sore throats, skin infections, and arrest bleeding from injuries.^{9,10} The volatile substances found in the aerial parts of *C. coelestinum* consist of germacrene D, (2E)-hexenal, β -caryophyllene, and (Z, Z)- α -farnesene. On the other

*Corresponding author. E mail: odegheothukebensandy@delsu.edu.ng
Tel: +2348066768804

Citation: Odeghe OB*, Onobrudu DA, Orororo OC, Ikoya S, Apitikori-Owumi JE, Egbune EO, Nwogweze BC, Ojugbeli ET, Agboola OE, Awhin EP, Ekakitie LI, Ovir MO, Ofoke IH, Dunkwu CC, Enyi KC, Derhie VA, Isaac PO, Mordi JC and Adegoke AO. Phytochemical Profiling, GC-MS Analysis, and Antioxidant Activity of Polar Solvent Extracts of *Conoclinium coelestinum* Leaves. Trop. J. Nat. Prod. Res., 2025 9(6): 2784 - 2794 <https://doi.org/10.26538/tjnpr/v9i6.58>

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria

hand, the non-volatile substances include intermedine, gardenin A, nobiletin, amorphane sesquiterpenoid 5, 8-epoxy-4, 6-dihydroxy-3-cadinanone, 7-hydroxytoxol 7-O-glucoside, luteolin, and its derivatives.¹⁰ The increase in awareness of the relevance of medicinal plants use and the growing request for detailed information regarding the potency and side effects of plant extract employed traditionally in the treatment of diseases necessitate this study. Currently, there is no documented information regarding the phytochemical profile, GC-MS analysis and *in vitro* antioxidant potentials of *C. coelestinum* polar leaves, hence this study was designed to determine and provide the basic scientific knowledge relating to this gap.

Materials and Methods

Collection, identification and preparation of plant material

Fresh leaves of *C. coelestinum* was collected from Abraka Community at GPS coordinates 5°47' 24.828" N and 6° 6' 17.208 E in Ethiopia East Local Government Area, Delta State, Nigeria in June, 2024. Taxonomic identification and authentication of the plant was done by Mr. Wilson O. Egboduku of the Department of Botany, Faculty of Science, Delta State University, Abraka, Nigeria. Herbarium specimen with voucher number DELSUN 269 was deposited at the University Herbarium in the Department of Botany, Faculty of Science. The leaves were washed and shade dried at room temperature in a clean and hygienic conditions. The dried leaves (500 g) were ground into powder and extracted separately with methanol (MeOH), ethanol (EtOH) and water (H₂O) using Soxhlet apparatus based on the modified method described by Hussien and Endalew (2023).¹¹ The resulting plant extracts were kept at 4°C until further experiment.

Phytochemical screening

Qualitative assay

The Qualitative phytochemical screening of the leaf extracts was done based on previously described methods.^{1,11,12}

Quantitative Assay by GC-MS

The GC-MS Analysis was done according to the method previously described by¹³

In vitro antioxidant assays

DPPH radical scavenging assay

The antioxidant potential of *C. coelestinum* leaf extracts was determined using 2,2-diphenyl-1-picryl hydrazyl (DPPH) radical scavenging assay.¹⁴ Different dilutions of the extracts (25, 50, 100, 200 and 400 µg/mL) were prepared, and 2 mL each of the different concentrations of the extract was added to 3 mL of 0.1 mM DPPH solution in methanol. The reaction mixture was incubated at room temperature in the dark for 30 minutes. The absorbance of the mixture was measured at 517 nm using a UV-Visible spectrophotometer. Ascorbic acid at similar concentrations as the extracts was used as the reference standard. Methanol, ethanol, and distilled water were used as the blank for the respective extracts. All determinations were performed in triplicate. The percentage DPPH radical scavenging activity was calculated using the formula below:

$$\% \text{ Inhibition of DPPH} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100$$

Where, A_{control} is the absorbance of the mixture of methanol/ethanol/water and DPPH solution, and A_{sample} is the absorbance of the mixture of sample extract and DPPH solution. The percentage inhibition was plotted against concentration from which IC₅₀ values were calculated.

Ferric reducing antioxidant power (FRAP) assay

The FRAP of the different extracts of *C. coelestinum* leaves was evaluated using the modified method of¹⁵ The extracts solution (100 µL aliquots) was added to 2.5 mL of phosphate buffer (0.2 M, pH 6.6) and 2.5 mL of potassium ferricyanide (1%). After 20 minutes of incubation at 50°C, 2.5 mL of trichloroacetic acid (10%) was added. From the mixture, 2.5 mL was taken and mixed with 2.5 mL of water

and 0.5 mL of 1% FeCl₃. The resulting solution was allowed to stand for 30 minutes, then the absorbance was measured at 593 nm. Trolox was used as the reference standard. The results were expressed as Trolox equivalent antioxidant capacity (TEAC) in mM/L. TEAC was calculated by preparing a Trolox standard curve (Equation: $y = 0.0007x + 0.0645$, $R^2 = 0.9998$) using 31.25 µg/mL to 1.0 mg/mL of standard Trolox solution. Each experiment was carried out in triplicate.

ABTS radical scavenging assay

The ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) radical scavenging activity of the different extracts of *C. coelestinum* leaves was assessed using a modified method of¹⁶ The ABTS radical cation (ABTS^{•+}) was produced after dissolving ABTS in water to a final concentration of 7 mM by reacting ABTS stock solution with 2.45 mM potassium persulfate. The ABTS^{•+} solution was diluted with water until an absorbance of 0.70 (±0.02) at 734 nm was achieved. The reaction mixture comprised 0.07 mL of extract and 3 mL of the ABTS radical solution. The reaction mixture was incubated for 6 minutes, and immediately the absorbance was measured at 734 nm. The ABTS radical scavenging activity was calculated using the formula below:

$$\% \text{ ABTS Inhibition} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100$$

Where; A_{control} is the absorbance of a mixtures of 10 mL of 7 mM ABTS, 2.45 mM K₂S₂O₈ with blank solvents, and A_{sample} is the absorbance of the mixture of sample extract/standard and ABTS. The scavenging activity of *C. coelestinum* leaf extract against ABTS was expressed as IC₅₀.

H₂O₂ radical scavenging assay

The H₂O₂ radical scavenging activity was determined by the method of¹ A potassium phosphate buffer was prepared by mixing 0.2 M potassium dihydrogen phosphate and 0.2 M sodium hydroxide solutions. Equal volume of hydrogen peroxide and potassium phosphate buffer were mixed to generate free radicals, and the solution was kept at room temperature for 5 minutes for the reaction process to be completed. The hydrogen peroxide radical solution (0.6 mL) was added to the 1 mL of the extract, and the absorbance was measured using a spectrophotometer at 230 nm against phosphate buffer solution blank. The percentage inhibition of H₂O₂ was calculated using the formula:

$$\text{Percentage H}_2\text{O}_2 \text{ Inhibition} = \frac{A_1 - A_2}{A_1} \times 100$$

Where A₁ is the control absorbance and A₂ is the absorbance in the presence of the extract and standard

Lipid peroxidation inhibition assay

The ability of *C. coelestinum* leaf extracts to inhibit lipid peroxidation was evaluated *in vitro* following the method described by¹⁷ Briefly, the reaction mixtures contained 2.0 mL of the trichloroacetic acid-thiobarbituric acid-hydrochloric acid (TCA-TBA-HCl) reagent (15% [w/v] TCA, 0.375% [w/v] TBA, and 0.25 N HCl) and 1 mL of the extracts or standard (L-ascorbic acid) at concentrations of 25 µg/mL, 50 µg/mL, 100 µg/mL, 200 µg/mL and 400 µg/mL. The resulting mixtures were incubated in a water bath set at 90°C for 10 minutes, cooled, and centrifuged at 10 000 rpm for 15 minutes. The absorbance of the supernatant was measured at 532 nm. The percentage inhibition of lipid peroxidation (LP) was calculated using the formula below:

$$\% \text{ LP Inhibition} = \frac{\text{Absorbance of control} - \text{Absorbance of test sample}}{\text{Absorbance of control}} \times 100$$

Nitric oxide radical scavenging assay

The ability of the extracts to scavenge nitric oxide (NO) generated from sodium nitroprusside (SNP) was evaluated using the modified method of¹² The reaction mixture (SNP, with or without extract) was incubated close to visible polychromatic light source for 180 minutes at 25°C. The produced NO radical was interacted with oxygen and the resulting nitrite ion (NO₂⁻) was investigated at 30 min intervals by mixing equal

amount of Griess reagent (1% sulphanilamide in 5% phosphoric acid and 0.1% naphthylethylenediamine dihydrochloride). The mixture was incubated, the absorbance of the purple azo dye produced during the diazotization of nitrite ions with sulphanilamide and subsequent coupling with naphthylethylenediamine dihydrochloride was determined at 546 nm. The nitrite formed with or without the leaf extract was calculated based on sodium nitrite solutions of known concentrations using a standard curve.

Superoxide radical scavenging assay

The scavenging of superoxide ($O_2^{\cdot-}$) anion radical was determined by the modified method described by¹⁸ The reaction mixture consisted of 1 mL of alkaline DMSO, 0.3 mL of the leaf extract and 0.1 mL of Nitro blue tetrazolium (NBT). The absorbance of the reaction mixture was measured at 560 nm using a UV-VIS spectrophotometer. Pure DMSO was used as the blank, while ascorbic acid was used as the standard.

Statistical analysis

Data were analyzed using statistical package for social sciences (SPSS) for windows version 23.0. The data were expressed as mean \pm standard error of mean (Mean \pm SEM, $n = 3$). Significant differences between means were determined by one-way analysis of variance (ANOVA) followed by post hoc analysis. P-value < 0.05 was considered statistically significant.

Results and Discussion

Percentage yield of the extracts

The crude extract yields from 500 g of *C. coelestinum* leaves were 69 g for methanol extract, 58 g for ethanol extract, and 35 g for aqueous extract. These results indicate that methanol produced the highest yield compared to the other solvents (Table 1).

Table 1: Extraction yields of *Conoclinium coelestinum* leaves

Sample	Weight (g)	Percentage yield (%)
Methanol Extract	69	42.59
Ethanol Extract	58	35.80
Aqueous Extract	35	21.60
Total Yield	162	100

Phytochemical constituents

This study examined the phytochemical composition, chemical profile, and antioxidant activity of the aqueous, methanol and ethanol extracts of *C. coelestinum* leaves. The purpose of this analysis was to provide scientific validation for the plant's traditional medicinal applications, particularly its antioxidant potentials. A preliminary qualitative phytochemical screening of the leaf extracts revealed the presence of alkaloids, glycosides, saponins, proteins, flavonoids, triterpenoids, steroids, and tannins (Table 2). These compounds are extensively documented for their ability to counteract oxidative stress, a condition associated with neurodegenerative disorders and inflammation.^{19,20} The GC-MS analysis offered a more detailed identification of these phytochemicals, emphasizing their structural and functional diversity, which plays a crucial role in the observed antioxidant properties.²¹

Gas chromatography-mass spectrometric profile of *C. coelestinum*

The GC-MS analysis identified 27 bioactive constituents from the methanol extract of *C. coelestinum* leaves. The chromatogram is presented in Figure 1, while the bioactive constituents with their structures, molecular formula, molecular weight (MW), peak area (%), and retention time (RT) are presented in Table 3.

The chemical compounds identified are Hexadecanoic acid ethyl ester, Pyrrole, 2-methyl-5-phenyl, Butanal, Oxime, Lidocaine, 1,2,4-Trithiolane,3,5-diethyl, Carpaine, Cedrene, 1,2,4-Thiadiazole-3,5-diamine, gamma-Murolene, 2(3H)-Furanone,dihydro-5-pentyl, Alpha-Terpineol, IsoCaryophyllene, trans-Sesquisabinene hydrate,

Table 2: Phytochemical constituents of *Conoclinium coelestinum* leaves

Bioactive Compound	Observation	Inference		
		MeOH	EtOH	H ₂ O
		Extract	Extract	Extract
Alkaloids	Creamy precipitate	+	+	+
Glycosides	Reddish brown precipitate	+	+	+
Saponins	Persistent foaming	+	+	+
Proteins	White precipitate	-	+	+
Flavonoids	Yellow colouration	+	+	+
Terpenoids	Reddish brown	+	+	+
Steroids	Deep yellow	+	+	+
Tannins		+	-	+

Key: - Absent, + Positive.

Methacrolein, n-Caprylic acid, 2(5H)-Furanone,methyl, D-Limonene, 2-Furancarboxaldehyde, 2-Propanone,1-hydroxy, Methyl,8-Methyl-nonanoate, 8-Octadecenoic acid,methyl ester, 1-Butanol,3-methyl, Phenol,3-pentadecyl, Linolenic acid, Pterin-6-carboxylic acid, Tetraacetyl-d-xylonic nitrile, and Propanoic acid.

Among these, hexadecanoic acid ethyl ester exhibited the highest peak area, indicating its prominence as a major compound. Previous studies have reported this compound to possess antioxidant, anti-inflammatory, and hypocholesterolemic activities.¹⁵ Similarly, linolenic acid, a well-known polyunsaturated fatty acid, has demonstrated roles in mitigating oxidative stress and promoting cardiovascular health.¹³

Linolenic acid, with a peak area of 3.47%, is a polyunsaturated fatty acid critical for reducing inflammation and protecting against cardiovascular diseases. Its role in mitigating oxidative damage is well-documented, making it a valuable compound in preventing chronic illnesses.¹⁶ Also, gamma-murolene (3.36%), a sesquiterpene, has demonstrated antimicrobial and anti-inflammatory activities, supporting its potential application in managing infections and inflammatory disorders.²¹

Terpenoids, such as gamma-murolene and alpha-terpineol, identified in moderate quantities, are recognized for their anti-inflammatory and antimicrobial activities. These findings are consistent with the plant's traditional use in managing inflammation-related conditions. Flavonoids like D-limonene were also detected, widely acknowledged for their strong antioxidant and anticancer properties.²²

Other notable compounds include alpha-terpineol (3.63%), a monoterpene alcohol with significant antioxidant and anti-inflammatory properties, and phenol, 3-pentadecyl (1.59%), recognized for its robust free radical scavenging capacity. The presence of D-limonene (1.78%), a monoterpene, adds to the extract's anticancer and antioxidant potential, as reported in various studies.¹⁶

Compounds like 2-furancarboxaldehyde (0.62%) and 2-propanone, 1-hydroxy (3.94%) are known for their antioxidant activity and contribute to the extract's ability to neutralize reactive oxygen species.

























Figure 1: GC-MS Chromatogram of methanol extract of *Conoclinium coelestinum* leaves

Table 3: Compounds identified from the GC-MS analysis of *Conoclinium coelestinum* leaves

S/No.	Name of Compound	Molecular Formula	Molecular Weight (g/Mol)	Peak Area (%)	RT (Min)	Structure of Compound
1.	Hexadecanoic acid ethyl ester	C ₁₈ H ₃₆ O ₂	284.4	8.89	8.839	
2.	Pyrrole,2-methyl-5-phenyl	C ₁₁ H ₁₁ N	157.21	4.74	10.702	
3.	Butanal oxime	C ₄ H ₉ NO	87.12	5.18	10.948	
4.	Lidocaine	C ₁₄ H ₂₂ N ₂ O	234.34	3.26	12.313	
5..	1,2,4-Trithiolane,3,5-diethyl	C ₆ H ₁₂ S ₃	180.4	2.31	12.672	
6.	Carpaine	C ₂₈ H ₅₀ N ₂ O ₄	478.7	2.64	13.336	

7.	Cedrene	C ₁₅ H ₂₄	204.35	3.94	13.774	
8.	1,2,4-Thiadiazole-3,5-diamine	C ₂ H ₄ N ₄ S	116.15	4.21	14.552	
9.	Gamma-Muurolene	C ₁₅ H ₂₄	204.35	3.36	14.815	
10.	2(3 <i>H</i>)-Furanone, dihydro-5-pentyl	C₉H₁₆O₂	156.22	6.52	15.732	
11.	Alpha-Terpineol	C ₁₀ H ₁₈ O	154.25	3.63	17.718	
12.	Isocaryophyllene	C ₁₅ H ₂₄	204.35	4.92	18.473	
13.	trans-Sesquisabinene hydrate	C ₁₅ H ₂₆ O	222.37	1.74	18.890	
14.	Methacrolein	C ₄ H ₆ O	70.09	0.85	20.281	
15.	n-Caprylic acid	C ₈ H ₁₆ O ₂	144.21	3.01	20.504	
16.	2(5 <i>H</i>)-Furanone, methyl	C ₅ H ₆ O ₂	98.1	2.63	21.718	
17.	D-Limonene	C ₁₀ H ₁₆	136.23	1.78	22.295	
18.	2-Furancarboxaldehyde	C ₅ H ₄ O ₂	96.08	0.62	22.849	

19.	2-Propanone,1-hydroxy	C ₃ H ₆ O ₂	74.07	3.94	23.553	
20.	Methyl,8-Methyl-nonanoate	C ₁₁ H ₂₂ O ₂	186.29	2.51	23.791	
21.	8-Octadecenoic acid methyl ester	C ₁₉ H ₃₆ O ₂	296.51	1.79	24.342	
22.	1-Butanol,3-methyl	C ₅ H ₁₂ O	88.15	1.63	24.860	
23.	Phenol,3-pentadecyl	C ₂₁ H ₃₆ O	304.5	1.59	26.392	
24.	Linolenic acid	C ₁₈ H ₃₀ O ₂	278.43	3.47	26.774	
25.	Pterin-6-carboxylic acid	C ₇ H ₅ N ₅ O ₃	207.15	2.22	26.912	
26.	Tetraacetyl-D-xylonic nitrile	C ₁₄ H ₁₇ NO ₉	343.286	2.69	28.374	
27.	Propanoic acid	C ₃ H ₆ O ₂	74.08	1.24	28.726	

Additionally, terpenoids such as gamma-murolene and trans-sesquibabinene hydrate (1.74%) enhance the plant's pharmacological properties, particularly in antimicrobial and anti-inflammatory applications.⁹ Unique compounds like carpaine (2.64%), a plant alkaloid, have demonstrated cardioprotective and antimicrobial effects, while methyl linoleate (1.79%) is a known anti-inflammatory agent. The identification of beta-caryophyllene, a sesquiterpene, further corroborates the extract's potential for managing oxidative stress and inflammatory conditions.

In vitro antioxidant activity of *C. coelestinum* leave extracts

The *in vitro* antioxidant activity of *C. coelestinum* leaf extracts are shown in Tables 4 – 10. Antioxidants are chemicals capable of inhibiting or preventing the oxidation of lipids or other molecules by halting the initiation of oxidative chain reactions. Consequently, they can prevent or reverse damage caused by oxidative stress to the body cells.²³ Natural antioxidants are increasingly favored for their health-promoting properties and their role in disease prevention.²³ The significance of bioactive compounds in traditional medicinal plants cannot be overemphasized owing to their multifaceted therapeutic applications. Studies suggest that plants containing a diverse array of

secondary metabolites like those in *C. coelestinum* may exhibit enhanced pharmacological effects due to their synergistic and multifunctional activities. This aligns with the growing interest in utilizing plant-based extracts as natural remedies for managing oxidative stress and inflammation-driven disorders.

The antioxidant potential of *C. coelestinum* extracts is corroborated by compounds such as phenol, 3-pentadecyl, and D-limonene. Phenolic compounds are well-recognized for their strong radical scavenging abilities, as demonstrated in assays like ABTS (70.15% inhibition at 400 µg/mL) and FRAP (61.67% inhibition at 400 µg/mL). D-limonene, a monoterpene, has been extensively researched for its antioxidant and anti-carcinogenic properties.¹¹

The methanol extract exhibited significant antioxidant activities across various free radical scavenging assays, including DPPH, ABTS, FRAP, lipid peroxidation, hydrogen peroxide (H₂O₂), nitric oxide (NO), and superoxide radical scavenging assays. These findings emphasize the extract's capability to remove reactive oxygen and nitrogen species, which are critical contributors to oxidative stress.

In the DPPH assay, the methanol extract significantly ($p < 0.05$) displayed 94.15% inhibition at 400 µg/mL, with an IC₅₀ value of 95.65 (Table 4).

Table 4: DPPH radical scavenging activity of *Conoclinium coelestinum* leaf extracts

Conc. (µg/mL)	H ₂ O Extract				MeOH Extract				EtOH Extract				AA	
	A	Inhibition (%)	IC ₅₀ (µg/mL)	A	Inhibition (%)	IC ₅₀ (µg/mL)	A	Inhibition (%)	IC ₅₀ (µg/mL)	A	Inhibition (%)	IC ₅₀ (µg/mL)	A	Inhibition (%)
25	0.81 ± 0.09 [@]	70.15	120.87	0.22 ± 0.03 [@]	92.15	95.65	0.24 ± 0.04 [@]	91.85	95.97	0.12 ± 0.06	93.15	92.90		
50	0.78 ± 0.08 [@]	77.18		0.20 ± 0.03 [@]	92.42		0.22 ± 0.03 [@]	92.16		0.10 ± 0.05	93.56			
100	0.76 ± 0.07 [@]	84.13		0.19 ± 0.02 [*]	93.15		0.20 ± 0.02 [@]	92.47		0.09 ± 0.04	93.99			
200	0.72 ± 0.07 [@]	89.19		0.16 ± 0.01 [@]	93.86		0.19 ± 0.02 [@]	92.96		0.08 ± 0.03	94.52			
400	0.68 ± 0.05 [*]	94.45		0.11 ± 0.04 [*]	94.15		0.15 ± 0.01 [*]	93.45		0.08 ± 0.03	95.01			

Data represent mean ± S.E.M (n = 3). Values with superscript (*) denote significant difference (P < 0.05), while values with superscript (@) indicate no significant difference (P > 0.05). A = Absorbance, AA = Ascorbic Acid, IC₅₀ = Half-maximal inhibitory concentration.

DPPH radical has a deep violet colour in solution, turning colourless or pale yellow upon neutralization and conversion to DPPH-H.^{11,18} Numerous plant extracts have been reported to scavenge DPPH radicals in vitro.²⁴ Similarly, extracts of *C. coelestinum* scavenged DPPH radicals in a concentration-dependent manner. Comparable effects have been observed in various tea extracts rich in polyphenols, which also exhibit potent DPPH radical scavenging activity.¹⁸ Other phytochemicals such as mangiferin and naringin have been noted to scavenge DPPH radicals effectively in a concentration-dependent

manner.^{18,25} The DPPH scavenging ability of *C. coelestinum* can thus be attributed to its abundant flavonoids and polyphenols, as highlighted in this study. The reducing power of the extracts, as measured by the FRAP assay, further demonstrated the superiority of MeOH and EtOH extracts, with significant (p < 0.05) percentage inhibitions of 81.12% and 85.03%, respectively, compared to the aqueous extract (61.67%). Their IC₅₀ values followed a similar pattern, with MeOH (100.90 µg/mL) slightly outperforming EtOH (110.10 µg/mL) (Table 5).

Table 5: Ferric reducing antioxidant power (FRAP) of *Conoclinium coelestinum* leaf extracts

Conc. (µg/mL)	H ₂ O Extract				MeOH Extract				EtOH Extract				AA	
	A	Inhibition (%)	IC ₅₀ (µg/mL)	A	Inhibition (%)	IC ₅₀ (µg/mL)	A	Inhibition (%)	IC ₅₀ (µg/mL)	A	Inhibition (%)	IC ₅₀ (µg/mL)	A	Inhibition (%)
25	0.69 ± 0.07 [*]	60.15	192.77	0.49 ± 0.04 [@]	80.01	100.9	0.52 ± 0.03 [@]	83.73	110.1	0.39 ± 0.04	85.12	99.97		
50	0.68 ± 0.07 [@]	60.45		0.47 ± 0.02 [@]	80.23		0.50 ± 0.02 [@]	84.10		0.37 ± 0.03	85.59			
100	0.65 ± 0.06 [*]	60.93		0.45 ± 0.03 [@]	80.45		0.48 ± 0.03 [@]	84.45		0.35 ± 0.01	85.96			
200	0.62 ± 0.05 [@]	61.24		0.44 ± 0.00 [*]	80.75		0.45 ± 0.00 [*]	84.94		0.32 ± 0.02	86.32			
400	0.59 ± 0.03 [*]	61.67		0.39 ± 0.03 [*]	81.12		0.43 ± 0.01 [*]	85.03		0.29 ± 0.00	87.01			

Data represent mean ± S.E.M (n = 3). Values with superscript (*) denote significant difference (P < 0.05), while values with superscript (@) indicate no significant difference (P > 0.05). A = Absorbance, AA = Ascorbic Acid, IC₅₀ = Half-maximal inhibitory concentration.

This result underscores the extract's capability to reduce ferric (Fe³⁺) ions to ferrous (Fe²⁺) ions, indicative of its reducing power. Such activity is strongly linked to its phenolic and flavonoid contents, which stabilizes reactive oxygen species by donating electrons. Compounds like hexadecanoic acid and beta-tocopherol, identified in the GC-MS analysis, contribute significantly to the extract's reducing power and antioxidant properties. The FRAP assay has been widely used due to its simplicity and reliability in determining antioxidant activity.²⁶ Extracts of *C. coelestinum* showed a concentration-dependent increase in FRAP values, consistent with other plant extracts reported to exhibit high antioxidant activity in vitro.^{26,27} In the ABTS assay, the extracts demonstrated lower scavenging activity compared to DPPH and FRAP

assays. The MeOH extract exhibited the highest inhibition (70.15%), followed by EtOH (55.18%) and H₂O extract (48.15%), while ascorbic acid showed superior activity (90.15%). The higher IC₅₀ values of MeOH (190.97 µg/mL), EtOH (256.89 µg/mL), and H₂O extract (334.93 µg/mL) indicate reduced potency compared to DPPH and FRAP assays (Table 6).

This comparatively lower activity may be attributed to the reduced reactivity of certain compounds toward ABTS radicals. These assays reflect the extract's ability to reduce oxidized radicals and ferric ions, demonstrating its robust reducing potential. Compounds like beta-tocopherol and linolenic acid further support these findings.

Table 6: ABTS scavenging activity of *Conoclinium coelestinum* leaf extracts

Conc. ($\mu\text{g/mL}$)	H ₂ O Extract			MeOH Extract			EtOH Extract			AA		
	A	% Inhibitio n	IC ₅₀ ($\mu\text{g/mL}$)	A	% Inhibiti on	IC ₅₀ ($\mu\text{g/mL}$)	A	% Inhibiti on	IC ₅₀ ($\mu\text{g/mL}$)	A	% Inhibition	IC ₅₀ ($\mu\text{g/mL}$)
25	0.56 ± 0.04 [@]	14.45	353.97	0.43 ± 0.04 [*]	28.45	190.97	0.49 ± 0.02 [@]	21.15	287.97	0.32 ± 0.05	5.15	130.97
50	0.52 ± 0.03 [@]	18.98		0.40 ± 0.03 [@]	43.12		0.46 ± 0.02 [*]	32.06		0.29 ± 0.04	5.02	
100	0.49 ± 0.02 [@]	29.43		0.39 ± 0.03 [*]	58.15		0.43 ± 0.02	41.01		0.26 ± 0.02	7.11	
200	0.45 ± 0.01 [*]	35.89		0.36 ± 0.02 [*]	63.10		0.39 ± 0.01 [*]	48.14		0.24 ± 0.03	9.19	
400	0.39 ± 0.00 [@]	48.15		0.31 ± 0.01 [@]	70.15		0.35 ± 0.01 [@]	55.18		0.20 ± 0.01	0.15	

Data represent mean \pm S.E.M (n = 3). Values with superscript (*) denote significant difference (P < 0.05), while values with superscript (@) indicate no significant difference (P > 0.05). A = Absorbance, AA = Ascorbic Acid, IC₅₀ = Half-maximal inhibitory concentration.

The ABTS radical cation is generated through the reaction between ABTS and potassium persulfate, resulting in a blue cation that absorbs light at 734 nm.²⁸ ABTS is highly reactive toward antioxidants, including phenols, thiols, and vitamin C.²⁷ Extracts of *C. coelestinum* displayed concentration-dependent inhibition of ABTS radicals. A similar effect has been observed in extracts of *Syzygium cumini*, naringin, and mangiferin.¹⁸ Additionally, kaempferol, a flavonoid reported to scavenge ABTS radicals, may also contribute to the antioxidant activity of *C. coelestinum* leaves observed in this study.²⁹ Similarly, the H₂O₂ scavenging activity was significant, with 76.12% inhibition at 400 $\mu\text{g/mL}$ (Table 7). These outcomes align with the roles

of terpenoids and phenolics, which are well-known for their protective effects against oxidative stress. H₂O₂ is an essential substance because of its potential to pass through biological membranes. While H₂O₂ itself is not highly reactive, it can become toxic by generating hydroxyl radicals within cells.³⁰ The scavenging of H₂O₂ by the extracts may be attributed to the presence of phenolics, which donate electrons to neutralize H₂O₂ into water.³¹ The data suggests that all extracts exhibit robust H₂O₂ scavenging activity, likely owing to their antioxidant compounds. As effective electron donors, these components facilitate the conversion of H₂O₂ to H₂O.

Table 7: H₂O₂ scavenging activity of *Conoclinium coelestinum* leaf extracts

Conc. (μg/mL)	H ₂ O Extract			MeOH Extract			EtOH Extract			AA		
		%	IC ₅₀		%	IC ₅₀		%	IC ₅₀		%	IC ₅₀
	A	Inhibition	(μg/mL)	A	Inhibition	(μg/mL)	A	Inhibition	(μg/mL)	A	Inhibition	(μg/mL)
25	0.55 ± 0.03 [@]	61.56	285.77	0.46 ± 0.02 [@]	74.01	196.19	0.49 ± 0.02 [*]	69.73	228.10	0.33 ± 0.04	75.12	184.14
50	0.52 ± 0.02 [*]	61.91		0.43 ± 0.03 [*]	75.23		0.46 ± 0.02 [@]	70.10		0.31 ± 0.03	75.59	
100	0.50 ± 0.01 [@]	62.13		0.41 ± 0.02 [@]	75.45		0.44 ± 0.02 [*]	70.45		0.29 ± 0.03	75.96	
200	0.48 ± 0.03 [*]	62.67		0.38 ± 0.01 [*]	75.75		0.42 ± 0.01 [@]	70.94		0.27 ± 0.01	76.32	
400	0.46 ± 0.01 [*]	63.07		0.36 ± 0.00 [*]	76.12		0.39 ± 0.01 [*]	71.03		0.23 ± 0.02	7.01	

Data represent mean \pm S.E.M (n = 3). Values with superscript (*) denote significant difference (P < 0.05), while values with superscript (@) indicate no significant difference (P > 0.05). A = Absorbance, AA = Ascorbic Acid, IC₅₀ = Half-maximal inhibitory concentration.

The lipid peroxidation assay revealed a 64.07% inhibition at 400 $\mu\text{g/mL}$, demonstrating the extract's capacity to prevent oxidative damage to lipids, which is essential for maintaining cellular integrity (Table 8). The nitric oxide scavenging assay showed 60.07% inhibition at 400 $\mu\text{g/mL}$, indicating the extract's potential in mitigating inflammation mediated by NO radicals (Table 9). Furthermore, the superoxide radical scavenging activity was notably high, with 90.12% inhibition at the same concentration, highlighting the extract's effectiveness in countering mitochondrial oxidative stress (Table 10).

The nitric oxide radical (NO•) is known for its cytotoxicity, particularly when reacting with oxygen or superoxide anion radicals. The *C. coelestinum* extracts demonstrated a concentration-dependent reduction in NO• generation. Also, several plant extracts and formulations have shown the ability to scavenge NO• in a concentration-dependent manner.³² Compounds such as betanin, phylloactin, and betanidin have been previously documented to scavenge NO radicals effectively.³³

Table 8: Lipid peroxidation inhibitory activity of *Conoclinium coelestinum* leaf extracts

Conc. (µg/mL)	H ₂ O Extract			MeOH Extract			EtOH Extract			AA		
	% Inhibition		IC ₅₀ (µg/mL)	% Inhibition		IC ₅₀ (µg/mL)	% Inhibition		IC ₅₀ (µg/mL)	% Inhibition		IC ₅₀ (µg/mL)
	A			A			A			A		
25	0.63 ± 0.02*	62.26	276.28	0.49 ± 0.00@	68.01	186.12	0.55 ± 0.03@	66.73	195.10	0.45 ± 0.03	72.12	175.23
50	0.61 ± 0.03*	62.91		0.48 ± 0.03@	68.13		0.52 ± 0.01@	67.10		0.42 ± 0.04	72.48	
100	0.58 ± 0.02@	63.13		0.46 ± 0.02	68.44		0.50 ± 0.01@	67.45		0.40 ± 0.00	72.85	
200	0.57 ± 0.02@	63.67		0.45 ± 0.00*	68.76		0.48 ± 0.02@	67.74		0.37 ± 0.03	73.32	
400	0.55 ± 0.01*	64.07		0.42 ± 0.00*	69.11		0.47 ± 0.03*	68.00		0.35 ± 0.01	74.00	

Data represent mean ± S.E.M (n = 3). Values with superscript (*) denote significant difference (P < 0.05), while values with superscript (@) indicate no significant difference (P > 0.05). A = Absorbance, AA = Ascorbic Acid, IC₅₀ = Half-maximal inhibitory concentration.

Table 9: Nitric oxide scavenging activity of *Conoclinium coelestinum* leaf extracts

Conc. (µg/mL)	H ₂ O Extract			MeOH Extract			EtOH Extract			AA		
	% Inhibition		IC ₅₀ (µg/mL)	% Inhibition		IC ₅₀ (µg/mL)	% Inhibition		IC ₅₀ (µg/mL)	% Inhibition		IC ₅₀ (µg/mL)
	A			A			A			A		
25	0.75 ± 0.05*	58.36	302.13	0.59 ± 0.02*	62.24	201.12	0.66 ± 0.02*	61.73	235.76	0.52 ± 0.05	67.12	182.95
50	0.74 ± 0.04@	58.91		0.57 ± 0.03*	62.23		0.63 ± 0.00	62.33		0.50 ± 0.03	67.45	
100	0.72 ± 0.03@	59.22		0.54 ± 0.01*	62.45		0.62 ± 0.01*	62.45		0.48 ± 0.02	67.86	
200	0.69 ± 0.01*	59.65		0.53 ± 0.04@	63.75		0.59 ± 0.01@	62.94		0.45 ± 0.00	68.15	
400	0.67 ± 0.02*	60.07		0.49 ± 0.00*	64.12		0.57 ± 0.02*	63.03		0.42 ± 0.02	69.01	

Data represent mean ± S.E.M (n = 3). Values with superscript (*) denote significant difference (P < 0.05), while values with superscript (@) indicate no significant difference (P > 0.05). A = Absorbance, AA = Ascorbic Acid, IC₅₀ = Half-maximal inhibitory concentration.

Table 10: Superoxide radical scavenging activity of *Conoclinium coelestinum* leaf extracts

Conc. (µg/mL)	H ₂ O Extract			MeOH Extract			EtOH Extract			AA		
	% Inhibition		IC ₅₀ (µg/mL)	% Inhibition		IC ₅₀ (µg/mL)	% Inhibition		IC ₅₀ (µg/mL)	% Inhibition		IC ₅₀ (µg/mL)
	A			A			A			A		
25	0.72 ± 0.06@	80.36	139.00	0.49 ± 0.04*	89.01	120.12	0.57 ± 0.03	90.73	105.64	0.40 ± 0.06	92.12	94.87
50	0.71 ± 0.05@	80.91		0.46 ± 0.03@	89.23		0.53 ± 0.06*	91.13		0.39 ± 0.04	92.59	
100	0.69 ± 0.01@	81.13		0.42 ± 0.01*	89.45		0.51 ± 0.01*	91.46		0.37 ± 0.01	92.96	
200	0.67 ± 0.00*	81.67		0.40 ± 0.04*	89.75		0.49 ± 0.03*	91.92		0.33 ± 0.03	92.32	
400	0.42 ± 0.02*	82.07		0.38 ± 0.01*	90.12		0.43 ± 0.00*	92.03		0.29 ± 0.04	4.01	

Data represent mean ± S.E.M (n = 3). Values with superscript (*) denote significant difference (P < 0.05), while values with superscript (@) indicate no significant difference (P > 0.05). A = Absorbance, AA = Ascorbic Acid, IC₅₀ = Half-maximal inhibitory concentration.

Additionally, flavonoids like kaempferol, myricetin, epigallocatechin gallate, catechin, epicatechin, and resveratrol, along with delphinidin, pelargonidin, malvin, mangiferin, and naringin, have been reported to neutralize NO radicals in earlier studies.^{18,28,31,34}

Differences in observations could be attributed to variations in the cultivars used or the influence of the local environment where the plants were grown. For instance, contrary to these findings, it has been reported that polyphenol extracts from mango leaves failed to scavenge hydroxyl radicals across all tested concentrations.³⁵ However, DPPH scavenging of mango leaf alcoholic extracts, as reported by³⁶ aligns with the present observations. These authors emphasized that the type of extraction solvent and plant variety play pivotal roles in determining phenolic content and antioxidant activity.¹⁸ Four primary phenolic compounds identified in mango leaves contributed significantly to antioxidant properties.²⁶

The diverse phytochemicals identified through GC-MS analysis, coupled with the robust antioxidant activity, highlight the pharmacological potential of *C. coelestinum*. These compounds possess anti-inflammatory and cardioprotective activities. For instance, flavonoids are associated with antitumor, anti-inflammatory, and anti-diarrheal properties, while terpenoids exhibit activities such as anti-HIV, antitumor, and antimycobacterial effects.³⁷ These findings reinforce the therapeutic relevance of *C. coelestinum* for managing oxidative stress and associated disorders.

Conclusion

The GC-MS analysis of *C. coelestinum* leaf extracts highlight the richness and diversity of its bioactive compounds. The methanol and ethanol extracts of *C. coelestinum* leaves exhibit potent antioxidant activity, with methanol extract showing slightly superior efficacy. Among all the antioxidant assays, the DPPH and FRAP assays demonstrated the highest scavenging activities, highlighting the extracts' potential to neutralize free radicals and reduce oxidative stress. These findings, in conjunction with the strong radical scavenging activity observed in the assays, validate the plant's traditional use in medicine and underscore its potential as a source of natural antioxidants and anti-inflammatory agents. Future studies should focus on isolating and characterizing the bioactive compounds to better understand their individual and synergistic roles in therapeutic applications.

Conflict of interest

The author reports no conflicts of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

References

- Agrawal MY, Agrawal YP, Arora SK, Lahange P, Kshirsagar N. Phytochemical screening and evaluation of antioxidant activity of hydroalcoholic extract of *Justicia procumbens* leaf. *J Ayurv Herb Med*. 2021; 7(1):41-45.
- Orororo OC and Asagba SO. Effects of *Hibiscus sabdariffa* L. anthocyanins on body weight, lipid peroxidation, and lipid profile of cadmium-exposed rats. *Saudi J Forensic Med Sci*. 2024; 4(1):21-25.
- Candace JC. Harmful free radicals. *Lancet*. 2023; 23:680.
- Odeghe OB, Orororo OC, Agboola OE, Onyeso G, Dennis-Eboh U, Ekakitie LI, Awhin EP, Ojugbeli ET. Phytochemical profiling and *in vitro* antioxidant activity of methanolic extracts from *Fadogia cienkowski* leaves. *Asian Plant Res J*. 2024; 12(6):110-120.
- Onobrudu DA, Nzebude CP, Orororo OC. Comparative analysis of the phytochemicals, minerals, antioxidant, and antimicrobial activities of aqueous and ethanolic extracts of *Rauvolfia vomitoria* leaves. *Asian Plant Res J*. 2024; 12(5):67-81.
- Busari, AA, Obadiha CC, Orororo OC, Efejene IO, Aisuodionoe ME, Osirim E, DeLoache S, Ebu CI, Adejare AA, Badmos K. Assessment of Cardiovascular Function in Periodontitis and Diabetes Mellitus Induced Rats Treated with Black Seed (*Nigella sativa*) Oil. *Asian J Cardiol Res*. 2024; 7(1):230-244.
- Odeghe OB, Adikwu E, Ojiego CC. Phytochemical and antioxidant assessments of *Dioscorea bulbifera* stem tuber. *Biomed Biotechnol Res J*. 2020; 4:305-311.
- Kiersten I. Medicinal and traditional applications of plant-derived compounds. *Herbal Med J*. 2024; 17:89-94.
- Guo X, Guo Z, Duan C, Wang G, Lu Y, Li M, Zhang Z, Chen J, Li J. Long-term persistence of IgG antibodies in SARS-CoV-infected healthcare workers. *medRxiv*. 2020; 02(12):20021386.
- Lawson GM, McKenzie ME, Becker KD, Selby L, Hoover SA. The core components of evidence-based social emotional learning programs. *Prev Sci*. 2019; 20(4):457-467.
- Hussen EM and Endalew SA. *In vitro* antioxidant and free-radical scavenging activities of polar leaf extracts of *Vernonia amygdalina*. *BMC Complement Med Ther*. 2023; 23:146. doi: 10.1186/s12906-023-03923-y.
- Odeghe OB, Monanu MO, Anacletus FC. *In vitro* antioxidant abilities and inhibition of rabbit muscle lactate dehydrogenase by aqueous extract of *Hannoa klaineana* stem bark. *Int J Adv Res*. 2016; 4:509-515.
- Njoku OO, Umeh CG, Ogugofor MO. Phytochemical profiling and GC-MS analysis of aqueous methanol fraction of *Hibiscus asper* leaves. *Futur J Pharm Sci*. 2021; 7(59):1-5.
- Odeghe OB, Orororo CO, Egbune EO, Onobrudu DA, Nwilo BI, Bassey UE, Adegoke AO, Effiong GS, Ovir, MO, Ofoke, IH, Mordi, JC, Ejimole, OL. Phytochemical analysis, antioxidant capability, and inhibition studies of the aqueous extract of *Phyllanthus discoides* root. *Biomed Biotechnol Res J*. 2023; 7(3):411-419. doi: 10.4103/bbrj.bbrj_126_23.
- Shahinuzzaman M, Yaakob Z, Anuar FH, Akhtar PN, Kadir HA, Hasan AKM, Sobayel K, Nour M, Sindi H, Amin N, Sopian K, Akhtaruzzaman MD. *In vitro* antioxidant activity of *Ficus carica* L. latex from 18 different cultivars. *Sci Rep*. 2020; 10(1):10852. doi: 10.1038/s41598-020-67765-1.
- González-Palma I, Escalona-Buendía HB, Ponce-Alquicira E, Téllez-Téllez M, Gupta VK, Díaz-Godínez G, Soriano-Santos J. Evaluation of the antioxidant activity of aqueous and methanol extracts of *Pleurotus ostreatus* in different growth stages. *Front Microbiol*. 2016; 7:1099. doi: 10.3389/fmicb.2016.01099.
- Moriasi GA, Ireri AM, Ngugi MP. *In vivo* cognitive-enhancing, *ex vivo* malondialdehyde-lowering activities and phytochemical profiles of aqueous and methanolic stem bark extracts of *Piliostigma thonningii* (Schum.). *Int J Alzheimer's Dis*. 2020; 2020:1367075. doi: 10.1155/2020/1367075.
- Lalhmingshui K and Jagetia GC. Evaluation of the free-radical scavenging and antioxidant activities of *Chilauni*, *Schima wallichii* Korth *In vitro*. *Future Sci OA*. 2018; 4(2):FSO272. doi: 10.4155/foa-2017-0086.
- Idenyi OM, Orororo OC, Efejene IO, Aisuodionoe ME, Umoren UF. (2024). Evaluation of anti-inflammatory and analgesic effects of *Allophylus spicatus* leaf extract in mice. *J Complement Altern Med Res*. 2024; 25(9):59-68.
- Odeghe OB, Monanu MO, Anacletus FC. Phytochemistry of Achomanes Difformis Stem, Dioscorea Bulbifera Stem Bark, Fadogia Cienkowski Leaf, Hannoa Klaineana Stem and Vitex Simplicifolia Leaf. *AIM Sci Res*. 2016; 2(1):1-8.
- Kumari S, Yadav S, Yadav JP, Srivastava S. GC-MS analysis and evaluation of antioxidant activities of methanolic root extract of *Withania somnifera*. *J Appl Biol Biotechnol*. 2019; 7(4):1-7.
- Osafanne IL and Ngozi OP. Phytochemical screening and *in vitro* antioxidant properties of methanol and aqueous leaf extracts of *Geophila obvallata*. *Asian J Res Biochem*. 2018; 3(2):1-11.
- Neha K, Haider MR, Pathak A, Yar MS. Medicinal prospects of antioxidants: A review. *Eur J Med Chem*. 2019; 178:687-704. doi: 10.1016/j.ejmech.2019.06.010.
- Endris MH and Sisay AE. *In vitro* antioxidant and free radical scavenging activities of polar leaf extracts of *Vernonia amygdalina*. *BMC Complement Med Ther*. 2023; 23(1):146.
- Puangpronpitag D, Tankitjanon P, Sumalee A, Konsue A. Phytochemical screening and antioxidant activities of the seedling extracts from *Inca peanut (Plukenetia volubilis)*. *Phytochem J*.

- 2021;13(1):52–58.
26. Abeysinghe DT, Kumara KAH, Kaushalya KAD, Chandrika UG, Alwis DDDH. Phytochemical screening, total polyphenol, flavonoid content, *in vitro* antioxidant and antibacterial activities of Sri Lankan varieties of *Murraya koenigii* and *Micromelum minutum* leaves. *Heliyon*. 2021; 7(7):e0744.
 27. Minarti M, Ariani N, Megawati M, Hidayat A, Hendra M, Primahana G, Darmawan A. Potential antioxidant activity methods (DPPH, ABTS, FRAP), total phenol, and total flavonoid levels of *Macaranga hypoleuca* leaves extract and fractions. *Web Conf*. 2024; 503:07005.
 28. Hassan A, Akmal Z, Khan N. The phytochemical screening and antioxidant potential of *Schoenoplectus triqueter* L. Palla. *J Chem*. 2020; (1):3865139.
 29. Ahamad R, Singh N, Dixit V, Dudhe A, Dudhe R. Phytochemical analysis and antioxidant activity of whole plant of *Tridax procumbens* Linn. *J Adv Sci Res*. 2023; 14(1):61–68. doi: 10.55218/JASR.202314109.
 30. Adebayo SA, Ondua M, Shai LJ, Lebelo SL. Inhibition of nitric oxide production and free radical scavenging activities of four South African medicinal plants. *J Inflamm Res*. 2019; 12:195–203.
 31. Sharma P, Raj H, Choudhary D, Kumari P, Kaur R, Ankita, Kumar S. Phytochemical screening, *in vitro* antioxidant, and antibacterial activities on *Skimmia laureola* leaves extract. *Eur J Med Plants*. 2023; 34(11):40–55.
 32. Andrés CM, Pérez de la Lastra JM, Juan CA, Plou FJ, Pérez-Lebeña E. Chemistry of hydrogen peroxide formation and elimination in mammalian cells, and its role in various pathologies. *Stresses*. 2022; 2(3):256–274.
 33. Andrabi SM, Ali H, Khan NA, Fatima S, Malik AR. Nitric oxide: physiological functions, delivery, and biomedical applications. *Adv Sci*. 2023; 10(30):1–38.
 34. Rao H, Rao I, Saeed L, Aati HY, Aati S, Zeeshan M, Khan KUR. (2023). Phytochemical analysis and bioactivity assessment of five medicinal plants from Pakistan: Exploring polyphenol contents, antioxidant potential, and antibacterial activities. *Saudi J Biol Sci*. 2023; 30(10):103783.
 35. Ahemad AIA, Aejazuddin QMA, Khan GJ. Phytochemical screening and *in-vitro* free radical scavenging activity of Unani formulation *Habb-e-Asgand*. *J Pharm Res Int*. 2021; 33(59B):1–7.
 36. Ndel Famen LC, Tangué Talom B, Simo Tagné R, Tchente Kamsu G, Kodjio N, Tamekou LS, Gatsing D. *In vitro* antioxidant activities and effect of hydroethanolic and aqueous extracts of *Terminalia avicennioides* on *Salmonella*. *Microbiol Res J Int*. 2020; 30(1):1–14. doi: 10.9734/mrji/2020/v30i130185.
 37. Velmurugan G and Anand S. GC-MS analysis of bioactive compounds in traditional medicinal plants. *Asian J Res Biochem*. 2022; 11(2):1–9.