

Independent Project Diony Final Report

Diony Gamoso

r Sys.Date()

First will load packages

```
library(tidyverse)
```

```
## Warning: package 'ggplot2' was built under R version 4.3.3
```

```
## Warning: package 'stringr' was built under R version 4.3.3
```

```
library(openintro)  
library(ggplot2)  
library(dplyr)
```

Introduction

Eighty-seven percent of the world's flowering plants reproduce through the aid of pollinators, and the most important pollinators are bees (Ollerton et al., 2011). However, many bee species are in decline around the world due to disease, habitat destruction, pesticides, exotic species, urbanization, and other stressors (Biesmeijer et al., 2006; Burkle et al., n.d.; Leather, 2018; LeBuhn & Vargas Luna, 2021; Mathiasson & Rehan, 2020). To counter this decline, many organizations are increasing efforts to preserve and create habitat for wild bees in urban areas. Urban areas can serve as a refuge for wild bees (Frankie et al., 2019; Frankie et al., n.d.; Mach & Potter, 2018; Matteson & Langellotto, 2011; Tonietto et al., 2011). Urban food gardens, residential landscaping, city parks, weedy lots, pollinator gardens, and “ecosystem-oriented” urban landscapes are host to a diverse array of native bees (G. W. Frankie et al., n.d.; Pawelek et al., 2009; Tommasi & Entomologist, n.d.; Wojcik et al., 2008). In urban ecosystems, wild bees pollinate plants, which, in turn, provides food and shelter for other wildlife, and humans (Burghardt et al., 2009; Hausmann et al., 2016; Lowenstein et al., 2015; Narango et al., 2020; Potter & LeBuhn, 2015). Interested individuals, grassroots organizations, city and county parks, and even entire cities are starting to realize the importance of urban habitats for pollinators, and efforts are under way to provide patches of habitat within the urban jungle (Home - Million Pollinator Garden Challenge, n.d.; National Wildlife Federation, n.d.; The Xerces Society for Invertebrate Conservation, n.d.).

Bees can prefer native plants (Morandin & Kremen, 2013; Pardee & Philpott, 2014), but also use ornamental, or non-native plants (Belle, n.d.; G. Frankie et al., 2019; G. W. Frankie et al., 2005; Mach & Potter, 2018; Matteson & Langellotto, 2011; Pawelek et al., 2009)

Few studies have looked at the effectiveness of relatively small-scale, native plant additions in urban environments.

The Presidio, an urban national park within the city of San Francisco, CA U.S.A. , recently began an initiative to practice ecological horticulture (“EcoHort”) in the formal, designed landscape (Michael Boland, Chief Park

Officer, pers. comm). EcoHort aims to convert decorative planters and lawns, as well as under-utilized, weedy plots within the designed landscape, into ecologically richer habitats which are both aesthetically pleasing and ecologically valuable. Large-scale changes are just on the horizon, and hence, a giant experiment is about to unfold. Let us capitalize on this opportunity to study the effects of current, small-scale management actions, fine-tune them for greater effectiveness, use that to inform future planning and management on a larger scale, and share what is learned with others.

Given that the eventual goal of the Presidio Ecological Horticulture project is to develop a more ecologically rich environment within the designed landscape, inhabited by a diverse array of native pollinators, in this study, I will ask: 1. Does supplementing existing planter beds with native wildflowers increase pollinator/bee abundance?

Hypotheses: 1. Null: Planter beds augmented with native wildflowers will not have greater pollinator abundance. 1' Alternative: Planter beds augmented with native wildflowers will have greater pollinator abundance.

Methods for statistical analysis

I used the tidyverse package as well as ggplot2. To prepare the raw data for analysis, I first grouped the Host plant treatment and Non-host plant treatment into one category “Augmented” - meaning both of these treatments were augmented with native wildflowers. Control plots, which did not get augmented were classified as “Unaugmented”. My analysis is focused on comparing pollinator abundance in “Augmented” versus “Unaugmented” plots. I then selected the relevant variables using “group_by” to create a simplified dataframe. From there, I used the “summarise” function to get the sums of pollinator Counts for each individual sampling event which is identified by the Date-Block-Treatment code “DateBlkTrtmnt”. Using this new dataframe with summed counts, I then plotted the data using ggplot2 to create boxplots.

After, I ran an ANOVA to test for significance between the pollinator abundance per individual sampling event in Augmented versus Unaugmented plots. I used an ANOVA to compare the two means, rather than a t-test, because I used a blocked design, and the ANOVA would allow me to factor in the Block as a random effect.

link to data on Github:<https://github.com/PresidioEcoHort/Bee-study/tree/main/Data>

link to R-script on Github: <https://github.com/PresidioEcoHort/Bee-study/tree/main/Rscript>

Now will read in data files

```
# Read in the bee observation data
Bees2023 <- read.csv("C:/Users/dgamoso/OneDrive - The Presidio Trust/Diony's Stuff/Graduate school/Presidio Ecological Horticulture/Bees2023.csv")
# Next, create a new data frame where both Host and Non-host treatments are clumped under "Augmented" (Control is "Unaugmented")
Bees2023Augmented <- Bees2023 %>% dplyr::mutate(Augmentation = case_when(Treatment == "C" ~ "NoExtraFlowers", Treatment == "H" ~ "Augmented", Treatment == "NH" ~ "Augmented"))
```

... # Now, want to group the data by DateBlkTrtmnt, and sum the insect counts per each DBT :

```
# let me just group by DateBlkTrtmnt these 4 categories
DBT_groups<-group_by(Bees2023Augmented,DateBlkTrtmnt, Augmentation, Block)

# Next, I want to get a total count of observed or netted bees per group, our groups being DateBlkTrtmnt, Augmentation, Block
Sums_counts_DBT <-summarize(DBT_groups, SumsOfCounts = sum (Count))
```

```
## 'summarise()' has grouped output by 'DateBlkTrtmnt', 'Augmentation'. You can
## override using the '.groups' argument.
```

```
#getting rid of the NA
rowsnotNA <- which(!is.na(Sums_counts_DBT$Augmentation))
Sums_counts_DBT<-Sums_counts_DBT[rowsnotNA,]

#What is the mean of each treatment?
means_AugVsUnaug <-aggregate(SumsOfCounts~Augmentation, Sums_counts_DBT, mean)
means_AugVsUnaug
```

```
##           Augmentation SumsOfCounts
## 1 NativeFlowersAdded      4.156716
## 2      NoExtraFlowers      1.208333
```

Visualizing the data with a boxplot

```
plot_means_all_pollinators<- ggplot(data = Sums_counts_DBT, aes (x = Augmentation, y = SumsOfCounts, fill = Augmentation)) +
  geom_boxplot()+
  labs(x = "Added Native Wildflowers?", y="Mean/Median Pollinatorcounts", title = "Does adding wildflowers increase pollinator counts?",
  stat_summary(fun.y="mean",color="red", shape=13)
```


```
## Warning: The 'fun.y' argument of 'stat_summary()' is deprecated as of ggplot2 3.3.0.
## i Please use the 'fun' argument instead.
## This warning is displayed once every 8 hours.
## Call 'lifecycle::last_lifecycle_warnings()' to see where this warning was
## generated.
```

```
plot_means_all_pollinators
```

```
## Warning: Removed 2 rows containing non-finite outside the scale range
## ('stat_boxplot()').
```

```
## Warning: Removed 2 rows containing non-finite outside the scale range
## ('stat_summary()').
```

```
## Warning: Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
```



➤ Augmented (plots with native wildflowers added), and Unaugmented (control) plots


```
#If I want to look at all blocks...
plots_all_blocks<-ggplot(data = Sums_counts_DBT, aes (x = Augmentation, y = SumsOfCounts, fill = Augmentation)) +
  geom_boxplot()+
  labs(x = "Added Native Wildflowers?", y="Mean/Median Pollinatorcounts", title = "Does adding wildflowers make a difference?") +
  stat_summary(fun.y="mean",color="red", shape=13)+
  facet_wrap(~Block)
plots_all_blocks
```

```
## Warning: Removed 2 rows containing non-finite outside the scale range
## ('stat_boxplot()').
```

```
## Warning: Removed 2 rows containing non-finite outside the scale range
## ('stat_summary()').
```

```
## Warning: Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
## Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
## Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
## Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
## Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
## Removed 2 rows containing missing values or values outside the scale range
```

```
## ('geom_segment()').
## Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
## Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
## Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
## Removed 2 rows containing missing values or values outside the scale range
## ('geom_segment()').
```



› Augmented (plots with native wildflowers added), and Unaugmented (control) plots

Now want to run an ANOVA, but only comparing two groups (ANOVA rather than t-test because ANOVA allows me to factor in the Block)

```
# ANOVA
anova_augmentation <- aov(SumsOfCounts ~ Augmentation + Block, data= Sums_counts_DBT)
summary(anova_augmentation)
```

```
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Augmentation  1  407.1    407.1  31.376 7.16e-08 ***
## Block         9  421.3     46.8   3.607 0.000352 ***
## Residuals    195 2530.3     13.0
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## 2 observations deleted due to missingness
```

Results

The results of the ANOVA show highly significant effect of “Augmentation” (adding native wildflowers to a planter). The p-value was 0.0000000716, which is highly significant, and therefore we can reject the null hypothesis. In addition, the effect of block was also significant with a p-value of 0.000352.

Discussion

This experiment is being performed to test the effectiveness of augmenting ornamental planter beds with native plants. The Presidio Trust is interested in this question because they are embarking on a new land management initiative, “Ecological Horticulture” to transform the formal, designed landscape into “ecoscapes” using native plants both for their aesthetic value as well as their ecological value. This highly significant result from the data showing that adding native plants to planter beds increases pollinator abundance helps confirm that these Ecological Horticulture practices are fulfilling their desired, ecological goals. However, do note that the design of the experiment does not tease out the effect of adding native wildflowers, from the effect of just adding more flowers in general (regardless of nativity or not). We did not augment planter beds with more non-native, ornamental flowers which would have helped tease out the importance of nativity. More research on that would be desirable in the future. I am glad that I ended up choosing to do a block design. It’s very much like creating a “food court” with the 3 “restaurant” choices offered up to pollinators within each block. Looking at all of the blocks, the Augmented “restaurant” gets more “customers”, although the degree to which the Augmented plot gets more pollinators varies by block. The blocks where Augmented and Unaugmented blocks did not differ very much in mean number of pollinators visiting were typically those blocks in which the native plantings had not grown very much, and had very small blooms.

These preliminary results bode well for future pollinator abundance as these planter beds develop over time. The current data was taken from plots whose native plants were essentially “infantile” - being small in size and having relatively few blooms. As these plantings get more robust, I am expecting more flowers, and likely an even bigger effect on pollinator abundance.