

Addendum: Advanced Modalities in Artificial Oxygen Carriers and Intravenous Carbon Dioxide Mitigation for Extreme Hypoxic Environments

Introduction to the Biochemical Engineering of Survival

The engineering of pressurized vessels, deep-well environments, and closed-loop life support systems traditionally focuses on maintaining a breathable atmosphere through mechanical ventilation, ambient gas scrubbing, and external oxygen supply. However, in catastrophic entrapment scenarios—such as submarine failure, mining collapses, or deep-well entrapment—external mechanical life support apparatuses frequently fail or become entirely inaccessible. In these acute emergencies, the limits of human survivability are dictated by two inexorable biochemical clocks: the rapid depletion of cellular oxygen (hypoxia) and the toxic accumulation of metabolic carbon dioxide (hypercapnia).

When ambient oxygen cannot be restored, or when catastrophic blood loss prevents native erythrocytes from delivering available oxygen to tissues, the paradigm of rescue must shift from environmental engineering to systemic biological engineering. This addendum expands upon existing research regarding pressurized environments by detailing cutting-edge pharmacological and biochemical interventions designed to sustain human life from within the vascular compartment. Specifically, this report synthesizes the latest advancements in Artificial Oxygen Carriers (AOCs)—encompassing advanced Hemoglobin-Based Oxygen Carriers (HBOCs), Perfluorocarbons (PFCs), and novel injectable oxygen gas microparticles—alongside parallel strategies for intravenous and chemical carbon dioxide mitigation. By bridging the gap between point-of-injury and definitive extrication, these technologies possess the potential to drastically extend the survivability window in hypoxic and hypercapnic environments.

The Historical Evolution of Blood Surrogates

The quest for a suitable alternative to human blood dates back to the seventeenth century, when pioneers like Sir Christopher Wren proposed the intravenous infusion of ale, wine, and opium as rudimentary blood substitutes¹. Over the centuries, an assortment of fluids, including milk, plant resins, and sheep blood, were utilized with predictably fatal hemolytic reactions¹. It was not until the early twentieth century, following Karl Landsteiner's discovery of blood groups, that allogeneic blood transfusion became a safe and established medical procedure¹.

However, allogeneic blood possesses severe logistical limitations for emergency entrapment scenarios. Donated human blood requires strict refrigeration (the cold chain), has a maximum

shelf life of approximately 42 days, requires time-consuming type-and-crossmatch compatibility testing, and carries the risk of transmitting blood-borne pathogens¹. In austere environments—such as a collapsed mine or a battlefield—these constraints render whole blood transfusion nearly impossible⁴.

The modern foundation for artificial blood was laid in 1949 when Amberson and colleagues reported the first infusion of cell-free hemoglobin for resuscitation in a patient with postpartum hemorrhage¹. While the concept was groundbreaking, the unrefined, stroma-containing free hemoglobin resulted in severe renal failure, coagulopathy, and fatal vasoconstriction¹. Decades later, the field experienced a brief triumph in 1989 when the United States Food and Drug Administration (FDA) approved Fluosol-DA-20, a perfluorocarbon-based product manufactured by Green Cross of Japan². However, due to the complexity of administration—requiring concurrent hyperbaric oxygen therapy—and significant side effects, it was withdrawn in 1994, remaining the only oxygen therapeutic ever fully approved by the FDA².

The late twentieth and early twenty-first centuries saw massive investments by the United States military and various pharmaceutical entities to develop safe Hemoglobin-Based Oxygen Carriers (HBOCs). Products such as HemAssist (a diaspirin cross-linked hemoglobin) and PolyHeme (a polymerized human hemoglobin) entered late-stage clinical trials⁶. Unfortunately, these trials culminated in a devastating setback. A 2008 meta-analysis published in the Journal of the American Medical Association (JAMA) pooled data across five distinct HBOC molecules—HemAssist, PolyHeme, Hemolink, Hemopure, and Hemospan—and revealed a 30 percent increase in the risk of death and a nearly threefold increase in the risk of myocardial infarction across the pooled trials³. The FDA subsequently placed the entire class of HBOC research on clinical hold³.

Only recently has the field begun to recover, driven by advances in nanotechnology, polymer chemistry, and the discovery of naturally occurring extracellular hemoglobins. These next-generation modalities seek to bypass the toxicological failures of the past, offering the promise of a universally compatible, shelf-stable powder that can be reconstituted in a well or trench to sustain life.

The Pathophysiology of Confined Hypoxia and Hypercapnia

To understand the engineering requirements of a synthetic blood substitute, one must first delineate the dual physiological crises of entrapment: hypoxemia and hypercapnia.

Under normal physiological conditions, oxygen is transported primarily by chemically binding to the iron-containing heme groups of hemoglobin within red blood cells (RBCs), while a minute fraction dissolves directly in the plasma². The average adult human red blood cell contains approximately 270 million hemoglobin molecules⁸. When entrapment limits ambient oxygen, tissues rapidly deplete their adenosine triphosphate (ATP) reserves. This forces cellular metabolism to shift from aerobic oxidative phosphorylation to anaerobic glycolysis, producing lactic acid as a byproduct⁹. The resulting metabolic acidosis leads to ischemic cell death, myocardial depression, and irreversible central nervous system damage⁹.

Simultaneous with oxygen starvation is the insidious threat of hypercapnia. Carbon dioxide, a constant byproduct of cellular respiration, is transported back to the lungs via three mechanisms: dissolved in plasma, bound to hemoglobin as carbaminohemoglobin, and predominantly converted into bicarbonate ions (HCO_3^-) by the enzyme carbonic anhydrase within red blood cells¹². In a confined space where the ambient atmosphere is not scrubbed, or where the patient is unable to ventilate (e.g., airway occlusion or burial), carbon dioxide accumulates in the bloodstream at a rate of 3 to 4 mmHg per minute¹⁴. Because carbon dioxide is highly lipid-soluble, it readily crosses the blood-brain barrier. Within the cerebrospinal fluid, it reacts with water to form carbonic acid, which immediately dissociates into hydrogen ions, plunging the local pH into severe respiratory acidosis¹⁵. This acidosis induces cerebral vasodilation, radically increasing intracranial pressure, while systemically causing neurological narcosis, lethal cardiac arrhythmias, and hemodynamic collapse¹⁵. Therefore, any theoretical "blood substitute" deployed in a confined environment must not only deliver oxygen but must interface seamlessly with the body's mechanisms for managing carbon dioxide and pH.

First and Second Generation Hemoglobin-Based Oxygen Carriers (HBOCs)

The fundamental premise of HBOC development is that hemoglobin can be extracted from outdated human donor blood or bovine (cow) blood, purified, and infused directly into the plasma as an acellular fluid¹. However, evolution designed hemoglobin to function exclusively within the protective envelope of the red blood cell membrane.

The Toxicity of Cell-Free Hemoglobin

When unmodified human hemoglobin is infused freely into the plasma, the 64 kDa tetrameric protein rapidly dissociates into 32 kDa dimers⁸. These dimers are small enough to be filtered by the renal glomeruli, leading to severe renal tubular obstruction, hemoglobinuria, and acute renal failure¹.

More disastrously, cell-free hemoglobin acts as a potent scavenger of endothelial nitric oxide (NO)⁷. Nitric oxide is a critical vasodilator that maintains microvascular perfusion. Because the free hemoglobin is no longer shielded by the RBC membrane, it extravasates into the interstitial spaces of the vascular endothelium, chemically stripping nitric oxide from the smooth muscle cells¹. This unopposed scavenging results in severe, systemic vasoconstriction, profound hypertension, reduced oxygen delivery to the microcirculation, and increased risks of myocardial infarction—the precise complications that halted the early HBOC trials⁶.

Polymerization and Cross-Linking: Hemopure

To prevent tetramer dissociation and mitigate nitric oxide scavenging, biochemists developed second-generation HBOCs by chemically cross-linking and polymerizing the hemoglobin molecules into larger macromolecular complexes⁷. The most prominent surviving product of this generation is Hemopure (HBOC-201), manufactured by HbO2 Therapeutics².

Hemopure utilizes a glutaraldehyde-polymerized bovine hemoglobin suspended in a balanced electrolyte solution². Because it is entirely acellular, it lacks human blood group antigens, meaning it requires no typing or cross-matching³. At a molecular weight of roughly 200–250 kDa, the polymerized hemoglobin is too large to easily extravasate into the endothelium, significantly reducing (though not entirely eliminating) the vasoactive side effects seen in first-generation products⁸.

Hemopure possesses highly advantageous logistical properties for emergency entrapment scenarios. It is shelf-stable at room temperature for at least three years, entirely bypassing the cold chain³. Packaged in a 250 mL bag, it is theoretically ideal for deployment in rescue vehicles, subterranean mining medical caches, or deep-sea rescue vessels³. While the 2008 FDA clinical hold relegated Hemopure to expanded access (compassionate use) protocols in the United States, it is fully approved and utilized as the standard of care for acute surgical anemia in South Africa and is approved in Russia².

Extracellular Marine Hemoglobin: M101 (HEMO2life)

A radically different approach to HBOC development involves sourcing hemoglobin that evolved naturally to function outside of a cell membrane, negating the need for artificial chemical cross-linking²². Hemarina, a French biotechnology firm, has commercialized M101, an extracellular hemoglobin isolated from the marine lugworm *Arenicola marina*¹⁰.

Lugworms endure dramatic environmental shifts, surviving prolonged hypoxia during low tides by relying on a highly specialized hemoglobin structure²². Unlike human or bovine hemoglobin, which are tetramers capable of binding exactly four oxygen molecules, M101 is a massive 3600 kDa hexagonal-bilayer nanoparticle capable of binding up to 156 oxygen molecules simultaneously¹⁰. Because it evolved over 450 million years to exist freely in the plasma of the invertebrate, it naturally resists dissociation and operates on a simple oxygen partial pressure gradient without the need for biological cofactors²³.

Clinical evaluations demonstrate that M101 does not induce the hypertensive vasoconstriction seen in early mammalian HBOCs²³. Furthermore, it possesses an intrinsic superoxide dismutase (SOD)-like antioxidant activity, driven by internal copper and zinc ions, which actively protects tissues from oxidative stress and ischemia-reperfusion injury¹⁰. Currently utilized as a CE-marked medical device additive (HEMO2life) for the static cold storage and machine perfusion of donor organs, M101 drastically reduces delayed graft function²⁵. In the national multicenter OXYOP clinical study, patient survival at 48 months post-transplantation was significantly higher in the HEMO2life group (98.3%) compared to the control group (86%)²⁴. Its ongoing development into a direct systemic intravenous oxygen carrier (under the pipeline name Hemoxycarrier) holds immense promise for trauma due to its massive oxygen payload, intrinsic safety profile, and lack of immunogenicity¹⁰.

Attribute	Polymerized Bovine	Extracellular Marine HBOC (M101 /
-----------	--------------------	-----------------------------------

	HBOC (Hemopure)	HEMO2life)
Origin	Extracted from bovine red blood cells ²² .	Harvested from <i>Arenicola marina</i> (lugworm) ²³ .
Molecular Structure	Glutaraldehyde-polymerized tetramers (200-250 kDa) ⁸ .	Natural hexagonal-bilayer nanoparticle (3600 kDa) ¹⁰ .
Oxygen Binding Capacity	4 O_2 molecules per tetramer subunit.	Up to 156 O_2 molecules per macromolecule ¹⁰ .
Vasoconstrictive Risk	Reduced compared to free Hb, but still present ⁸ .	Negligible; functions naturally as an extracellular protein ²³ .
Additional Properties	Acts as an immediate volume expander ⁸ .	Intrinsic SOD-like antioxidant activity (copper/zinc) ²⁴ .

Third Generation: Nanoscale Bio-Synthetic Artificial Red Cells

The most sophisticated advancement in AOC design abandons the idea of circulating free or polymerized hemoglobin entirely, instead attempting to synthetically recreate the full architecture of the human red blood cell. *ErythroMer*, currently in advanced pre-clinical development by the biotech firm KaloCyte (supported by DARPA, the NIH, and the DoD), is a "nano-cyte" that physically encapsulates purified human hemoglobin inside a soft, biocompatible lipid-oligomeric nanoparticle shell⁵.

Structural Architecture and Microvascular Penetration

At approximately 200 nanometers in diameter, ErythroMer is roughly 1/50th the size of a native human red blood cell⁵. This toroidal morphology provides a distinct rheological advantage: the nanoparticles can easily navigate compromised, constricted, or partially occluded microvasculature that native RBCs cannot penetrate, making it highly effective for reperfusion of ischemic tissues following trauma or shock⁵.

The soft lipid shell serves three critical protective functions. First, it contains the hemoglobin payload, preventing renal toxicity³⁰. Second, it possesses a specific surface charge (zeta potential) that prevents particle aggregation and improves overall biocompatibility³⁰. Third, and most importantly, the lipid shell provides a physical and chemical barrier to nitric oxide, completely preventing the NO-scavenging vasoconstriction that doomed earlier HBOC

generations¹⁹.

Dynamic Allosteric Control and the Bohr Effect Mimicry

The evolutionary brilliance of the native red blood cell lies in its ability to know *where* to release oxygen. Native hemoglobin relies on the allosteric effector 2,3-diphosphoglycerate (2,3-DPG) and local pH variations to alter its oxygen affinity (the Bohr effect)¹⁸. In the lungs, where pH is high, hemoglobin holds tightly to oxygen; in peripheral tissues, where cellular metabolism generates acidic carbon dioxide and lactic acid, the resulting low pH forces hemoglobin to rapidly offload its oxygen¹⁸.

Previous encapsulated hemoglobins lacked this mechanism, resulting in an abnormally low P_{50} (the partial pressure of oxygen at which 50% of hemoglobin is saturated). This meant the artificial cells captured oxygen in the lungs but refused to release it effectively to the tissues that needed it most¹⁹.

ErythroMer solves this physiological hurdle through the integration of a proprietary synthetic effector molecule designated KC1003 (sometimes interacting with modifiers like RSR13)²⁸.

KC1003 is packaged inside the lipid shell alongside the hemoglobin payload. It regulates oxygen affinity by dynamically capturing and releasing adenosine triphosphate (ATP) based on the ambient pH of the surrounding blood³⁰.

- **In the Lungs (High pH):** KC1003 binds to ATP. Without ATP interacting with the hemoglobin, the protein assumes a high-affinity state, maximizing oxygen capture from the alveoli³⁰.
- **In Hypoxic Tissues (Low pH):** As ErythroMer travels to peripheral tissues suffering from ischemia, the acidic environment triggers KC1003 to release the ATP. The free ATP binds allosterically to the hemoglobin, drastically shifting the oxygen dissociation curve to the right (increasing the P_{50}) and forcing the rapid, targeted offloading of oxygen directly into the oxygen-starved tissue³⁰.

This context-responsive control represents a massive leap forward, matching the vital oxygen-binding features of human red blood cells within 10 percent of natural efficacy²⁹.

Storage, Reconstitution, and Clinical Trajectory

For emergency entrapment scenarios, ErythroMer's formulation is revolutionary. The design incorporates low molecular weight polyethylene glycol (PEG) and carbohydrate-based lyoprotectants, allowing the entire nanoparticle suspension to be freeze-dried (lyophilized) into a stable powder⁸.

Once approved, ErythroMer will be supplied in a dual-chamber intravenous bag containing the dried red powder and sterile water. It is shelf-stable for years at ambient temperatures without refrigeration³⁰. In a disaster scenario—such as a mine collapse or battlefield trauma—a responder simply breaks the seal, mixes the solution, and administers a universally compatible blood substitute within minutes³⁰.

Based on rigorous in vivo modeling, ErythroMer restores normal hemodynamics and systemic oxygen delivery in animal models subjected to 40 percent blood volume loss¹⁹. The product

clears the body within 24 hours, with the lipid shell metabolized by phospholipases into endogenous phospholipids (like phosphatidylcholine), and the hemoglobin processed naturally by the liver³⁰. As of 2026, KaloCyte has secured over \$17 million in defense and federal grants, completing FDA Investigational New Drug (IND) enabling studies, including GLP-safety pharmacology and non-GLP pharmacokinetic biodistribution mapping, positioning the product for imminent Phase 1 human clinical trials²⁸.

Nanoparticle Property	ErythroMer Specification	Physiological Benefit in Entrapment
Size	~200 nanometers ⁵ .	Penetrates occluded microvasculature unavailable to native RBCs ²² .
Shell Composition	PEGylated lipid bilayer ⁸ .	Blocks NO scavenging; prevents vasoconstriction ¹⁹ .
Effector Molecule	KC1003 ³⁰ .	Senses low pH in hypoxic tissue; releases ATP to force O_2 offloading (Bohr mimicry) ³⁰ .
Storage State	Lyophilized powder ²⁹ .	No refrigeration needed; instant reconstitution in austere environments ³⁰ .

Perfluorocarbons (PFCs) and the Paradigm of Liquid Ventilation

Parallel to hemoglobin-based biochemistry is the utilization of Perfluorocarbons (PFCs) as synthetic oxygen carriers. PFCs are entirely synthetic, heavily fluorinated liquid hydrocarbons. Unlike hemoglobin, which binds oxygen chemically, PFCs dissolve respiratory gases through pure physical solution, mediated by exceedingly weak van der Waals forces¹⁸.

Gas Solubility Dynamics

The physical properties of PFCs make them uniquely suited for extreme gas exchange. While human blood carries roughly 20 mL of O_2 per 100 mL, specific saturated cycloaliphatic PFC

variants (such as Oxybyte) can dissolve up to 66 mL of O_2 per 100 mL³⁵. Even more astonishing is their affinity for carbon dioxide; a PFC solution can dissolve between 122 and 255 mL of CO_2 per 100 mL, rendering it exponentially more efficient at carrying CO_2 than water or biological plasma³⁵.

Because gas dissolution in PFCs strictly follows Henry's Law—meaning solubility is directly proportional to the partial pressure of the gas—PFCs require the patient to breathe

supplemental pure oxygen (F_iO_2 of 1.0) to achieve maximal oxygen loading². Because PFCs are highly hydrophobic, they will not mix with blood plasma and must be rigorously emulsified into micro-droplets (roughly 1/40th the diameter of an RBC) using lipid surfactants before intravenous administration². Prominent formulations include Perfluorodecalin (used in the Russian product Perftoran), Perfluorooctyl bromide (Perflubron), and experimental radiotherapeutics like NVX-108².

Total and Partial Liquid Ventilation

Beyond intravenous injection, the most striking application of PFCs in extreme medicine is Liquid Ventilation. Because PFCs hold vast quantities of dissolved gases, mammalian lungs can physically "breathe" the liquid².

In Total Liquid Ventilation (TLV), the lungs are filled to functional residual capacity (FRC)—approximately 30 mL/kg—with oxygenated PFC. A specialized mechanical liquid ventilator is then used to generate tidal breathing, cycling the heavy fluid in and out of the pulmonary tree³⁶. In Partial Liquid Ventilation (PLV), PFC is instilled into the lungs to replace the gas FRC, eliminating the alveolar air-liquid interface, while standard mechanical positive-pressure gas ventilation continues over the fluid column³⁶.

Because PFCs possess a remarkably low surface tension (14-18 dyne/cm) and a high density (nearly twice that of water), the liquid gravitates to the most dependent, collapsed regions of the lungs³⁶. It acts as a liquid continuous positive airway pressure (CPAP), forcing open atelectatic alveoli, recruiting lung volume, and establishing a barrier against inflammatory leukocyte infiltration³⁶. In entrapment scenarios complicated by severe blast-induced acute respiratory distress syndrome (ARDS) or toxic smoke inhalation, liquid ventilation provides a mechanism to deliver oxygen directly across the injured alveolar membrane while simultaneously lavaging heavy toxic exudates and cellular debris from the airways³⁶.

Clearance Mechanisms and the Carbon Dioxide Bottleneck

Unlike biological fluids, the human body lacks the metabolic machinery to break down PFCs. Instead, they are cleared physically. Intravenous PFC droplets are slowly phagocytized by macrophages of the reticuloendothelial system (RES), primarily in the liver and spleen³⁵. From there, the PFCs are gradually returned to the bloodstream, where their high vapor pressure allows them to transition into a gas upon reaching the pulmonary capillaries. The chemical is then permanently exhaled over a period of 48 hours to several weeks².

However, PFCs face a severe limitation regarding carbon dioxide mitigation. Despite their

massive theoretical capacity for CO_2 solubility, the heavy density and kinematic viscosity of these liquids (roughly twice that of water) create a severe "diffusion barrier"³⁵. During liquid breathing, if the mechanical exchange of the fluid is not meticulously optimized, the elimination of CO_2 lags significantly behind oxygen delivery. This kinematic bottleneck causes the partial pressure of CO_2 in the blood to spike rapidly, leading to fatal hypercapnia and respiratory acidosis despite adequate oxygenation³⁵.

Intravenous Oxygen Gas Delivery: Lipidic Oxygen Microparticles (LOMs)

While HBOCs and PFCs require the blood to circulate through functioning lungs to acquire oxygen, deep entrapment environments often feature entirely unbreathable atmospheres—such as complete airway obstruction, structural collapse, or submerged compartments. In these absolute emergencies, the lungs are rendered useless, and an intravenous carrier fluid cannot be oxygenated.

To address this, an alternative strategy focuses on injecting oxygen gas directly into the venous bloodstream. Historically, injecting raw gas into a vein leads to the coalescence of large bubbles, triggering massive pulmonary vascular obstruction, air embolism, and immediate cardiac arrest³⁸. However, researchers at Boston Children's Hospital, led by Dr. John Kheir, engineered a method to bypass this using microfluidic packaging³⁸.

Manufacture and Mechanism of LOMs

Utilizing a high-shear in-line homogenizer/sonicator, researchers forcefully combined oxygen gas with specific lipid molecules—primarily 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC) and cholesterol in a 1:1 molar ratio—creating an injectable foam suspension known as Lipidic Oxygen Microparticles (LOMs)³⁸.

The resulting LOMs consist of a single, highly deformable, semipermeable lipid monolayer encompassing a microscopic core of pure oxygen gas¹⁴. The microbubbles possess a median diameter of 0.93 to 4 micrometers, ensuring they are smaller than native red blood cells and can pass safely through the pulmonary microvasculature without causing mechanical obstruction¹⁴. The bulk emulsion can be manufactured rapidly (up to 4 liters in 10 minutes) and consists of roughly 60% to 70% oxygen gas by volume, providing an oxygen payload three to four times denser than that of native whole blood³⁸.

Efficacy in Total Asphyxia

When LOMs are injected intravenously, they rapidly mix with the deoxygenated venous blood. The massive concentration gradient forces the encapsulated oxygen gas to diffuse out of the lipid shell and bind instantly to the deoxyhemoglobin in the patient's native red blood cells. *In vitro* mixing studies demonstrate that this oxygen transfer is virtually instantaneous, completing within 3.95 to 4 seconds of contact¹⁴. As the oxygen gas vacates the lipid shell, the particle safely collapses by domain folding into small vesicles, avoiding hemolysis or complement

activation⁴⁰.

In extreme pre-clinical trials involving completely asphyxiated animal models (utilizing tracheal occlusion to simulate total airway blockage), continuous LOM infusion maintained systemic arterial oxygenation, preserved normal blood pressure, and prevented cardiac arrest and ischemic organ injury for 15 minutes without a single breath of air being taken¹⁴.

The Hypercapnia Limitation

While LOMs represent a miraculous bridging therapy—capable of buying paramedics or rescue divers a critical 15 to 30-minute window to establish an airway or extricate a trapped victim—they are fundamentally a one-way delivery system. They provide massive volumes of oxygen, but they possess zero capacity to absorb or remove carbon dioxide¹⁴. In the absence of ventilation, CO_2 rises rapidly, affording only a brief window before lethal hypercarbia and acidosis ensue¹⁴. Consequently, while LOMs stave off death by hypoxia, concurrent CO_2 mitigation becomes strictly necessary to ensure long-term survivability.

Intravenous and Chemical Mitigation of Carbon Dioxide (Hypercapnia)

The management of severe hypercapnia in environments where the lungs are incapacitated is arguably a more difficult physiological engineering challenge than oxygen delivery. In standard pressurized environments, CO_2 is traditionally mitigated using ambient chemical scrubbers such as Lithium Hydroxide (LiOH) or Soda Lime, which react with gaseous CO_2 to form stable carbonates³⁵. However, if the victim is buried or suffering from airway occlusion, CO_2 must be scrubbed directly from the bloodstream.

Pharmacological Buffering: THAM (Tromethamine)

When CO_2 cannot be exhaled, it accumulates in the blood and reacts with water (catalyzed by the enzyme carbonic anhydrase) to form carbonic acid (H_2CO_3), which rapidly dissociates into hydrogen ions (H^+) and bicarbonate (HCO_3^-). This flood of free hydrogen ions plunges the blood pH into severe acidosis³¹.


Traditionally, metabolic acidosis is treated with intravenous Sodium Bicarbonate. However, in a trapped or asphyxiated patient, Sodium Bicarbonate is lethal. The administration of exogenous bicarbonate drives the chemical equilibrium backward, generating massive amounts of *new* CO_2 gas. If the patient cannot exhale this generated gas, the intracellular acidosis worsens drastically¹⁵.

To solve this, emergency protocols utilize an intravenous amino alcohol buffer known as Tromethamine (Tris-hydroxymethyl aminomethane, commercially known as THAM)¹⁵.

Biochemical Mechanism of THAM

Unlike bicarbonate, THAM is a pure proton acceptor containing no sodium, and it does not generate carbon dioxide when it buffers acid¹⁶. At normal body temperature (37°C), THAM possesses a pK_a of 7.8 (compared to bicarbonate's 6.1), making it a highly effective physiological buffer¹⁵.

When infused intravenously as a 0.3 Molar solution, THAM directly targets carbonic acid:



By binding the free hydrogen ions of carbonic acid, THAM actively pulls dissolved CO_2 out of the plasma and converts it into stable bicarbonate³¹. In essence, THAM acts as an intravenous chemical CO_2 scrubber. Approximately 30% of THAM remains un-ionized at physiological pH, allowing it to penetrate cell membranes and cross the blood-brain barrier. This provides vital intracellular buffering, rapidly decreasing the partial pressure of carbon dioxide in the brain and reversing the intracranial pressure caused by CO_2 -induced cerebral vasodilation¹⁵. The protonated THAM-complex is subsequently excreted by the kidneys, inducing an osmotic diuresis that flushes the trapped acid from the body at a rate slightly exceeding standard creatinine clearance¹⁵.

Dosage and Clinical Limitations

The empirical dosage for THAM is calculated precisely based on the patient's base deficit and lean body weight:

$$\text{Volume of 0.3M THAM (mL)} = \text{Lean Body W}$$

(Note: Clinical guidelines often multiply this by a factor of 1.1 to account for buffering capacity reduction due to the acetic acid used to balance the solution's pH to 8.6)¹⁵.

While highly effective at stalling death by respiratory acidosis, THAM presents specific side effects. Because it rapidly decreases $PaCO_2$ and raises pH, it eliminates the biological drive to breathe, causing profound respiratory depression¹⁵. Furthermore, the un-ionized form of THAM stimulates insulin release and increases insulin sensitivity, posing a significant risk of severe, prolonged hypoglycemia¹⁵. Extravasation of the highly alkaline solution can also cause severe local tissue necrosis and chemical phlebitis, necessitating administration through a central line or large peripheral vein⁴³.

Extracorporeal and Intravenous Mechanical CO2 Removal

If pharmacological buffering is insufficient for prolonged entrapment, mechanical venous interception is required.

- **Intravenous Oxygenator and CO₂ Removal Device (IVOX):** Surgically deployed into the inferior or superior vena cava via a femoral or jugular venotomy, the IVOX consists of a bundle of highly permeable hollow fibers connected to external gas conduits. As venous blood flows over the fibers, CO_2 is removed via a vacuum pump gradient, and oxygen is infused⁴⁵. While highly localized, it removes approximately 30% of basal CO_2 production without requiring a traditional extracorporeal pump⁴⁵.
- **Impeller Percutaneous Respiratory Assist Catheter (IPRAC):** An advancement over passive fibers, the IPRAC utilizes an array of high-speed rotating impellers mounted on an intravenous catheter. Placed percutaneously, the impellers vigorously mix the blood against the surrounding gas exchange surfaces, breaking up the boundary layer and drastically enhancing CO_2 transfer efficiency. This maximization allows the catheter profile to be reduced from 25 French (8.3 mm) to 20 French (6.7 mm), making rapid insertion feasible in austere emergency environments⁴⁶. Advanced IPRAC models integrate carbonic anhydrase enzyme biocatalytic coatings on the gas exchange surfaces to actively strip CO_2 from the blood at vastly accelerated rates⁴⁶.

Nanomaterial and Amino Acid Chemical CO₂ Scrubbing

Outside the body, advances in nanomaterials and chemistry offer new avenues for ambient CO₂ scrubbing in confined spaces. Amino acid salt solutions (AAS), such as GAPL (a blend of

Glycine, Alanine, Proline, and Lysine), have demonstrated a 19% higher CO_2 absorption

capacity (0.91 mol CO_2 /mol amino acids) compared to standard monoethanolamine (MEA) industrial scrubbers⁴⁷. These solutions are nonvolatile, stable, noncorrosive, and nontoxic, making them ideal for regenerating life-support systems in sealed vessels⁴⁷.

Furthermore, carbon-based nanomaterials, including hollow mesoporous carbon spheres and graphene oxide decorated with silver nanoparticles (Ag NPs), are being engineered to capture

and catalytically reduce CO_2 buildup⁴⁸. While currently employed primarily in hemodialysis to remove protein-bound toxins (hemoperfusion)⁴⁹, the integration of highly conductive silver nanoparticles within defective metal-organic frameworks (e.g., Ag@UiO-66-SH) has shown an

astounding Faradaic efficiency of 74% for electrocatalytically converting ambient CO_2 back into carbon monoxide and potentially usable fuels, offering a closed-loop atmospheric regeneration system for deep-entrapment pressure vessels⁵⁰.

Peritoneal Dialysis for CO₂ Clearance

An unconventional but highly effective method for clearing CO_2 without establishing central

vascular access involves utilizing the highly vascularized peritoneal membrane of the abdomen⁵¹. In situations where CO_2 levels are lethal, infusing hyperosmotic dialysis solutions (e.g., 15% dextrose) tailored with a highly alkaline pH (7.6) into the peritoneal cavity creates a massive diffusion gradient. Dissolved CO_2 diffuses rapidly from the mesenteric blood supply into the abdominal fluid pool, which is subsequently drained out of the body⁵². While this technique has primarily been studied in the context of mitigating complications from carbon dioxide pneumoperitoneum during laparoscopic surgery⁵¹, the thermodynamic principles of peritoneal CO_2 clearance offer a highly viable, low-tech temporizing measure for acute hypercapnia in austere trauma settings where hemodialysis or ECMO is impossible⁵¹.

CO2 Mitigation Modality	Mechanism of Action	Clinical Application in Entrapment	Primary Limitation
THAM (Tromethamine)	Intravenous amino alcohol binds H^+ to convert CO_2 to bicarbonate ³¹ .	Immediate buffering of severe respiratory acidosis ¹⁵ .	Induces respiratory depression and severe hypoglycemia ¹⁵ .
IPRAC Catheter	Intravenous impellers maximize blood contact with gas exchange fibers ⁴⁶ .	Percutaneous insertion for mechanical CO_2 extraction ⁴⁶ .	Requires invasive vascular access and external vacuum drive ⁴⁵ .
Amino Acid Scrubbing	GAPL solutions chemically bind ambient CO_2 in closed systems ⁴⁷ .	High-capacity, non-toxic atmospheric scrubbing for sealed vessels ⁴⁷ .	Environmental only; does not treat blood acidosis directly.
Peritoneal Dialysis	Hyperosmotic/alkaline fluid in the abdomen draws CO_2 from mesenteric	Low-tech alternative to ECMO for systemic CO_2 clearance ⁵² .	Requires abdominal catheterization and fluid cycling ⁵¹ .

	blood ⁵² .		
--	-----------------------	--	--

Synthesis: A Comprehensive Paradigm for Entrapment Survival

To contextualize the integration of these technologies, one must consider the operational reality of a catastrophic entrapment: a miner trapped in a collapsed, non-ventilated subterranean void, or a submariner in a disabled vessel where the atmosphere has become critically hypoxic and hypercapnic. Standard rescue operations involving heavy extrication or deep-sea recovery may require hours or days, far exceeding the biological limits of the trapped individuals.

With the advent of these pharmacological and biochemical tools, the survivability timeline is fundamentally altered, moving life support from the external environment into the vascular compartment:

1. **Immediate Rescue (0–30 Minutes):** Through a small access bore-hole or umbilical, paramedics can pass a pre-loaded syringe of Lipidic Oxygen Microparticles (LOMs). Direct intravenous injection provides an immediate, life-sustaining delivery of concentrated oxygen gas directly to the brain and myocardium. This buys a critical 15 to 30-minute window for further intervention without requiring the victim to breathe the toxic ambient atmosphere³⁸.
2. **Sustained Support and Acidosis Reversal (Hours):** To combat the inevitable, rapid rise in CO_2 resulting from the absence of ventilation and the LOM infusion, a localized IV drip of Tromethamine (THAM) is initiated. THAM acts as an intravenous chemical CO_2 sink inside the plasma, pulling the gas into stable bicarbonate and reversing the fatal intracranial pressure and respiratory acidosis¹⁵. If the situation is prolonged, rescue workers can instill hyperosmotic alkaline fluid into the peritoneal cavity, utilizing the abdomen as a makeshift lung to dialyze CO_2 out of the body⁵².
3. **Hemorrhage Control and Microvascular Reperfusion:** If the victim has sustained crush injuries, traumatic amputations, or massive hemorrhage, native red blood cells are lost, rendering standard oxygen therapy useless. In this scenario, lyophilized ErythroMer (stored ambiently in the rescue kit) is rapidly reconstituted with sterile water³⁰. The 200-nanometer nanoparticles bypass crushed microvasculature, utilizing their KC1003 effector molecules to sense the highly acidic, hypoxic environments of the damaged tissue⁵. This provides a massive volume expansion with highly efficient, pH-regulated oxygen delivery, counteracting hemorrhagic shock and maintaining perfusion until heavy machinery can complete the extrication²⁷.

By leveraging the extracellular carrying capacity of marine hemoglobin (M101), the bio-mimicry of nano-synthetic red blood cells (ErythroMer), the raw gas delivery of LOMs, and the chemical buffering power of THAM, modern trauma medicine is no longer bound entirely by the constraints of ambient atmospheric pressure. These advancements represent a profound shift in

the engineering of human survival, allowing responders to sustain the fundamental biochemical requirements of life even when the external environment has completely failed.

Works cited

1. Artificial Blood: A Futuristic Dimension of Modern Day Transfusion Sciences - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6864588/>
2. Blood substitute - Wikipedia, https://en.wikipedia.org/wiki/Blood_substitute
3. Hemopure: The Blood Substitute That Almost Was, <https://www.bloodbytesbeyond.com/post/hemopure-the-blood-substitute-that-almost-was>
4. Banking on Artificial Blood | ASH Clinical News | American Society of Hematology, <https://ashpublications.org/ashclinicalnews/news/8465/Banking-on-Artificial-Blood>
5. KaloCyte: A biotech developing a first of its kind blood substitute | Meet Life Sciences, <https://www.meetlifesciences.com/media-hub/blog/kalocyte--a-biotech-developing-a-universal--first-of-its-kind-blood-substitute/>
6. Hemoglobin-Based Oxygen Carriers: Where Are We Now in 2023? - MDPI, <https://www.mdpi.com/1648-9144/59/2/396>
7. Blood Substitute Anyone? - Centre for Blood Research - The University of British Columbia, <https://cbr.ubc.ca/blood-substitute-anyone/>
8. Biosynthetic blood surrogates: Current status and future opportunities - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12617565/>
9. A Novel Oxygen Carrier (M101) Attenuates Ischemia-Reperfusion Injuries during Static Cold Storage in Steatotic Livers - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8395216/>
10. The oxygen carrier M101 alleviates complement activation, which may be beneficial for donor organ preservation - Frontiers, <https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2022.1006761/full>
11. Artificial oxygen carriers in emergency medicine | Kobińska | Medical Research Journal, https://journals.viamedica.pl/medical_research_journal/article/view/96727
12. Therapeutic delivery of oxygen using artificial oxygen carriers demonstrates the possibility of treating a wide range of diseases - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11742488/>
13. [PDF] Elimination of CO₂ in patients with carbonic anhydrase II, <https://www.semanticscholar.org/paper/46f798b0aed6e91cb0ef29157ae570daefd3785>
14. Kheir 1..10, <http://www.medlive.cn/uploadfile/2012/0629/20120629041757584.pdf>
15. (PDF) Guidelines for the Treatment of Acidaemia with THAM: - ResearchGate, https://www.researchgate.net/publication/51303669_Guidelines_for_the_Treatment_of_Acidaemia_with_THAM
16. Buffer therapies: sodium bicarbonate, Carbicarb and THAM | Deranged Physiology, <https://derangedphysiology.com/main/required-reading/electrolyte-disorders/Chapt>

- [er-223/buffer-therapies-sodium-bicarbonate-carbicarb-and-tham](#)
17. High-Flow Nasal Oxygen in Contemporary Airway Management: Enhancing Physiological Safety in Anaesthesia—A Narrative Review - MDPI, <https://www.mdpi.com/2227-9032/14/14/2206>
 18. Artificial Blood: The History and Current Perspectives of Blood Substitutes - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7086064/>
 19. Erythromer (EM), a Nanoscale Bio-Synthetic Artificial Red Cell | Request PDF, https://www.researchgate.net/publication/362554182_Erythromer_EM_a_Nanoscale_Bio-Synthetic_Artificial_Red_Cell
 20. Hemoglobin-Based Oxygen Carriers – Are We There Yet? - Red Cross Plus Online Blog, <https://redcrossplus.blog/hemoglobin-based-oxygen-carriers-are-we-there-yet/>
 21. Artificial Blood and Oxygen Carriers: The Next Revolution in Intraoperative Safety, <https://globalrph.com/2026/02/artificial-blood-and-oxygen-carriers-the-next-revolution-in-intraoperative-safety/>
 22. Red Blood Cell Substitutes - HbO2 Therapeutics, <https://www.hbo2therapeutics.com/post/red-blood-cell-substitutes>
 23. HEMOXYCarrier® - Hemarina, <https://www.hemarina.com/ours-products/hemoxycarrier/>
 24. Long Term Results of the Use of the Oxygen Carrier M101 as an Additive to Organ Preservation Solution - World Scientific Publishing, <https://www.worldscientific.com/doi/pdf/10.1142/S2972497X24200013?download=true>
 25. HEMO2life® - Hemarina, <https://www.hemarina.com/ours-products/hemo2life/>
 26. Efficacy of the natural oxygen transporter HEMO2 life® in cold preservation in a preclinical porcine model of donation after cardiac death - PubMed, <https://pubmed.ncbi.nlm.nih.gov/30924562/>
 27. Erythromer - Wikipedia, <https://en.wikipedia.org/wiki/Erythromer>
 28. ErythroMer: Nanoscale BioSynthetic Red Cell Substitute - NIH RePORTER, <https://reporter.nih.gov/project-details/11159433>
 29. ErythroMer as a Step Forward in Artificial Blood - Fight Aging!, <https://www.fightingaging.org/archives/2016/12/erythromer-as-a-step-forward-in-artificial-blood/>
 30. ERYTHROMER | KaloCyte, <https://kalocyte.com/erythromer/>
 31. THAM SOLUTION Tromethamine Injection, <https://labeling.pfizer.com/ShowLabeling.aspx?id=4642>
 32. Effects of a Synthetic Allosteric Modifier of Hemoglobin Oxygen Affinity on Outcome From Global Cerebral Ischemia in the Rat - American Heart Association Journals, <https://www.ahajournals.org/doi/10.1161/01.str.29.8.1650>
 33. Nano blood cells could offer convenient, portable alternative to transfusion, <https://grainger.illinois.edu/news/stories/29196>
 34. KaloCyte Selected to Join BLUE KNIGHT - UM Ventures, <https://www.umventures.org/news/kalocyte-selected-join-blue-knight>
 35. https://drive.google.com/open?id=1bMyg3XEC8X8neHSWTQvHhWK78FLDpJPCF3_iwWMYA3Q

36. Liquid ventilation - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4258983/>
37. Liquid Ventilation - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3191624/>
38. Researchers Develop Alternative Method for Oxygenating Blood - Respiratory Therapy,
<https://respiratory-therapy.com/disorders-diseases/infectious-diseases/other-infections/researchers-develop-alternative-method-for-oxygenating-blood/>
39. Injected microbubbles could be a safe way to deliver emergency oxygen,
<https://answers.childrenshospital.org/injected-microbubbles-could-be-a-safe-way-to-deliver-emergency-oxygen/>
40. Bulk manufacture of concentrated oxygen gas-filled microparticles for intravenous oxygen delivery - PubMed, <https://pubmed.ncbi.nlm.nih.gov/23471884/>
41. (PDF) Oxygen Gas-Filled Microparticles Provide Intravenous Oxygen Delivery - ResearchGate,
https://www.researchgate.net/publication/228089270_Oxygen_Gas-Filled_Microparticles_Provide_Intravenous_Oxygen_Delivery
42. What is a Refuge Chamber Scrubbing System? - MineARC Systems,
<https://minearc.com/what-is-a-refuge-chamber-scrubbing-system/>
43. Tromethamine Injection - DailyMed,
<https://dailymed.nlm.nih.gov/dailymed/fda/fdaDrugXsl.cfm?setid=57db5686-7f7f-4999-87e3-e1cbbda4a98d&type=display>
44. Tromethamine | CHEO ED Outreach, <https://outreach.cheo.on.ca/manual/1727>
45. Lung replacement therapies for acute respiratory failure,
<https://hrcak.srce.hr/file/251020>
46. Richard Jefferies - Swanson School of Engineering,
<https://www.engineering.pitt.edu/subsites/programs/cbtp/people/past-trainees/jefferies-richard/>
47. Enhanced Carbon Dioxide Capture Using a Mixed Amino Acid Salt Solution,
<https://pubs.acs.org/doi/10.1021/acs.iecr.2c03533>
48. Metal Nanoparticles and Carbon-Based Nanomaterials for Improved Performances of Electrochemical (Bio)Sensors with Biomedical Applications - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8585379/>
49. Functional Carbon-Based Materials for Blood Purification: Recent Advances Toward Improved Treatment of Renal Failure and Patient Quality of Life - MDPI,
<https://www.mdpi.com/2306-5354/12/8/893>
50. Silver Nanoparticle-Decorated Defective Zr-Based Metal-Organic Frameworks for Efficient Electrocatalytic Carbon Dioxide Reduction with Ultrahigh Mass Activity - ACS Publications, <https://pubs.acs.org/doi/10.1021/acsaem.3c00299>
51. Videolaparoscopic peritoneal dialysis catheter implant and rescue procedures under local anesthesia with nitrous oxide pneumoperitoneum - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/10649698/>
52. Peritoneal transfer of carbon dioxide in the rat - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/1450480/>
53. a. Carbon dioxide (CO₂) inflation using the peritoneal dialysis... - ResearchGate,
https://www.researchgate.net/figure/a-Carbon-dioxide-CO-2-inflation-using-the-peritoneal-dialysis-catheter-b_fig1_380854028