

Anthropocanine Sympatry: Neurobiological, Archaeogenetic, and Ethological Co-Modulation in *Canis familiaris* and *Homo sapiens*

Introduction to Interspecies Neuroethology and Sympatric Evolution

The evolutionary trajectory of *Canis lupus familiaris* and *Homo sapiens* represents an unprecedented instance of interspecies neurobiological and genomic co-modulation. The sympatric development of these species extends significantly beyond rudimentary domestication and utilitarian paradigms, encompassing reciprocal physiological adaptations, shared autonomic nervous system synchronization, and complex neuroendocrine feedback loops. Current research delineates highly specific modalities through which canines and humans interact, heavily implicating the autonomic nervous system, somatosensory processing pathways, and highly specialized chemosensory and magnetoreceptive apparatus capable of detecting subtle environmental and physiological frequencies. Furthermore, the genomic architecture of the modern domestic dog reflects a composite of natural population bottlenecks—often resulting from Pleistocene environmental cataclysms—and subsequent, intense anthropogenic selective pressures that exploited inherent genomic instabilities.

Autonomic Co-Modulation and Heart Rate Variability

The physiological linkage between canines and humans manifests acutely in the autonomic nervous system (ANS), which regulates involuntary bodily functions via its sympathetic (arousal/alertness) and parasympathetic (rest/digestion) branches¹. This interspecies linkage is quantified empirically through Heart Rate Variability (HRV), defined as the fluctuation in time intervals between adjacent heartbeats (RR intervals), serving as a reliable index of cardiac vagal tone¹.

Synchronization of the Autonomic Nervous System

High HRV indicates a state of relaxation and recovery, mediated by parasympathetic dominance, whereas low HRV signifies sympathetic dominance, indicative of stress, stimulation, or strain¹. Recent electrophysiological studies utilizing electrocardiogram (ECG) electrodes on freely behaving dog-owner dyads reveal a pronounced physiological synchrony, formally termed "co-modulation"³.

During interspecies interaction, the HRV of a dog and its human handler mirror one another³. When a human exhibits high HRV, the canine counterpart concurrently exhibits high HRV,

demonstrating interspecific emotional contagion². Notably, this synchronization is heavily context-dependent and temporally distinct. During free-form resting periods, overall HRV correlations are most pronounced, whereas physical activity levels synchronize primarily during predefined interaction tasks, such as playing or structured petting¹.

The statistical predictability of this co-modulation is highly specific. Canine overall HRV acts as the singular predictive factor for human overall HRV¹. Conversely, human negative affectivity—a temperamental trait reflecting a human's propensity for concern, anxiety, or hyper-vigilance—correlates positively with canine HRV¹. This counterintuitive finding suggests that canines, particularly herding breeds, develop compensatory attachment bonds that elevate their own sense of safety and parasympathetic tone in the presence of anxious handlers, effectively acting as an autonomic buffer¹.

Metric	Physiological Definition	Indication in Human-Canine Dyads
HRmean	Mean Heart Rate	Decreases progressively during familiar, positive handling and proximity ² .
RMSSD	Root mean square of successive differences	Increases over repeated familiar interactions, indicating rising parasympathetic tone ² .
HFlog	High-frequency absolute power component (log transformed)	Peaks during states of interspecies emotional contagion and mutual relaxation ² .
Cortisol	Primary glucocorticoid stress hormone	Decreases in both species during tactile and visual interaction ⁵ .
Oxytocin	Hypothalamic neuropeptide	Increases via positive feedback loop driven by mutual gaze and tactile stimulation ⁹ .

The Somatosensory Processing of Affiliation: C-Tactile Afferents

Human-canine physical interactions—such as humans laying on the ground, engaging in playfulness, permitting dogs to lick their faces, and rubbing different parts of the canine body—activate highly specific somatosensory pathways. The physiological and emotional responses to this tactile stimulation, or allogrooming, are governed by the activation of distinct cutaneous sensory nerves¹⁰.

C-Tactile Afferents and the Insular Cortex

Mammalian hairy skin contains specialized low-threshold, unmyelinated peripheral nerve fibers known as C-tactile (CT) afferents¹⁰. These mechanoreceptors are evolutionarily tuned to process affiliative, social touch. CT afferents are maximally excited by slow, caress-like stroking at velocities of 1 to 10 cm per second, applied with a gentle force at typical conspecific body temperatures¹⁰.

Unlike fast-conducting, myelinated A-beta afferents that project to the primary somatosensory cortex to process discriminative touch, CT afferents project via the spinothalamic tract directly to the anterior insular cortex and the posterior superior temporal sulcus¹³. The insular cortex operates as a neural hub integral to interoception, emotional processing, and the conscious perception of pleasantness¹⁰. Consequently, tactile stimulation such as rhythmic rubbing or petting bypasses strictly analytical sensory pathways, routing directly into the affective neural circuitry of both species.

The Oxytocinergic Two-Step Activation Model

The application of tactile stimulation initiates a complex neuropeptide cascade, facilitating interspecies bond maintenance. Gentle touch activates a two-neuron pathway involving tachykinin 1-neurons originating in the periaqueductal gray (PAG), which subsequently project to the supraoptic (SON) and paraventricular (PVN) nuclei of the hypothalamus¹⁰. This projection triggers the release of oxytocin (OT) into the central nervous system, where it can act both as a classical neurotransmitter and via paracrine diffusion, reaching distant oxytocin receptors (OXTR)¹⁰.

A secondary phase of tactile interaction, characterized by firmer pressure (e.g., deep rubbing or sustained physical contact such as laying together), triggers a second pool of cutaneous afferents¹⁰. This deeper mechanoreception engages distinct oxytocinergic pathways associated with profound anti-stress effects, significant reductions in pain, lowering of blood pressure, and enhanced vagal activation of the gastrointestinal endocrine system, driving systemic restorative processes¹⁰. The act of canine facial licking and reciprocal human petting stimulates the release of endogenous opioids and dopamine alongside oxytocin, cementing a state of reward, mutual psychological security, and a lowering of baseline aggression¹³.

This synchronization is sustained biochemically by a positive oxytocin-cortisol feedback loop⁷. The release of OT in both species effectively inhibits the hypothalamic-pituitary-adrenal (HPA) axis, reducing the secretion of glucocorticoids (GCs) such as cortisol⁵. Research investigating visual pathways demonstrates that mutual gazing between humans and canines triggers this exact oxytocinergic feedback loop, effectively hijacking the mammalian maternal-infant bonding mechanism¹¹. Notably, this interspecies gaze-bonding mechanism is absent in hand-reared

wolves, indicating that the cognitive architecture for this hormonal hijack evolved strictly post-domestication¹¹.

Chemosensation, Forensic Applications, and Environmental Threat Detection

Canines possess profound capabilities to detect danger, shifts in emotional frequencies, and hazardous environments, helping humans navigate and survive otherwise imperceptible threats. These abilities are fundamentally rooted in their advanced chemosensory systems, which operate far beyond human perceptual thresholds.

The Vomeronasal Organ and Volatile Organic Compounds

While the main olfactory system processes general environmental odors, the vomeronasal organ (VNO), or Jacobson's organ, specializes in detecting non-volatile and volatile organic compounds (VOCs), including pheromones and chemical markers of physiological states²⁰. Located in the soft tissue of the nasal septum just above the hard palate, the VNO communicates directly with the brain via cranial nerve zero (CN 0), projecting to the accessory olfactory bulb, the amygdala, and the anterior hypothalamus²².

Recent immunohistochemical analyses indicate that canine VNO neurons predominantly express V1R (VR1) receptor genes, contrasting with rodents that express both V1R and V2R²³. Despite this genomic regression, the transduction pathways within the canine VNO are highly sophisticated. Signal transduction operates via a dual-pathway mechanism following the activation of phospholipase C (PLC)²⁰.

1. **TRPC2 Pathway:** PLC cleaves phosphatidylinositol 4,5-bisphosphate into inositol 1,4,5-trisphosphate (IP3) and diacylglycerol (DAG). DAG directly activates the Transient Receptor Potential Cation Channel, Subfamily C, Member 2 (TRPC2), leading to immediate depolarization and calcium influx²⁰.
2. **Arachidonic Acid Pathway:** Operating independently of TRPC2, DAG is metabolized by DAG lipase into arachidonic acid (AA), which subsequently activates a distinct, Ca²⁺-permeable channel²⁰.

This dual redundancy ensures extreme sensitivity to specific VOCs. Through the VNO and the main olfactory epithelium, dogs detect human fear, stress, and anxiety via the precise quantification of human cortisol and adrenaline metabolites secreted through apocrine sweat glands and respiration²¹. When humans enter states of high autonomic arousal—often conceptualized colloquially as existing on "dangerous frequencies"—canines perceive these neurochemical shifts instantaneously. Brain scans confirm that canine neural networks generate distinct, unique activity patterns that actively differentiate human expressions and chemosensory signatures of fear, anger, and sadness²⁸.

This chemosensory acuity is highly leveraged in forensic and operational human assistance. Human remain detection (HRD) dogs utilize these olfactory mechanisms to identify VOCs emitted from dry, weathered bones. Studies employing comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry (GCxGC-TOFMS) have confirmed that

canines can isolate and alert to specific VOC profiles originating from skeletal remains possessing virtually no residual organic matter, providing critical assistance in disaster, forensic, and cataclysmic recovery scenarios²³.

Magnetoreception and Non-Standard Sensory Modalities

Beyond chemical sensing, canines demonstrate an ability to perceive shifts in geomagnetic fields, interacting with environmental "frequencies" that evade human detection. While peripheral investigations into human-canine interactions have explored fringe concepts—such as the "psychic pet phenomenon" popularized by the experimental replication tests of Wiseman and Sheldrake regarding anticipatory behavior—empirical science grounds these seemingly telepathic connections in hyper-acute, unrecognized sensory cues³⁰. Among the most profound of these is quantum magnetoreception.

Mammalian magnetoreception is theorized to operate via a quantum radical pair mechanism³¹. Researchers have identified the light-sensitive flavoprotein *cryptochrome 1a* (Cry1a) within the outer segments of the ultraviolet/blue-sensitive S1 cone photoreceptors in the canine retina³². Cryptochromes are the only known animal photopigments capable of forming spin-correlated radical pair intermediates³¹.

When a photon of blue light excites a flavin adenine dinucleotide (FAD) molecule within the cryptochrome, an electron is transferred from a nearby tryptophan amino acid, creating a radical pair with quantum entangled electrons³¹. The singlet-triplet interconversion rate of this radical pair is exceedingly sensitive to the alignment of external magnetic fields³¹. Because Cry1a is found in dog-like carnivores (Canidae) but is absent in cat-like carnivores (Felidae) and many other mammalian orders, researchers hypothesize that canines perceive magnetic inclinations as a superimposed visual pattern across their field of vision³³.

Furthermore, the canine vibrissal system, including macrovibrissae on the muzzle and highly innervated microvibrissae on the upper lip, functions as a network of hypersensitive mechanoreceptors³⁷. These structures detect microscopic variations in air currents and vibrational frequencies, augmenting their awareness of environmental anomalies and corroborating reports of canines alerting humans to impending environmental dangers.

Archaeogenetics: The *Kóryos* and the Proto-Indo-European *Männerbund*

The historical entwinement of humans and dogs involves complex socio-religious integration, predating modern conceptualizations of pet ownership and stretching into the militaristic foundations of European civilization. A critical vector for this integration occurred in the Late Bronze Age Eurasian steppes among Proto-Indo-European (PIE) cultures.

Linguistic and archaeological evidence reconstructs a central PIE institution known as the *Kóryos* (from the PIE root **kóro-*, meaning "cutting, section, or division")³⁸. The *Kóryos* functioned as a *Männerbund*—an alliance of unmarried male adolescents, typically aged 12 to

19, who operated as a landless, roving warrior band³⁸.

This demographic cohort existed in a state of societal liminality. As a rite of passage into adulthood and full societal integration, these youths were expelled from the host community into the wild during the winter months³⁹. During this period, the youths underwent a mystical self-identification with wolves and dogs, animals symbolizing death, lawlessness, liminality, and warrior fury³⁹.

Archaeological excavations at the Srubnaya culture settlement of Krasnosamarskoe (circa 1750 BCE) in the Samara River valley of Russia yielded profound empirical evidence of this institution⁴⁰. Excavators David W. Anthony and Dorcas R. Brown identified a highly specific midwinter New Year festival dedicated to warrior initiations involving mass dog sacrifices³⁸. Canine and wolf remains constituted over 30% of all osteological material at the Krasnosamarskoe site—a statistical anomaly compared to the <3% typical of surrounding sites⁴⁰. The butchery patterns were strictly ritualistic, not nutritional. Snouts were systematically severed into three pieces, and the crania were meticulously fragmented into geometric shapes⁴⁰. Dogs of advanced age (up to 12 years) were sacrificed exclusively during the winter solstice⁴⁰.

In the PIE cosmology, multi-headed dogs guarded the entrance to the Afterworld (e.g., Cerberus, Saranyu)⁴³. By consuming dog flesh and donning wolf or dog skins, the *Kóryos* initiates symbolically died to their boyhood identities, embodying the predatory nature of the canine⁴¹. Upon completion of the raiding season, the removal of the canine skins marked their resurrection and reintegration into civilized society as adult warriors³⁹. This foundational mythological and practical alliance facilitated the violent expansion of Indo-European languages and cultures across Eurasia. The institutionalized violence and strict hierarchical structures of the *Kóryos* eventually evolved into the martial and aristocratic classes that established the foundational monarchies of Europe, intertwining human socio-political evolution inextricably with canine symbolism and utility³⁸.

Genomic Variance: Cataclysmic Bottlenecks vs. Anthropogenic Morphological Engineering

The immense phenotypic variance observed in modern *Canis familiaris* invites inquiry into its origins: specifically, whether this variance is the byproduct of ancient environmental cataclysms and resultant population bottlenecks in wolves, or the result of deliberate anthropogenic selection.

Genomic studies indicate that while natural selection and cataclysmic population bottlenecks—such as Pleistocene glaciation events—shaped the baseline genetic architecture of the ancestral wolf populations, the extreme morphological variance seen today is almost exclusively the result of intense, human-directed artificial selection⁴⁵. Humans utilized the inherent instability of the canine genome—characterized by rapidly mutating microsatellites and abundant short interspersed nuclear elements (SINEs) and long interspersed nuclear elements (LINEs)—to engineer specific morphological and behavioral traits over centuries⁴⁷.

The *FGF4* Retrogene and Chondrodysplasia

A definitive example of this evolutionary manipulation is the presence of breed-defining disproportionate dwarfism, or chondrodysplasia (CDPA). The genetic etiology of CDPA is not a standard point mutation, but rather an RNA-mediated retrotransposition event⁴⁵.

During the evolution of the domestic dog, the processed mRNA transcript of the Fibroblast Growth Factor 4 (*FGF4*) gene was reverse-transcribed and inserted as a functional retrogene into a LINE on a completely different chromosome⁴⁸.

1. **FGF4L1 (Chromosome 18):** An insertion on CFA18 causes chondrodysplasia (CDPA). This phenotype defines at least 19 breeds, including the Dachshund, Corgi, and Basset Hound, resulting in significantly shortened long bones and varying degrees of carpal valgus⁴⁵.
2. **FGF4L2 (Chromosome 12):** A separate *FGF4* retro-insertion on CFA12 causes chondrodystrophy (CDDY)⁴⁸.

While both retrogenes restrict long bone growth, the *FGF4* insertion on CFA12 is a highly deleterious pleiotropic locus. The extra gene copy overexpresses the FGF4 protein tenfold within the intervertebral discs⁴⁹. This triggers premature calcification of the nucleus pulposus in the intervertebral discs as early as 10 weeks of age, drastically elevating the risk of intervertebral disc disease (IVDD) and spinal cord herniation⁴⁹. Because CDDY is an autosomal dominant trait, even a single copy (heterozygosity) increases the probability of surgical disc herniation by a factor of 5 to 15⁴⁹.

Genomic Variant	Locus	Phenotypic Expression	Clinical Implications
FGF4L1 Retrogene	CFA18	Chondrodysplasia (CDPA): Shortened limbs, carpal valgus ⁵⁰ .	Breed-defining standard for Dachshunds, Corgis; not associated with elevated IVDD risk ⁵⁰ .
FGF4L2 Retrogene	CFA12	Chondrodystrophy (CDDY): Shortened limbs, increased head width ⁵⁰ .	Premature calcification of discs, highly elevated risk of IVDD and spinal herniation ⁴⁹ .

RSPO2 Mutation	Varies	Furnishings / Improper Coat (IC) ⁵³ .	Determines presence of facial furnishings (mustache/eyebrows) and wire hair texture; autosomal dominant ⁵³ .
WBS Orthologs	Varies	Hypersociality, extreme gregariousness ⁵⁴ .	Neurological obligation to seek interspecies social bonds, mirroring human Williams-Beuren syndrome ⁵⁴ .

Structural Variants and Hypersociality

Canine neurogenetics also explains the species' profound orientation toward human affection. Recent comparative genomic studies reveal that the hypersocial behavior of domestic dogs is strictly linked to structural variants in genomic regions orthologous to those causing Williams-Beuren syndrome (WBS) in humans⁵⁴. In humans, WBS is a rare multisystem genetic disorder characterized by extreme, indiscriminate gregariousness and hyper-sociality. The selective breeding of canines over millennia inadvertently selected for identical genomic deletions and insertions within these specific loci. Consequently, the domestic dog is neurologically obligated by its very genome to seek out interspecies social bonds⁵⁴. The vast morphological and behavioral tapestry of the modern dog underscores a reality in which *Canis familiaris* was not merely domesticated by humanity in the wake of environmental cataclysms, but rather structurally engineered by—and concurrently engineered the neurophysiology of—*Homo sapiens*, resulting in an obligate, symbiotic existence.

Works cited

1. Behavioral and emotional co-modulation during dog–owner interaction measured by heart rate variability and activity - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11502769/>
2. Assessing the Relationship Between Emotional States of Dogs and Their Human Handlers, Using Simultaneous Behavioral and Cardiac Measures - Frontiers, <https://www.frontiersin.org/journals/veterinary-science/articles/10.3389/fvets.2022.897287/full>
3. Dogs and owners share emotional and physiological changes - Harvard Health, <https://www.health.harvard.edu/heart-health/dogs-and-owners-share-emotional-and-physiological-changes>
4. The heart rate variability of a dog and its owner are interconnected during

interaction, and it indicates the state of the autonomic nervous system. The owner's high heart rate variability was connected to the dog's high heart rate variability, and vice versa : r/science - Reddit,
https://www.reddit.com/r/science/comments/1gl2lrm/the_heart_rate_variability_of_a_dog_and_its_owner/

5. (PDF) Behavioral and emotional co-modulation during dog–owner interaction measured by heart rate variability and activity - ResearchGate,
https://www.researchgate.net/publication/385216357_Behavioral_and_emotional_co-modulation_during_dog-owner_interaction_measured_by_heart_rate_variability_and_activity
6. A Healing Bond: How Dog-Human Synchrony Enhances Co-Regulation | Dr. Leslie Korn,
<https://drlesliekorn.com/research/dog-human-synchrony-mental-health/>
7. Psychosocial and Psychophysiological Effects of Human-Animal Interactions: The Possible Role of Oxytocin - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3408111/>
8. Evidence for a synchronization of hormonal states between humans and dogs during competition - DigitalCommons@UNO,
<https://digitalcommons.unomaha.edu/cgi/viewcontent.cgi?referer=&httpsredir=1&article=1141&context=psychfacpub>
9. Oxytocin promotes social bonding in dogs - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4078815/>
10. Oxytocin release patterns and effects, with a focus on touch, social interaction, and closeness - Frontiers,
<https://www.frontiersin.org/journals/psychology/articles/10.3389/fpsyg.2026.1771799/full>
11. Dog Gazes Hijack the Brain's Maternal Bonding System - Smithsonian Magazine,
<https://www.smithsonianmag.com/science-nature/dog-gazes-hijack-brains-maternal-bonding-system-180955019/>
12. Behavior 102 For Animal Sanctuaries: Communication,
<https://opensanctuary.org/animal-behavior-communication/>
13. Why do most mammals find being stroked/patted pleasurable? : r/askscience - Reddit,
https://www.reddit.com/r/askscience/comments/28mxt9/why_do_most_mammals_find_being_strokedpatted/
14. The right touch: Stroking of CT-innervated skin promotes vocal emotion processing - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC5709431/>
15. Calming Effects of Touch in Human, Animal, and Robotic Interaction—Scientific State-of-the-Art and Technical Advances - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7672023/>
16. The Neurobiology Shaping Affective Touch: Expectation, Motivation, and Meaning in the Multisensory Context - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4701942/>
17. Oxytocin release patterns and effects, with a focus on touch, social interaction, and closeness - Frontiers,

- <https://www.frontiersin.org/journals/psychology/articles/10.3389/fpsyg.2026.1771799/pdf>
18. Hearts in Sync: The Bond Between Dogs and Humans - Warrior Canine Connection,
<https://warriorcanineconnection.org/hearts-in-sync-the-bond-between-dogs-and-humans/>
 19. The Role of Oxytocin in the Dog–Owner Relationship - Max Planck Institute for Evolutionary Anthropology,
https://www.eva.mpg.de/documents/MDPI/Marshall-Pescini_The-role_Animals_2019_3179284.pdf
 20. Odors activate dual pathways, a TRPC2 and a AA-dependent pathway, in mouse vomeronasal neurons - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2867386/>
 21. Your dog's amazing Jacobson Organ – and how it enhances their super power of 'SMELL' and maybe even taste., <https://healthydogtreats.com.au/jacobson-organ/>
 22. Vomeronasal organ - Wikipedia, https://en.wikipedia.org/wiki/Vomeronasal_organ
 23. The dog vomeronasal organ: A review - ResearchGate,
https://www.researchgate.net/publication/325381241_The_dog_vomeronasal_organ_A_review
 24. The dog vomeronasal organ: a review - Dog behavior,
<https://dogbehavior.it/dogbehavior/article/view/27>
 25. Odors activate dual pathways, a TRPC2 and a AA-dependent pathway, in mouse vomeronasal neurons - PubMed, <https://pubmed.ncbi.nlm.nih.gov/20147653/>
 26. TRICK or TRP? What Trpc2–/– mice tell us about vomeronasal organ mediated innate behaviors - Frontiers,
<https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2015.00221/full>
 27. Odor-Based Recognition of Canine Stress: Investigating Human Olfactory Perception - MPG.PuRe,
https://pure.mpg.de/rest/items/item_3628961_3/component/file_3628962/content
 28. Dog Brains Differentiate Human Anger, Fear, and Sadness - Neuroscience News,
<https://neurosciencenews.com/dog-brains-human-emotion-31202/>
 29. The dog vomeronasal organ: a review - Semantic Scholar,
<https://www.semanticscholar.org/paper/The-dog-vomeronasal-organ%3A-a-review-Coli-Stornelli/dcf7270c1043c6dad06f2e7f5c178084f9ace01>
 30. [PDF] Can animals detect when their owners are returning home,
<https://www.semanticscholar.org/paper/d90c9ffadc13b4dda08d3ac0da4b7b9033bc476e>
 31. Magnetoreception - Wikipedia, <https://en.wikipedia.org/wiki/Magnetoreception>
 32. Cry1* label in the S1 cones of the dog and orangutan retina but not in... | Download Scientific Diagram - ResearchGate,
https://www.researchgate.net/figure/Cry1-label-in-the-S1-cones-of-the-dog-and-orangutan-retina-but-not-in-the-mouse-retina_fig2_295688050
 33. Magnetoreception in the wood mouse (*Apodemus sylvaticus*): influence of weak frequency-modulated radio frequency fields - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4413948/>

34. Magnetoreception molecule found in the eyes of dogs and primates,
<https://maxplanckneuroscience.org/magnetoreception-molecule-found-in-the-eyes-of-dogs-and-primates/>
35. Magnetoreception molecule found in the eyes of dogs and primates,
<https://www.mpg.de/10319591/magnetoreception-eye>
36. Magnetoreception molecule found in the eyes of dogs, primates - ScienceDaily,
<https://www.sciencedaily.com/releases/2016/02/160225105221.htm>
37. (PDF) The canine vibrissal system as a highly innervated and functional sensory organ,
https://www.researchgate.net/publication/389915531_The_canine_vibrissal_system_as_a_highly_innervated_and_functional_sensory organ
38. Kóryos - Wikipedia, entziklopedia askea.,
<https://eu.wikipedia.org/wiki/K%C3%B3ryos>
39. Kóryos - Wikipédia, <https://fr.wikipedia.org/wiki/K%C3%B3ryos>
40. The call of the myth - Pet Connection Veterinary Clinic,
<https://petconnection.ae/the-call-of-the-myth/>
41. Männerbund - Wikipedia, <https://en.wikipedia.org/wiki/M%C3%A4nnerbund>
42. The Trinity and the Three-Tiered Structure in Proto-Indo-European Organizations - researchmap,
https://researchmap.jp/read0026230/published_papers/32050497/attachment_file.pdf
43. The Horse, the Wheel, and Language: How Bronze-Age Riders from the Eurasian Steppes Shaped the Modern World [Course Book ed.] 9781400831104 - DOKUMEN.PUB,
<https://dokumen.pub/the-horse-the-wheel-and-language-how-bronze-age-riders-from-the-eurasian-steppes-shaped-the-modern-world-course-booknbsped-9781400831104.html>
44. The Paippalādasamhitā of the Atharvaveda : a new critical edition of the three 'new' Anuvākas of Kāṇḍa 17 with English translation and commentary - Scholarly Publications Leiden University,
<https://scholarlypublications.universiteitleiden.nl/access/item%3A2979360/view>
45. An expressed fgf4 retrogene is associated with breed-defining chondrodysplasia in domestic dogs - Princeton University,
<https://collaborate.princeton.edu/en/publications/an-expressed-fgf4-retrogene-is-associated-with-breed-defining-cho/>
46. An expressed fgf4 retrogene is associated with breed-defining chondrodysplasia in domestic dogs - PubMed, <https://pubmed.ncbi.nlm.nih.gov/19608863/>
47. An Expressed Fgf4 Retrogene Is Associated with Breed-Defining Chondrodysplasia in Domestic Dogs - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC2748762/>
48. OMIA:002542-9615: Chondrodysplasia, FGF4 retrogene-related in Canis lupus familiaris (dog) - Online Mendelian Inheritance in Animals,
<https://omia.org/OMIA002542/9615/>
49. Managing the Genetics of Chondrodystrophy (CDDY) - Veterinary Genetics Laboratory - UC Davis,

- <https://vgl.ucdavis.edu/news/managing-genetics-chondrodystrophy>
50. Chondrodysplasia, chondrodystrophy, and hereditary increased risk of intervertebral disc disease - TGH-Hundegesundheit,
<https://www.tgh-hundegesundheit.de/en/2026/04/15/chondrodysplasia-chondrodystrophy-and-hereditary-increased-risk-of-intervertebral-disc-disease/>
 51. The Effects of FGF4 Retrogenes on Canine Morphology - MDPI,
<https://www.mdpi.com/2073-4425/13/2/325>
 52. Phenotypic Effects of FGF4 Retrogenes on Intervertebral Disc Disease in Dogs - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6627552/>
 53. Furnishings-Improper Coat Locus (FURN-IC LOCUS) - GenSol Diagnostics,
<https://www.gensoldx.com/tests/furnishings-improper-coat-locus-furn-ic-locus/>
 54. Decoding the Canine Mind - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC7192336/>