

# HYDROGEN DIPOLE MATRIX ELEMENTS FROM BICONE GEOMETRY WITH AN $\alpha$ -SENSITIVITY DETERMINATION VIA $SO(4, 2)$

JEREMY EBERT

**ABSTRACT.** This note is pedagogical: an elegant Schwinger-boson packaging of known  $SO(4)/SO(4, 2)$  hydrogen results, not a claim of new physics. The  $SO(4)$  dynamical symmetry (Pauli, Fock) and its  $SO(4, 2)$  extension with parabolic-coordinate dipole matrix elements (Barut–Kleinert) are 1920s–1970s material; our contribution is the “bicone” geometric framing — two Schwinger  $su(2)$  cones glued at a common  $T_0$ -circle — and the observation that the parabolic basis  $|n_1 n_2 m\rangle$  is this construction’s *native* Fock basis (simultaneous  $(A_z, L_z)$  eigenstates), so no differential equation is solved for the states.

Within this framing, the hydrogen  $2p \rightarrow 1s$  dipole  $|\langle R_{10}|r|R_{21}\rangle|/a_0 = 768/(243\sqrt{6})$  follows from the bicone selection rule ( $|\langle R_{10}|r|R_{21}\rangle| = \sqrt{3}|C|$ ) and the exact rational  $\langle 000|z|100\rangle = -128/243$ , with no spherical radial solution and no Gordon hypergeometric integral. The geometric dipole  $Y_{\text{geom}} = 768/(243\sqrt{6})$  is  $\alpha$ -independent. Using a *measured* Lyman- $\alpha$  line as empirical input (this is not a priori numerology), elimination of  $a_0$  via  $a_0 = \bar{\lambda}_c/\alpha$  gives the  $\alpha$ -independent  $\alpha = 4\omega_{21}^3 \bar{\lambda}_c^2 Y_{\text{geom}}^2 / (9c^2 A_{21}) \approx 1/137.14$ . The point is conceptual non-circularity, not competitive precision: 0.07% ( $\sim 3$  figures) against the  $\sim 10$ -figure metrology standard. What is shown is that the  $SO(4, 2)$  dipole geometry determines  $\alpha$  from one spectral line with no circular input — a clean classroom demonstration of dynamical symmetry doing real work.

## 1. THE BICONE AND THE DIPOLE FORM

The bicone construction used here is self-contained: two cones glued base-to-base at a common  $T_0$ -circle form a bicone inside the sphere  $X^2 + Y^2 + (T - T_0)^2 = T_0^2$  (Fig. 1). Each cone carries a Schwinger  $su(2)$  (denoted  $J, K$ ); with  $L = J + K$ ,  $A = J - K$ ,

$$[L_i, L_j] = i\epsilon_{ijk}L_k, \quad [L_i, A_j] = i\epsilon_{ijk}A_k, \quad [A_i, A_j] = i\epsilon_{ijk}L_k,$$

the  $so(4)$  of the hydrogen  $n$ -shell. The gluing  $j_1 = j_2$  is the Kustaanheimo–Stiefel constraint geometrized;  $n = 2T_0$ .

Hydrogen states are  $|n; l, m\rangle \leftrightarrow |(j, j); l, m\rangle$ ,  $j = (n - 1)/2$ , via  $|j, m_1\rangle \otimes |j, m_2\rangle$  coupled with  $L = J + K$ . In particular  $|1s\rangle$  is the vacuum and  $|2p, m\rangle$  is the triplet from  $|\frac{1}{2}\rangle \otimes |\frac{1}{2}\rangle$  (Fig. 2).

**Proposition 1.1** (Bicone dipole form). *By explicit ladder algebra  $(a_i|1\rangle = |0\rangle)$ , with  $R_{-1} = a_1 b_1$ ,  $R_0 = (a_1 b_2 + a_2 b_1)/\sqrt{2}$ ,  $R_{+1} = a_2 b_2$ ,*

$$\langle 1s|R_q|2p, m\rangle = \delta_{q, -m} \cdot (+1),$$

*the Wigner–Eckhart vector selection rule, emerging from computation. Writing  $r_q = C(R_q + R_q^\dagger)$  with  $C$  a length,  $SO(3)$  Wigner–Eckhart gives*

$$(1) \quad |\langle R_{10}|r|R_{21}\rangle| = \sqrt{3}|C|.$$

*The  $\sqrt{3}$  and the  $\delta_{q, -m}$  pattern are bicone output.*

*Remark.* The  $ab$ -type bilinears do *not* form proper  $SO(3)$  tensor operators under  $L$  (ladder relations fail; Schwinger operators are dual spinors, not tensors). Hence  $r_q$  is not in the enveloping algebra




FIGURE 1. The bicone in the sphere: two cones glued at the gold  $T_0$ -circle, carrying  $su(2) \oplus su(2) \rightarrow so(4)$ . Four views:  $T$  (top-down, the  $T_0$ -circle),  $X$  and  $Y$  sides, and 3D.

of  $su(2) \oplus su(2)$  in simple bilinear form; the scale  $|C|$  requires the larger  $SO(4, 2)$ , where the dipole sits as a vector generator (Barut).

## 2. THE PARABOLIC $SO(4, 2)$ SCALE

In parabolic coordinates  $x = \sqrt{\xi\eta} \cos \phi$ ,  $y = \sqrt{\xi\eta} \sin \phi$ ,  $z = (\xi - \eta)/2$ ,  $r = (\xi + \eta)/2$  — the  $SO(4, 2)$ -natural basis, also the Stark/parabolic basis — the dipole component  $z = (\xi - \eta)/2$  is *linear* (vs.  $\hat{r}$  in spherical). The integrand is then polynomial  $\times$  exponential, not hypergeometric.

**Proposition 2.1** (Parabolic scale). *In parabolic coordinates  $x = \sqrt{\xi\eta} \cos \phi$ ,  $y = \sqrt{\xi\eta} \sin \phi$ ,  $z = (\xi - \eta)/2$ ,  $r = (\xi + \eta)/2$  — the  $SO(4, 2)$ -natural basis — the dipole component  $z = (\xi - \eta)/2$  is linear (vs.  $\hat{r}$  in spherical). The parabolic states  $|n_1 n_2 m\rangle$  are the simultaneous  $(A_z, L_z)$  eigenbasis*



FIGURE 2. The dipole transition  $|1s\rangle \rightarrow |2p\rangle$  between bicone shells  $n = 1$  ( $T_0 = 1/2$ ) and  $n = 2$  ( $T_0 = 1$ ).

of the bicone, with  $A_z = J_z - K_z$  the Stark operator: they are algebraic states of the cone's  $so(4)$ , whose position-space forms coincide with the Schrödinger parabolic solutions. The integrand is then polynomial $\times$ exponential, not hypergeometric.

$$I_1 = \iint e^{-3(\xi+\eta)/4} (\xi - \eta)(1 - \xi/2)(\xi + \eta) d\xi d\eta = -\frac{2048}{243}$$

by elementary gamma integrals (the  $(\xi^2 - \eta^2)$  term vanishes by  $\xi \leftrightarrow \eta$  symmetry), whence with  $N = 1/(4\sqrt{\pi})$ ,

$$\langle 000|z|100\rangle = \frac{1}{16} \left( -\frac{2048}{243} \right) = -\frac{128}{243},$$

an exact rational. Since  $|2p_z\rangle = (|100\rangle - |010\rangle)/\sqrt{2}$ ,

$$\frac{|C|}{a_0} = \frac{128\sqrt{2}}{243} = \frac{256}{243\sqrt{2}} \approx 0.744936.$$

*Remark (Framing).* No spherical radial solution and no Gordon/hypergeometric integral is used; the matrix element follows in the natural parabolic  $SO(4, 2)$  representation. The parabolic basis  $|n_1 n_2 m\rangle$  is the bicone's  $(A_z, L_z)$  eigenbasis — algebraic states of the cone's  $so(4)$ , not solutions imported from the Schrödinger equation in spherical coordinates. Their position-space forms coincide with the parabolic Schrödinger solutions, but the *basis itself* is cone-native: the whole construction is parabolic slices of the  $SO(4)$  geometry.

Combining (1) with the Proposition,

$$(2) \quad \frac{|\langle R_{10}|r|R_{21}\rangle|}{a_0} = \sqrt{3} \cdot \frac{256}{243\sqrt{2}} = \frac{768}{243\sqrt{6}} \approx 1.290266,$$

the textbook Gordon number, here from bicone $\times$ parabolic- $SO(4, 2)$ .

3. DETERMINING  $\alpha$  WITHOUT  $a_0$  OR  $\epsilon_0$ 

The Lyman- $\alpha$  Einstein coefficient

$$A_{21} = \frac{\omega_{21}^3 |d_{21}|^2}{3\pi\epsilon_0\hbar c^3}, \quad |d_{21}|^2 = e^2 |\langle 1s|\mathbf{r}|2p\rangle|^2,$$

with  $|\langle 1s|\mathbf{r}|2p\rangle|^2 = |C|^2$  (only  $q = -m$  contributes). The  $SO(3)$  Wigner–Eckhart radial-to-vector factor is explicit:

$$|C|^2 = \frac{1}{3} |\langle R_{10}|r|R_{21}\rangle|^2 = \frac{1}{3} Y_{\text{geom}}^2 a_0^2, \quad Y_{\text{geom}} = \frac{768}{243\sqrt{6}}.$$

Hence

$$(3) \quad e^2 = \frac{9\pi\epsilon_0\hbar c^3 A_{21}}{\omega_{21}^3 a_0^2 Y_{\text{geom}}^2},$$

the  $1/3$  now explicit (absorbed in the  $9\pi$ ).

For  $\alpha = e^2/(4\pi\epsilon_0\hbar c)$ , the  $\epsilon_0$  cancels:

$$\alpha = \frac{9c^2 A_{21}}{4\omega_{21}^3 a_0^2 Y_{\text{geom}}^2}.$$

Using  $a_0 = \bar{\lambda}_c/\alpha$  with the reduced Compton wavelength  $\bar{\lambda}_c = \hbar/(m_e c)$  —  $\hbar, c$  exact (SI 2019),  $m_e$  from Penning-trap cyclotron ratios and XRCD silicon spheres, all  $\alpha$ -independent — and solving for  $\alpha$ :

$$(4) \quad \boxed{\alpha = \frac{4\omega_{21}^3 \bar{\lambda}_c^2 Y_{\text{geom}}^2}{9c^2 A_{21}}}$$

No  $a_0$ , no  $\epsilon_0$ , no CODATA global fit, no  $\alpha$  input. Every quantity is either measured ( $\omega_{21}, A_{21}$ : Lyman- $\alpha$  spectroscopy;  $\bar{\lambda}_c$ : Penning trap + XRCD), exact ( $c$ ), or pure  $SO(4, 2)$ /bicone geometry ( $Y_{\text{geom}}$ ).

With  $\omega_{21} = 1.55 \times 10^{16} \text{ s}^{-1}$ ,  $A_{21} = 6.2648 \times 10^8 \text{ s}^{-1}$  (NIST ASD),  $\bar{\lambda}_c = 3.861593 \times 10^{-13} \text{ m}$ :

$$\alpha \approx \frac{1}{137.14} \quad \text{vs. known } 1/137.036,$$

0.07% (limited by  $A_{21}$  rounding). This is a genuine  $\alpha$ -independent determination:  $a_0 \equiv \hbar/(m_e c \alpha)$  cannot be measured without  $\alpha$  by definition, but (4) eliminates it entirely.

## 4. CALIBRATION: WHAT IS DERIVED, WHAT IS MEASURED

- **Derived ( $\alpha$ -independent):** the dipole form  $\sqrt{3}|C|$  (bicone ladder algebra); the scale  $|C|/a_0 = 256/(243\sqrt{2})$  (parabolic  $SO(4, 2)$  gamma integrals); hence  $Y_{\text{geom}} = 768/(243\sqrt{6})$ . The  $\alpha$  determination (4) uses only  $\alpha$ -independent inputs: Lyman- $\alpha$  ( $\omega_{21}, A_{21}$ ), the Compton wavelength  $\bar{\lambda}_c$  (Penning trap + XRCD),  $c$  (exact), and  $Y_{\text{geom}}$  (geometry). No  $a_0$  (which is  $\hbar/(m_e c \alpha)$  by definition), no  $\epsilon_0$ , no CODATA global fit.
- **Not claimed:** a derivation of  $\alpha \approx 1/137$  from pure geometry (impossible by conformal invariance —  $SO(4, 2)$  is scale-invariant by definition, so one measurement is the theoretical minimum); “zero Schrödinger input” in any absolute sense (see Remark above).

The  $SO(4, 2)$  Casimirs in hydrogen’s irrep are pure numbers ( $W_2 = 3$ ,  $W_3 = 0$ ,  $W_4 = -18$ ) and cannot contain  $e^2$ : every  $e^2 > 0$  Coulomb problem realizes the same irrep. What the program supplies is the *form* of every transition; the coupling is then overdetermined by the spectrum — one  $e^2$  must work for all  $Y_{\text{geom}}$  — and fixed by a single line.

## ACKNOWLEDGMENT

Large language model tools were used for numerical verification, figure generation, drafting, and literature search.

## REFERENCES

- [1] A. O. Barut and H. Kleinert, *Dynamical group  $O(4, 2)$  for the hydrogen atom*, Phys. Rev. 156 (1967).
- [2] F. Nendzig, *The Quantum Theory of the Hydrogen Atom*, lecture notes (Apr. 2024), [http://www.fnendzig.de/physics/HydrogenAtom\\_Nendzig.pdf](http://www.fnendzig.de/physics/HydrogenAtom_Nendzig.pdf) (reviewed for  $SO(4, 2)$  generators, Ch. 7).
- [3] W. Gordon, *Der Comptoneffekt nach der Schrödingerschen Theorie*, Z. Phys. 40 (1926) (dipole matrix elements).

FRANKLIN COUNTY, OHIO, USA