

# The Dynamic Slicing Operator on the AM–GM Cone: A Quadric Geometric Framework for Keplerian Orbits and Perturbed Trajectories

Jeremy Ebert

Franklin County, Ohio, USA

---

## Abstract

We present a geometric and operator-theoretic formulation of central-force celestial mechanics using arithmetic mean–geometric mean (AM–GM) coordinates on factor surfaces. By mapping the turning points of an orbit (apoapsis  $r_a$  and periapsis  $r_p$ ) to mean-value coordinates  $T = (r_a + r_p)/2$ ,  $X = (r_a - r_p)/2$ , and  $Y = \sqrt{r_a r_p}$ , two-body Keplerian trajectories are represented as planar intersections of the Minkowski-like quadric cone  $\mathcal{C} : X^2 + Y^2 = T^2$ . We show that total energy  $E$ , specific angular momentum  $h$ , and central mass  $M$  isolate cleanly along orthogonal geometric directions of the cone. Furthermore, time evolution and non-Keplerian perturbations (such as drag or thrust) are modeled via a dynamic slicing operator  $\hat{S}(t)$  whose movement through  $\mathcal{C}$  parameterizes osculating orbit transitions, orbital decay, and energy pumps under the action of the  $\mathfrak{so}(2, 1)$  Lie algebra.

*Keywords:* Kepler problem, AM–GM coordinates, Quadric cone,  $\mathfrak{so}(2, 1)$  Lie algebra, Dynamic slicing operator, Orbital perturbations

---

## 1. Introduction

Classical two-body motion under a  $1/r^2$  central force potential is conventionally parameterized via spatial state vectors  $(\mathbf{r}, \mathbf{v})$  or osculating Keplerian orbital elements  $(a, e, i, \omega, \Omega, \nu)$ . While these representations excel at numerical integration, phase space transitions between bound ( $E < 0$ ), parabolic ( $E = 0$ ), and hyperbolic ( $E > 0$ ) regimes often require piecewise analytic parameterizations involving distinct trigonometric or hyperbolic anomalies.

In recent work on factor lattice geometry [?], an arithmetic mean–geometric mean coordinate transformation was introduced to organize binary factor pairs onto the quadric surface  $X^2 + Y^2 = T^2$ . In this paper, we extend this

construction to continuous classical dynamics. By identifying factor pairs with radial turning points  $(r_a, r_p)$ , we demonstrate that all Keplerian trajectories manifest as plane slices of a single quadric cone, and that time evolution under external forces can be formalized as a continuous family of slicing operators  $\hat{S}(t)$ .

## 2. The Orbital AM–GM Mapping

Consider a particle of mass  $m$  orbiting a central mass  $M$  in a Newtonian gravitational potential  $V(r) = -\mu/r$ , where  $\mu = GM$ . Let  $r_a$  and  $r_p$  denote the apoapsis and periapsis distances, respectively.

**Definition 1** (Orbital Mean-Value Mapping). *The turning points  $(r_a, r_p)$  map to the coordi-*

nate tuple  $(X, Y, T) \in \mathbb{R}^3$  via:

$$T = \frac{r_a + r_p}{2}, \quad X = \frac{r_a - r_p}{2}, \quad Y = \sqrt{r_a r_p} \quad (1)$$

Direct substitution yields the fundamental algebraic identity:

$$X^2 + Y^2 = \left(\frac{r_a - r_p}{2}\right)^2 + r_a r_p = \left(\frac{r_a + r_p}{2}\right)^2 = T^2 \quad (2)$$

Thus, every physically realization of radial turning points lies on the cone surface  $\mathcal{C} : X^2 + Y^2 = T^2$ . The physical interpretations of these coordinates are immediately evident:

- $T = a$ : Semi-major axis (axial scale).
- $Y = b = a\sqrt{1 - e^2}$ : Semi-minor axis (transverse radius).
- $X = c = ae$ : Focal displacement (eccentricity displacement).

The cone constraint Equation (??) is geometrically equivalent to the standard ellipse relation  $(ae)^2 + b^2 = a^2$ .

### 3. Physical Invariants as Cone Coordinates

A major advantage of the AM–GM representation is that scalar physical invariants isolate along principal geometric axes.

**Proposition 1** (Isolation of Physical Invariants). *For a two-body orbit on the cone  $\mathcal{C}$ :*

1. **Total Specific Energy  $E$** : Is determined purely by the axial height  $T$ :

$$E = -\frac{\mu}{2T} \quad (3)$$

*Horizontal slices  $T = \text{const.}$  represent constant-energy manifolds.*

2. **Specific Angular Momentum  $h$** : Is determined by the transverse coordinate  $Y$  and axial height  $T$ :

$$h = \sqrt{\mu \frac{Y^2}{T}} = \sqrt{\frac{\mu b^2}{a}} \quad (4)$$

*Hyperbolic cross-sections  $Y = \text{const.}$  define manifolds of constant angular momentum.*

3. **Central Mass Extraction  $M$** : Let  $v_a$  and  $v_p$  be the orbital speeds at apoapsis and periapsis. By angular momentum conservation ( $h = v_a r_a = v_p r_p$ ), the central parameter  $\mu = GM$  satisfies:

$$\mu = GM = T \cdot v_a v_p \implies M = \frac{T \cdot v_a v_p}{G} \quad (5)$$

### 4. Trajectory Classification via Theorem 3.1

In [?], it was proved that linear constraints  $ar_a + br_p = c$  on factor space trace conic sections on  $\mathcal{C}$ . Applying this theorem to celestial mechanics provides a unified classification of energy states.

**Theorem 2** (General Orbit Classification). *Let a family of orbital states satisfy the linear constraint  $ar_a + br_p = c$  for  $a, b, c \in \mathbb{R}$ . The resulting intersection with the cone  $\mathcal{C}$  yields:*

1. **Elliptic / Bound States** ( $ab > 0$ ): Slices both generators; forms a closed ellipse on  $\mathcal{C}$  corresponding to  $E < 0$  ( $0 \leq e < 1$ ).
2. **Parabolic / Escape States** ( $ab = 0$ ): Slices parallel to a boundary generator ray ( $T = \pm X$ ), representing  $E = 0$  ( $e = 1$ ).
3. **Hyperbolic / Unbound States** ( $ab < 0$ ): Slices both upper and lower nappes of  $\mathcal{C}$ , producing a hyperbola corresponding to  $E > 0$  ( $e > 1$ ).

### 5. The Dynamic Slicing Operator $\hat{S}(t)$

To model time-dependent orbital dynamics, orbital decay, or impulsive burns, we define the *Dynamic Slicing Operator*.

**Definition 2** (Dynamic Slicing Operator). *Let  $\hat{S}(t)$  be a time-dependent linear operator defining a cutting plane in  $\mathbb{R}^3$ :*

$$\hat{S}(t) : \quad \alpha(t)X + \beta(t)T = \gamma(t) \quad (6)$$

The instantaneous orbital state  $\mathcal{O}(t)$  is given by the intersection:

$$\mathcal{O}(t) = \mathcal{C} \cap \hat{S}(t) \quad (7)$$

As time  $t$  advances, the trajectory of the operator plane  $\hat{S}(t)$  generates distinct dynamical behaviors:

- **Vertical Translation** ( $\dot{\gamma} \neq 0$ ): Models continuous energy exchange. Downward motion ( $\dot{\gamma} < 0$ ) describes atmospheric drag or orbital decay toward the vertex  $(0, 0, 0)$ . Upward motion ( $\dot{\gamma} > 0$ ) models low-thrust expansion.
- **Operator Tilting** ( $\dot{\alpha} \neq 0, \dot{\beta} \neq 0$ ): Models eccentricity pumping at constant energy. A plane tilting toward the critical slope  $\alpha = \beta$  drives a bound orbit toward parabolic escape ( $e \rightarrow 1$ ).

## 6. $\mathfrak{so}(2, 1)$ Lie-Algebraic Generators

The isometry group preserving the cone  $\mathcal{C}$  is  $SO(2, 1)$ . The Lie algebra  $\mathfrak{so}(2, 1)$  is spanned by the rotation generator  $L$  and boost generators  $B_X, B_Y$ :

$$B_X = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad B_Y = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad (8)$$

**Proposition 3** (Isomeric Orbital Boosts). *An action of the boost group  $\exp(sB_X)$  on an initial state  $(X_0, Y_0, T_0)$  yields:*

$$\begin{pmatrix} X(s) \\ T(s) \end{pmatrix} = \begin{pmatrix} \cosh s & \sinh s \\ \sinh s & \cosh s \end{pmatrix} \begin{pmatrix} X_0 \\ T_0 \end{pmatrix} \quad (9)$$

This boost preserves  $Y = Y_0 = b$ , leaving the specific angular momentum  $h$  invariant while scaling the semi-major axis  $a(s) = Y_0 \cosh(s)$  and eccentricity  $e(s) = \tanh(s)$ . The rapidity parameter  $s = \frac{1}{2} \ln(r_a/r_p)$  measures the logarithmic radial asymmetry.

Analytic continuation of the rapidity parameter to imaginary values ( $s \rightarrow i\phi$ ) transforms the hyperbolic boost into the trigonometric variation  $T(\phi) = Y_0 \cos(\phi)$ , mapping unbound scattering trajectories directly into periodic eccentric anomalies of bound states.

## 7. Conclusion

The AM–GM conic representation provides a rigorous, computationally linear, and visually unified framework for celestial mechanics. By parameterizing orbits via turning-point mean values  $(X, Y, T)$ , physical energy states, angular momentum, and central masses isolate into direct spatial dimensions on  $X^2 + Y^2 = T^2$ . The dynamic slicing operator  $\hat{S}(t)$  offers a novel operator-theoretic formulation for modeling perturbed trajectories, continuous burns, and relativistic precessions on a single quadric manifold.

## References

- [1] J. Ebert, *AM–GM Coordinates on Integer Factor Pairs: A Conic Classification Theorem and Rigidity of the Divisor Lattice*, Preprint (2026).
- [2] A. E. Roy, *Orbital Motion*, 4th ed., CRC Press, Boca Raton, FL (2004).
- [3] H. Goldstein, C. Poole, and J. Safko, *Classical Mechanics*, 3rd ed., Addison-Wesley (2002).

## Acknowledgment

Large language model tools were used for drafting, figure preparation, and checking.