

TOTIENT HYPERBOLAS AND THE $V_4 \leftrightarrow QR(12)$ BIJECTION A COMPANION NOTE TO PAPER A

JEREMY EBERT

ABSTRACT. Paper A [1] classifies the conic sections cut from the factor cone $X^2 + Y^2 = T^2$ by lines $ax + by = c$ in factor coordinates. For the units $r \in U(12) = \{1, 5, 7, 11\}$, the sections $X = (r - 1)/2$ are hyperbolas passing through the unit-shell points $(\pm(r - 1)/2, \pm\sqrt{r}, (r + 1)/2)$. Their vertices $X \in \{0, 2, 3, 5\}$ and their landing-circle radii $T \in \{1, 3, 4, 6\}$ square to $QR(12) = \{0, 1, 4, 9\}$ modulo 12, giving two bijections $U(12) \rightarrow QR(12)$. The two bijections are complementary, $\varphi_+ + \varphi_- \equiv 1 \pmod{12}$, and both preserve the genus bipartition $\{1, 11\} \mapsto \{0, 1\}$, $\{5, 7\} \mapsto \{4, 9\}$. This note records the construction and its exact scope.

1. PAPER-A SECTIONS

We use Paper A's two-sided lift

$$X = \frac{x - y}{2}, \quad Y = \pm\sqrt{xy}, \quad T = \frac{x + y}{2}, \quad X^2 + Y^2 = T^2.$$

A line $ax + by = c$ with $ab < 0$ and $c \neq 0$ cuts a nondegenerate hyperbola; for $c = 0$ the section degenerates to the generator pair [1]. The lines

$$x - y = 2c, \quad \text{i.e. } a = 1, \ b = -1,$$

give the vertical planes $X = c$, whose sections are

$$T^2 - Y^2 = c^2,$$

hyperbolas opening along T (degenerate for $c = 0$).

2. THE UNIT SHELL

The units modulo 12 are

$$U(12) = \{1, 5, 7, 11\},$$

and since every element squares to 1 mod 12,

$$U(12) \cong V_4,$$

the Klein four-group. The factor pairs $(r, 1)$ and $(1, r)$ lift to the unit-shell points

$$(1) \quad T_r = \frac{r + 1}{2}, \quad X_r = \pm\frac{r - 1}{2}, \quad Y_r = \pm\sqrt{r},$$

so the four labels occupy the heights

$$T \in \{1, 3, 4, 6\}.$$

3. THE FOUR TOTIENT HYPERBOLAS

Definition 3.1. For $r \in U(12)$, the totient hyperbola H_r is the Paper-A section $X = (r - 1)/2$ (the $+$ branch; the $-$ branch is its image under factor exchange $(x, y) \leftrightarrow (y, x)$).

Proposition 3.2. On the cone, H_r is the hyperbola

$$T^2 - Y^2 = \left(\frac{r-1}{2}\right)^2,$$

nondegenerate for $r > 1$ and the degenerate generator pair $T = \pm Y$ for $r = 1$. It contains the shell points $(\pm(r-1)/2, \pm\sqrt{r}, (r+1)/2)$.

Proof. Intersecting $X = (r-1)/2$ with $X^2 + Y^2 = T^2$ gives the displayed equation directly. At $T = (r+1)/2$,

$$Y^2 = \left(\frac{r+1}{2}\right)^2 - \left(\frac{r-1}{2}\right)^2 = r,$$

so the shell points (1) lie on H_r . □

Figure 1 shows the four hyperbolas. Each colored arc is drawn from its vertex (open circle) to its shell points (squares): the vertex is where φ_- is read off, the landing circle where φ_+ is read off.

4. THE DOUBLE $V_4 \leftrightarrow QR(12)$ BIJECTION

The quadratic residues modulo 12 are

$$QR(12) = \{0, 1, 4, 9\}.$$

Each totient hyperbola determines two canonical integers: its vertex $X = (r-1)/2$ (closest approach to the T -axis) and its landing-circle radius $T = (r+1)/2$ (through the shell points).

Proposition 4.1. The maps

$$\varphi_-(r) = \left(\frac{r-1}{2}\right)^2 \bmod 12, \quad \varphi_+(r) = \left(\frac{r+1}{2}\right)^2 \bmod 12,$$

are bijections $U(12) \rightarrow QR(12)$.

Proof. Direct verification:

r	$(r-1)/2$	$\varphi_-(r)$	$(r+1)/2$	$\varphi_+(r)$
1	0	0	1	1
5	2	4	3	9
7	3	9	4	4
11	5	1	6	0

Both image columns are $\{0, 1, 4, 9\} = QR(12)$ with all values distinct. □

Thus the hyperbola bridges $V_4 \leftrightarrow QR(12)$ twice: once at its vertex, once on its landing circle. The cone's quadratic form is the mechanism: the half-integer offsets $(r \pm 1)/2$ become squares upon squaring, and modulo 12 those squares are exactly the quadratic residues.

The two bijections are not independent. Their sum is constant:

Proposition 4.2. For every $r \in U(12)$,

$$\varphi_+(r) + \varphi_-(r) \equiv 1 \pmod{12},$$

i.e. $\varphi_+ = (1 - \cdot) \circ \varphi_-$: vertex-square plus landing-square is 1.

Proof. As integers, $\varphi_+(r) + \varphi_-(r) = ((r+1)/2)^2 + ((r-1)/2)^2 = (r^2 + 1)/2$. Every unit mod 12 is odd and prime to 3, hence $r^2 \equiv 1 \pmod{8}$ and $r^2 \equiv 1 \pmod{3}$, so $r^2 \equiv 1 \pmod{24}$ and $(r^2 + 1)/2 \equiv 1 \pmod{12}$. $\square$

The bijections also respect the genus bipartition. Let $\chi(r) = (\frac{12}{r})$ be the Kronecker genus character; on $U(12)$ it is $+1$ on $\{1, 11\}$ and -1 on $\{5, 7\}$.

Proposition 4.3. *Both φ_- and φ_+ send*

$$\{1, 11\} \mapsto \{0, 1\}, \quad \{5, 7\} \mapsto \{4, 9\}.$$

Moreover $QR(12)$ is stable under $x \mapsto 1 - x$, with orbits $\{0, 1\}$ and $\{4, 9\}$, and $\varphi_+ = (1 - \cdot) \circ \varphi_-$ swaps within each orbit.

Proof. For $r = 12k \pm 1$, $((r-1)/2)^2 \equiv 0, 1 \pmod{12}$; for $r = 12k \pm 5$, $((r-1)/2)^2 \equiv 4, 9 \pmod{12}$, by direct expansion. Since $\chi(r) = +1 \iff r \equiv \pm 1 \pmod{12}$ for $r \in U(12)$, φ_- respects the partition, and $\varphi_+ = 1 - \varphi_-$ does too. $\square$

Remark. The character χ is splitting in the discriminant-12 field: $\chi(r) = +1$ exactly when r splits in $\mathbf{Q}(\sqrt{3})$ ($\{1, 11\}$ split; $\{5, 7\}$ are inert). So each $\sqrt{r}$ ray, defined over $\mathbf{Q}(\sqrt{r})$, carries the $\mathbf{Q}(\sqrt{3})$ -splitting label of r — its genus — and both totient bijections preserve it.




FIGURE 1. The four totient hyperbolas, four views of one object: (a) down the T axis, (b) down Y , (c) down X , (d) perspective. For each unit $r \in \{1, 5, 7, 11\}$ the colored arc is the Paper-A hyperbolic section $X = (r-1)/2$ (yellow $r = 11$ to red $r = 1$, degenerate), running from the open-circle vertex $((r-1)/2, 0, (r-1)/2)$ to the unit-shell squares $(\pm(r-1)/2, \pm\sqrt{r}, (r+1)/2)$ on the landing circles $T = (r+1)/2$ (thin gray); black connectors join the shell points at constant $Y = \sqrt{r}$. The vertices $X \in \{0, 2, 3, 5\}$ and the radii $T \in \{1, 3, 4, 6\}$ square to $QR(12) = \{0, 1, 4, 9\} \pmod{12}$, the two bijections of Proposition 4.1.

5. SCOPE

The placement realizes the bijection; it does not define the group law of $U(12)$, whose multiplication is prior arithmetic. The content of Proposition 4.1 is exact finite arithmetic read off the cone geometry, in the same spirit as Paper A's exact conic classification.

Remark. The author views this double bridge — the same four units read off twice, once at the hyperbola vertices and once on the landing circles — as the geometric form of mod-12 uniqueness in the cone project: the reason the number 12, and not a neighboring integer, is the one whose unit shell closes exactly onto its own quadratic residues. It is also the picture the project grew out of: the unit shell came first, and the discriminant-12 program was built outward from it.

ACKNOWLEDGMENT

Large language model tools were used for machine verification, figure preparation, and drafting.

REFERENCES

- [1] J. Ebert, *The Cutting Plane: Every Line Through the Multiplication Table Is a Conic Section*, Preprint (2026).

FRANKLIN COUNTY, OHIO, USA