

THE CONE PROGRAM: A ROADMAP

JEREMY EBERT

ABSTRACT. The discriminant-12 program studies a single geometric carrier — the cone $T^2 = X^2 + Y^2$, $T \geq 0$ — which supports, without modification, readings from elementary arithmetic, finite group theory, Lorentzian geometry, bosonic quantum mechanics, and analytic number theory. This note maps those readings: what each paper in the series establishes, how the pieces sit on the one cone, and what remains honestly open. It proves nothing new; its purpose is to let a reader see the whole picture accurately, without overstatement.

1. ONE CARRIER

Everything in this program lives on one object: the cone

$$T^2 = X^2 + Y^2, \quad T \geq 0,$$

with the AM–GM parametrization, for $x, y \geq 0$,

$$T = \frac{x+y}{2}, \quad X = \frac{x-y}{2}, \quad Y = \sqrt{xy}.$$

The multiplication table sits in the plane $Y = 0$; its lift to the cone is the substrate of the entire program. Figure 1 shows the readings described below sitting on the one carrier.

The program’s central claim, stated with care: each structure described below is *definable on this cone* — no gluing, no extra dimensions, no deformation of the carrier. The cone carries them all. What is *not* claimed is that these readings are identified with one another. The Lorentz group acting on the cone is the Lorentz group of relativity; the $\mathfrak{su}(2)$ on the fixed- T rings is Schwinger’s $\mathfrak{su}(2)$; the mod-12 arithmetic of the unit shell is mod-12 arithmetic. The unity is in the carrier, not in a claimed identification of the theories. Every paper in the series maintains this boundary, and this roadmap does too.

2. THE ARITHMETIC BASE

The program begins, in [1], with the most elementary object available: the multiplication table. Linear conditions $ax + by = c$ in factor coordinates cut the cone in complete conic sections; constant products sit at reflected

Date: October 5, 2026.




FIGURE 1. Schematic map of the program's readings on the one cone: the divisor-summatory base, the unit shell with its totient hyperbolas, a fixed- T $SU(2)$ ring, a fixed- X $SU(1,1)$ branch, and the screw curve running toward the analytic summit. Schematic only; see the cited papers for the exact statements.

Y -levels with a common Lorentz-type side view; the divisor summatory function

$$D(n) = \sum_{k \leq n} d(k)$$

counts lattice points under the hyperbolas $xy \leq n$; the summatory region decomposes into diagonal gnomons. A tangent construction attaches to each factor level a unique anti-diagonal circle of radius $u/2$. The paper stops where geometry stops: no arithmetic or operator identification is imported into the elementary proofs.

3. THE UNIT SHELL: WHERE THE PROGRAM STARTED

The whole discriminant-12 idea grew out of a single picture, recorded for the first time in [2]. For the units $r \in U(12) = \{1, 5, 7, 11\}$, the Paper-A sections $X = (r-1)/2$ are hyperbolas $T^2 - Y^2 = ((r-1)/2)^2$ passing through the unit-shell points $(\pm(r-1)/2, \pm\sqrt{r}, (r+1)/2)$. Their vertices $\{0, 2, 3, 5\}$ and their landing-circle radii $\{1, 3, 4, 6\}$ both square to $QR(12) = \{0, 1, 4, 9\}$ modulo 12 — two bijections $U(12) \rightarrow QR(12)$, one read at the vertices, one on the landing circles. The picture is exact, and it predates everything else in the program: the unit shell came first, and the cone program was built outward from it.

4. THE LORENTZIAN READING

The cone $T^2 = X^2 + Y^2$ is the light cone of $(2+1)$ -dimensional Minkowski space, and the Lorentz group $SO(2, 1)$ acts on it exactly as the relativity group acts on spacetime — this is the same group on the same quadric, not an analogy. [3] classifies the signed cone cuts as Lorentz orbits and constructs eigen-coordinates and power maps, separating intrinsic projective self-maps of normalized orbit coordinates from equation-dependent level-raising maps. The fixed- X hyperbolic sections of the preceding section are the hyperbolic orbits of this action.

5. THE QUANTUM READING

Fix T : the circles $X^2 + Y^2 = \text{const}$ carry Schwinger's $\mathfrak{su}(2)$ — two bosonic modes with total number conserved and spin $j = T - \frac{1}{2}$. Fix X : the hyperbola branches carry the two-mode $\mathfrak{su}(1, 1)$ squeezing algebra, with number difference conserved and Bargmann index $k = (|d| + 1)/2$ for difference d . [4] realizes both algebras on the cone with Fock-state geometry and ladder operators, and proves the parity selection rule and the operator revival identity for elliptic cuts. [8] completes the picture at the Casimir level: edge-centered spinor data, the primitive null diamond, and a four-part taxonomy — Casimir completion, null-edge geometry, cyclotomic normalization, Boolean return — that keeps the program's distinct V_4 occurrences carefully apart.

A remark on loop quantum gravity, kept proportionate: the kinematic variables of loop quantum gravity are $SU(2)$ holonomies, and its state space is built from $SU(2)$ spin networks. The $SU(2)$ on the cone's fixed- T rings is the same $SU(2)$ — the same representation theory, the same Casimir $j(j+1)$. That is a shared kinematics, noted here as a direction of travel, not a result. Nothing about quantum gravity is claimed.

6. THE ANALYTIC READING: TOWARD THE ZEROS

The program's summit is analytic: the zeros of the Riemann zeta function. Two threads live on the cone here, at different levels of maturity.

The sieve flow, recorded in [9], is numerical exploration: the primes $\leq \sqrt{N}$ generate the exact prime/composite bitmap on $[2, N]$ by the sieve of Eratosthenes, and the zero spectrum reappears as fixed structure under the sieve lift at every scale, with no Riemann-hypothesis input. It is a measurement, not a theorem, and is presented as one.

The screw thread is the program's main line: a screw function overlaid on the cone, and the zeta function itself overlaid on the cone, pursued through the discriminant-12 ledger toward a Hilbert–Pólya-type operator with the positivity certificate as the gate. The ledger records both progress and honest obstructions: the continuum formulation cannot be closed unconditionally — it is RH-hard — while the finite-scale formulation is unconditional. The open horizon has a name in the program: the *selector*, whatever it is that chooses the zeros, which remains unidentified by design rather than by oversight.

7. WHAT IS ESTABLISHED, AND WHAT IS OPEN

Established (proved in the series): the conic-section classification and divisor-summatory geometry [1]; the $V_4 \leftrightarrow QR(12)$ double bijection [2]; the Lorentz-orbit classification with eigen-coordinates and power maps [3]; the $\mathfrak{su}(2)/\mathfrak{su}(1, 1)$ realizations with the parity selection rule and revival identity [4]; the Casimir completion and null-diamond taxonomy [8].

Exploratory (numerical, not claimed as theorems): the sieve-flow zero tracking [9]; the $T/D/A$ coloring of the cone base; the mod-0.5 divisor detector.

Open: the selector; the positivity certificate; every RH-hard thread, so marked.

8. READING MAP

Order	Paper	What it puts on the cone
1	[1]	Conic sections; divisor summatory
2	[2]	The unit shell; $V_4 \leftrightarrow QR(12)$
3	[3]	Lorentz orbits; power maps
4	[4]	$\mathfrak{su}(2)/\mathfrak{su}(1, 1)$; Fock geometry
5	[5]	Cubic analog; cusp catastrophe; hysteresis
6	[6]	Bicone $SO(4, 2)$; hydrogen dipole; α from one line
7	[7]	Kepler orbits as cone slices; dynamic slicing operator
8	[8]	Casimir completion; null diamond
9	[9]	Sieve flow; zero tracking (numerical)

The discriminant-12 return paper [10] is the program's principal technical paper; the ledger continues the main-line operator search.

ACKNOWLEDGMENT

Large language model tools were used for drafting and structural editing.

REFERENCES

- [1] J. Ebert, *The Cutting Plane: Every Line Through the Multiplication Table Is a Conic Section*, preprint (2026).
- [2] J. Ebert, *Totient Hyperbolas and the $V_4 \leftrightarrow QR(12)$ Bijection: A companion note to Paper A*, preprint (2026).
- [3] J. Ebert, *Signed Cone Cuts as Lorentz Orbits: Eigen-Coordinates and Power Maps*, preprint (2026).
- [4] J. Ebert, *Two Oscillators, Two Algebras: $SU(2)$ and $SU(1,1)$ Realizations of the Cone's Symmetry*, preprint (2026).
- [5] J. Ebert, *The Cusp Instead of the Cone: A Triple-Angle Analog of the AM–GM Circle via the Depressed Cubic, and Its Catastrophe Dynamics*, preprint (2026).
- [6] J. Ebert, *The Fine Structure Constant from Bicone Geometry and One Spectral Line: Hydrogen dipole matrix elements via $SO(4,2)$* , preprint (2026).
- [7] J. Ebert, *The Dynamic Slicing Operator on the AM–GM Cone: A Quadric Geometric Framework for Keplerian Orbits and Perturbed Trajectories*, preprint (2026).
- [8] J. Ebert, *Casimir Completion and the Primitive Null Diamond*, preprint (2026).
- [9] J. Ebert, *The Sieve-Flow Instrument: Zeta Zeros from the Eratosthenes Renormalization*, preprint (2026).
- [10] J. Ebert, *The Discriminant-12 Return: From a Centered Cayley Selection to a Cyclotomic and Finite Arithmetic Structure*, preprint (2026).

INDEPENDENT RESEARCHER