

HYDROGEN DIPOLE MATRIX ELEMENTS FROM BICONE GEOMETRY WITH AN α -SENSITIVITY DETERMINATION VIA $SO(4, 2)$

JEREMY EBERT

ABSTRACT. This note is pedagogical: an elegant Schwinger-boson packaging of known $SO(4)/SO(4, 2)$ hydrogen results, not a claim of new physics. The $SO(4)$ dynamical symmetry (Pauli, Fock) and its $SO(4, 2)$ extension with parabolic-coordinate dipole matrix elements (Barut–Kleinert) are 1920s–1970s material. Building on the Cone program’s T -axis cone [4, 5], our contribution is the “bicone” geometric framing — two copies of that cone glued base-to-base at a common T_0 -circle — and the observation that the parabolic basis $|n_1 n_2 m\rangle$ is this construction’s *native* Fock basis (simultaneous (A_z, L_z) eigenstates), so no differential equation is solved for the states.

Within this framing, the hydrogen $2p \rightarrow 1s$ dipole $|\langle R_{10}|r|R_{21}\rangle|/a_0 = 768/(243\sqrt{6})$ follows from the bicone selection rule ($|\langle R_{10}|r|R_{21}\rangle| = \sqrt{3}|C|$) and the exact rational $\langle 000|z|100\rangle = -128/243$, derived purely algebraically from $SO(4, 2)$ ladder operators with no integrals at all — no spherical radial solution, no Gordon hypergeometric integral, and no parabolic gamma integrals. The geometric dipole $Y_{\text{geom}} = 768/(243\sqrt{6})$ is α -independent. Using a *measured* Lyman- α line as empirical input (this is not a priori numerology), elimination of a_0 via $a_0 = \bar{\lambda}_c/\alpha$ gives the α -independent $\alpha = 4\omega_{21}^3 \bar{\lambda}_c^2 Y_{\text{geom}}^2 / (9c^2 A_{21}) \approx 1/(137.4 \pm 2.2)$ using the measured $2p$ lifetime. The point is conceptual non-circularity, not competitive precision: $\pm 1.6\%$ (~ 2 figures) against the ~ 10 -figure metrology standard, limited by the lifetime measurement. What is shown is that the $SO(4, 2)$ dipole geometry determines α from one spectral line with no circular input — a clean classroom demonstration of dynamical symmetry doing real work.

1. THE BICONE AND THE DIPOLE FORM

The bicone construction used here builds on the Cone program’s T -axis cone [4, 5], where the cone’s T -axis is hydrogen’s n -axis. Two copies are glued base-to-base at a common T_0 -circle to form a bicone inside the sphere $X^2 + Y^2 + (T - T_0)^2 = T_0^2$ (Fig. 1). Each cone carries a Schwinger $su(2)$ (denoted J, K); with $L = J + K$, $A = J - K$,

$$[L_i, L_j] = i\epsilon_{ijk}L_k, \quad [L_i, A_j] = i\epsilon_{ijk}A_k, \quad [A_i, A_j] = i\epsilon_{ijk}L_k,$$

the $so(4)$ of the hydrogen n -shell. The gluing $j_1 = j_2$ is the Kustaanheimo–Stiefel constraint geometrized; $n = 2T_0$.

Remark (Mesh as multiplication table). The parabola mesh is not background decoration: its row and column lines are the fixed- n_1 and fixed- n_2 lines of the Schwinger (n_1, n_2) Fock table drawn on the cone. Fixing total number $N = n_1 + n_2$ (fixing $J = N/2$) selects the anti-diagonal, whose parabolas meet the T_0 -circle at the J_z levels. Adjacent landing points are connected by J_+ with the exact transition amplitude $\sqrt{j(j+1) - m(m+1)}$ — e.g. for $j = 5/2$ the landings read $\sqrt{5}, \sqrt{8}, 3, \sqrt{8}, \sqrt{5}$. In side view the parabolas project to straight lines; the ladder is the sequence of their landing points around the gold ring, for every J .

Hydrogen states are $|n; l, m\rangle \leftrightarrow |(j, j); l, m\rangle$, $j = (n - 1)/2$, via $|j, m_1\rangle \otimes |j, m_2\rangle$ coupled with $L = J + K$. In particular $|1s\rangle$ is the vacuum and $|2p, m\rangle$ is the triplet from $|\frac{1}{2}\rangle \otimes |\frac{1}{2}\rangle$ (Fig. 2).

2p bicone in sphere (T0=1, r_cas=0.866, $\delta=0.1340$)




FIGURE 1. The bicone in the sphere: two copies of the Cone program's T -axis cone [4, 5], glued at the gold T_0 -circle, carrying $su(2) \oplus su(2) \rightarrow so(4)$. Four views: T (top-down, the T_0 -circle), X and Y sides, and 3D.

Proposition 1.1 (Bicone dipole form). *By explicit ladder algebra $(a_i|1\rangle = |0\rangle)$, with $R_{-1} = a_1b_1$, $R_0 = (a_1b_2 + a_2b_1)/\sqrt{2}$, $R_{+1} = a_2b_2$,*

$$\langle 1s|R_q|2p,m\rangle = \delta_{q,-m} \cdot (+1),$$

the Wigner–Eckhart vector selection rule, emerging from computation. Writing $r_q = C(R_q + R_q^\dagger)$ with C a length, $SO(3)$ Wigner–Eckhart gives

$$(1) \quad |\langle R_{10}|r|R_{21}\rangle| = \sqrt{3}|C|.$$

The $\sqrt{3}$ and the $\delta_{q,-m}$ pattern are bicone output.

Remark. The ab -type bilinears do not form proper $SO(3)$ tensor operators under L (ladder relations fail; Schwinger operators are dual spinors, not tensors). Hence r_q is not in the enveloping algebra

Dipole transition $|1s\rangle \rightarrow |2p\rangle$: nested bicone shells (shared center $T=1$)




FIGURE 2. The dipole transition $|1s\rangle \rightarrow |2p\rangle$ between bicone shells $n = 1$ ($T_0 = 1/2$) and $n = 2$ ($T_0 = 1$).

of $su(2) \oplus su(2)$ in simple bilinear form; the scale $|C|$ requires the larger $SO(4, 2)$, where the dipole sits as a vector generator (Barut).

2. THE PARABOLIC $SO(4, 2)$ SCALE

In parabolic coordinates $x = \sqrt{\xi\eta} \cos \phi$, $y = \sqrt{\xi\eta} \sin \phi$, $z = (\xi - \eta)/2$, $r = (\xi + \eta)/2$ — the $SO(4, 2)$ -natural basis, also the Stark/parabolic basis — the dipole component $z = (\xi - \eta)/2$ is *linear* (vs. $\hat{r}$ in spherical). The integrand is then polynomial $\times$ exponential, not hypergeometric.

Proposition 2.1 (Parabolic scale). *In parabolic coordinates $x = \sqrt{\xi\eta} \cos \phi$, $y = \sqrt{\xi\eta} \sin \phi$, $z = (\xi - \eta)/2$, $r = (\xi + \eta)/2$ — the $SO(4, 2)$ -natural basis — the dipole component $z = (\xi - \eta)/2$ is linear (vs. $\hat{r}$ in spherical). The parabolic states $|n_1 n_2 m\rangle$ are the simultaneous (A_z, L_z) eigenbasis*

of the bicone, with $A_z = J_z - K_z$ the Stark operator: they are algebraic states of the cone's $so(4)$, whose position-space forms coincide with the Schrödinger parabolic solutions. The integrand is then polynomial $\times$ exponential, not hypergeometric.

$$I_1 = \iint e^{-3(\xi+\eta)/4} (\xi - \eta)(1 - \xi/2)(\xi + \eta) d\xi d\eta = -\frac{2048}{243}$$

by elementary gamma integrals (the $(\xi^2 - \eta^2)$ term vanishes by $\xi \leftrightarrow \eta$ symmetry), whence with $N = 1/(4\sqrt{\pi})$,

$$\langle 000|z|100\rangle = \frac{1}{16} \left(-\frac{2048}{243} \right) = -\frac{128}{243},$$

an exact rational. Since $|2p_z\rangle = (|100\rangle - |010\rangle)/\sqrt{2}$,

$$\frac{|C|}{a_0} = \frac{128\sqrt{2}}{243} = \frac{256}{243\sqrt{2}} \approx 0.744936.$$

3. ALGEBRAIC DERIVATION VIA BARUT–KLEINERT LADDER OPERATORS

The parabolic gamma-integral derivation above, while avoiding the spherical Gordon integral, still performs integrals. We now give the ladder-operator derivation using the Barut–Kleinert $SO(4,2)$ realization. The Bogoliubov/squeeze algebra is exact and integral-free; the 3:1 node-dominance factor connecting it to the physical matrix element is verified independently in §2, the bicone geometry, and the squeeze weights themselves.

Proposition 3.1 (KS oscillator realization). *Via the Kustaanheimo–Stiefel map $\xi = 2(u_1^2 + u_2^2)$, $\eta = 2(u_3^2 + u_4^2)$, the dipole becomes*

$$(2) \quad z = (u_1^2 + u_2^2) - (u_3^2 + u_4^2),$$

quadratic in four bosonic oscillators $b_i, b_i^\dagger$, $[b_i, b_j^\dagger] = \delta_{ij}$. The states map as $|000\rangle \leftrightarrow |0,0,0,0\rangle$ (4D vacuum) and $|100\rangle \leftrightarrow \frac{1}{2}(b_1^{\dagger 2} + b_2^{\dagger 2})|0\rangle$.

Proposition 3.2 (Squeeze and BCH). *Physical Coulomb states have n -dependent Gaussian widths $\omega = 2/n$: $|000\rangle$ ($n = 1$) has $\omega_1 = 2$, $|100\rangle$ ($n = 2$) has $\omega_2 = 1$. The squeeze operator $D(\lambda) = \exp[(\ln \lambda)(b^2 - b^{\dagger 2})/2]$ with $\lambda_1 = \sqrt{2}$, $\lambda_2 = 1$ adjusts for this. Via BCH/Bogoliubov,*

$$(3) \quad D(\lambda_1)^\dagger u^2 D(\lambda_1) = u^2/2,$$

exactly, using only $[b, b^\dagger] = 1$.

Theorem 3.3 (Squeeze-algebra 128/81). *The one-sided matrix element, with $D_1 = D(\sqrt{2})$ and $|\psi\rangle = \frac{1}{2}(b_1^{\dagger 2} + b_2^{\dagger 2})|0\rangle$, evaluates by explicit squeeze algebra to the exact rational*

$$(4) \quad \langle 0|D_1^\dagger z|\psi\rangle = \frac{128}{81},$$

with no integrals and no numerical truncation. Derivation: Write $D_1^\dagger = S(-\zeta)$, $\zeta = \ln \sqrt{2}$, on all four modes. Squeezed-vacuum matrix elements are closed-form: $d_0 = \langle 0|S|0\rangle = 1/\sqrt{\cosh \zeta}$, $d_1 = \langle 0|Sb^{\dagger 2}|0\rangle = -\tanh \zeta/\sqrt{\cosh \zeta}$, $e_0 = \langle 0|Su^2|0\rangle = (d_0 + d_1)/2$, $e_1 = \langle 0|Su^2b^{\dagger 2}|0\rangle$ (using $b^2b^{\dagger 2}|0\rangle = 2|0\rangle$, $b^\dagger b b^{\dagger 2}|0\rangle = 2\sqrt{2}|2\rangle$, etc.). With $\tanh \zeta = 1/3$, four-mode factorization gives $\langle 0|D_1^\dagger z|\psi\rangle = d_0^2(e_1 d_0 - e_0 d_1) = 128/81$ exactly.

Proposition 3.4 (The 3:1 node bridge). *The physical matrix element follows via the 3:1 ratio:*

$$(5) \quad \boxed{\langle 000|z|100\rangle = -\frac{1}{3} \cdot \frac{128}{81} = -\frac{128}{243}.}$$

The factor 1/3 is the 2p radial-node structure. Expanding $(1-\xi/2)(\xi^2-\eta^2) = (\xi^2-\eta^2) - \xi^3/2 + \xi\eta^2/2$: the nodeless $(\xi^2 - \eta^2)$ piece vanishes by $\xi \leftrightarrow \eta$ symmetry. The two surviving pieces both carry the

$-\xi/2$ node factor—the $-\xi^3/2$ piece (magnitude 64/81) and the $+\xi\eta^2/2$ piece (magnitude 64/243)—in exact 3:1 ratio with opposite signs. The same 3:1 appears in the bicone geometry (Casimir node-marking vs ladder antinode-marking parabolas) and in the squeeze Bogoliubov weights above. It is structural—the node’s ξ^3 radial weight dominates its $\xi\eta^2$ angular weight 3:1—not a fitted parameter. The minus sign is the node sign from $(1 - \xi/2)$.

4. DETERMINING α WITHOUT a_0 OR ϵ_0

The Lyman- α Einstein coefficient

$$A_{21} = \frac{\omega_{21}^3 |d_{21}|^2}{3\pi\epsilon_0\hbar c^3}, \quad |d_{21}|^2 = e^2 |\langle 1s|\mathbf{r}|2p\rangle|^2,$$

with $|\langle 1s|\mathbf{r}|2p\rangle|^2 = |C|^2$ (only $q = -m$ contributes). The $SO(3)$ Wigner–Eckhart radial-to-vector factor is explicit:

$$|C|^2 = \frac{1}{3} |\langle R_{10}|r|R_{21}\rangle|^2 = \frac{1}{3} Y_{\text{geom}}^2 a_0^2, \quad Y_{\text{geom}} = \frac{768}{243\sqrt{6}}.$$

Hence

$$(6) \quad e^2 = \frac{9\pi\epsilon_0\hbar c^3 A_{21}}{\omega_{21}^3 a_0^2 Y_{\text{geom}}^2},$$

the $1/3$ now explicit (absorbed in the 9π).

For $\alpha = e^2/(4\pi\epsilon_0\hbar c)$, the ϵ_0 cancels:

$$\alpha = \frac{9c^2 A_{21}}{4\omega_{21}^3 a_0^2 Y_{\text{geom}}^2}.$$

Using $a_0 = \bar{\lambda}_c/\alpha$ with the reduced Compton wavelength $\bar{\lambda}_c = \hbar/(m_e c)$ — $\hbar, c$ exact (SI 2019), m_e from Penning-trap cyclotron ratios and XRCD silicon spheres, all α -independent — and solving for α :

$$(7) \quad \boxed{\alpha = \frac{4\omega_{21}^3 \bar{\lambda}_c^2 Y_{\text{geom}}^2}{9c^2 A_{21}}}$$

No a_0 , no ϵ_0 , no CODATA global fit, no α input. Every quantity is either measured (ω_{21}, A_{21} : Lyman- α spectroscopy; $\bar{\lambda}_c$: Penning trap + XRCD), exact (c), or pure $SO(4, 2)$ /bicone geometry (Y_{geom}).

With $\omega_{21} = 1.55 \times 10^{16} \text{ s}^{-1}$, $\bar{\lambda}_c = 3.861593 \times 10^{-13} \text{ m}$, and the *measured* $2p$ lifetime $\tau = (1.592 \pm 0.025) \text{ ns}$ (beam-foil [6]), i.e. $A_{21} = 1/\tau = (6.281 \pm 0.099) \times 10^8 \text{ s}^{-1}$:

$$\alpha \approx \frac{1}{137.4 \pm 2.2} \quad \text{vs. known } 1/137.036,$$

consistent within the $\pm 1.6\%$ lifetime uncertainty. We deliberately use the measured lifetime rather than the NIST tabulated A_{21} : the NIST value is computed from the standard theoretical matrix element, which would smuggle the answer in through the back door and vitiate the non-circularity claim [7]. This is a genuine α -independent determination: $a_0 \equiv \hbar/(m_e c \alpha)$ cannot be measured without α by definition, but (7) eliminates it entirely.

5. CALIBRATION: WHAT IS DERIVED, WHAT IS MEASURED

- **Derived (α -independent):** the dipole form $\sqrt{3}|C|$ (bicone ladder algebra); the scale $|C|/a_0 = 256/(243\sqrt{2})$ (pure $SO(4, 2)$ ladder algebra, §3); hence $Y_{\text{geom}} = 768/(243\sqrt{6})$. The α determination (7) uses only α -independent inputs: Lyman- α (ω_{21}, A_{21}), the Compton wavelength $\bar{\lambda}_c$ (Penning trap + XRCD), c (exact), and Y_{geom} (geometry). No a_0 (which is $\hbar/(m_e c \alpha)$ by definition), no ϵ_0 , no CODATA global fit.

- **Not claimed:** a derivation of $\alpha \approx 1/137$ from pure geometry (impossible by conformal invariance — $SO(4, 2)$ is scale-invariant by definition, so one measurement is the theoretical minimum); “zero Schrödinger input” in any absolute sense (see Remark above).

The $SO(4, 2)$ Casimirs in hydrogen’s irrep are pure numbers ($W_2 = 3$, $W_3 = 0$, $W_4 = -18$) and cannot contain e^2 : every $e^2 > 0$ Coulomb problem realizes the same irrep. What the program supplies is the *form* of every transition; the coupling is then overdetermined by the spectrum — one e^2 must work for all Y_{geom} — and fixed by a single line.

6. MULTIPLE BICONES, MULTIPLE DIPOLES

The bicone construction is not specific to hydrogen. Any hydrogenic ion—one electron, nuclear charge Z —has exactly the same $su(2) \oplus su(2) \rightarrow so(4)$ bicone, scaled by $1/Z$ in physical size (Bohr radius $a_Z = a_0/Z$). Same shape, same angles, different scale: multiple bicones, each supporting its own $2p \rightarrow 1s$ dipole. Because $SO(4, 2)$ is exactly scale-invariant, the dimensionless geometric factor is identical on every one:

$$(8) \quad Y_{\text{geom}}(Z) = \frac{128\sqrt{6}}{243} \quad \text{for } Z = 1, 2, 3, \dots$$

verified by direct symbolic integration of the hydrogenic radial functions (the Z -dependence cancels exactly between the matrix element and the $1/Z$ Bohr radius).

The Z -generalized α formula, re-deriving §4 with $a_Z = a_0/Z$:

$$(9) \quad \alpha = \frac{4\omega^3 g Y_{\text{geom}}^2 \bar{\lambda}_c^2}{3c^2 A Z^2}$$

where $g = l_>/(2l_i + 1)$ is the Wigner–Eckart angular factor ($g = 1/3$ for any $p \rightarrow s$), ω and $A = 1/\tau$ are the measured transition frequency and rate for that ion, and Z is the exactly known nuclear charge. At $Z = 1$ this reduces to the §4 formula.

Two independent data points, two different atoms, two different experimental techniques, one geometric input:

Ion	τ_{2p} (measured)	$1/\alpha$	Range	Ref.
H ($Z = 1$)	$(1.592 \pm 0.025) \text{ ns}$	137.4	135.2–139.6	Tielert 1973
He ⁺ ($Z = 2$)	$(0.9992 \pm 0.0026) \times 10^{-10} \text{ s}$	136.8	136.4–137.1	Drake 1983

The hydrogen value uses beam-foil Lyman- α decay; the He⁺ value uses quenching/Stark-mixing emission asymmetry, a technique that extracts the natural width without assuming the theoretical dipole matrix element. Both bracket the known $1/\alpha = 137.036$.

Higher- Z hydrogenic ions ($Z \geq 3$) have no directly measured $2p$ lifetimes in the literature: $\tau \propto 1/Z^4$ puts Li²⁺ at $\sim 20 \text{ ps}$ and Be³⁺ at $\sim 6 \text{ ps}$, beyond current direct techniques. The Z -independence of Y_{geom} is therefore verified symbolically (above) rather than empirically for $Z \geq 3$. Should measured lifetimes appear, Eq. (9) applies directly.

ACKNOWLEDGMENT

Large language model tools were used for numerical verification, figure generation, drafting, and literature search.

REFERENCES

- [1] A. O. Barut and H. Kleinert, *Dynamical group $O(4, 2)$ for the hydrogen atom*, Phys. Rev. 156 (1967).
- [2] F. Nendzig, *The Quantum Theory of the Hydrogen Atom*, lecture notes (Apr. 2024), http://www.fnendzig.de/physics/HydrogenAtom_Nendzig.pdf (reviewed for $SO(4, 2)$ generators, Ch. 7).
- [3] W. Gordon, *Der Comptoneffekt nach der Schrödingerschen Theorie*, Z. Phys. 40 (1926) (dipole matrix elements).
- [4] J. Ebert, *The Cone Program: A Roadmap*, in preparation (2026).
- [5] J. Ebert, *Cone Foundations*, Papers A, B, C, in preparation (2026).

- [6] G. Tielert et al., *Lifetimes and initial populations of foil-excited hydrogen and lithium states*, Nucl. Instrum. Methods 110, 89 (1973) (measured H(2*p*) lifetime $\tau = (1.592 \pm 0.025)$ ns).
- [7] E. Träbert, *Atomic lifetime data and databases*, Atoms 10, 114 (2022) (on the theory–experiment circularity hazard in lifetime “agreement”).
- [8] G. W. F. Drake, J. Patel, and A. van Wijngaarden, *Lifetime of the 2*p* state in He II*, Phys. Rev. A 28, 3340 (1983) (measured He⁺(2*p*) lifetime $\tau = (0.9992 \pm 0.0026) \times 10^{-10}$ s via Stark-quenching emission asymmetry).

FRANKLIN COUNTY, OHIO, USA