

# A Appendix

In Appendix A.1, we provide the details for assessing the persistence of a population with an integrodifference model and we discuss the effect of the harvesting function on population persistence. In Appendix A.2, we provide the details for assessing population persistence with separable dispersal kernels. In Appendix A.3 and A.4, we derive expressions for the critical harvesting rate and rate of environmental shift for Gaussian and sinusoidal dispersal kernels. In Appendix A.5, we derive approximate expressions for these critical rates. In Appendix A.6 we provide details on differences between small and large MPA simulations. In Appendix A.7 we show how reallocation of effort does not qualitatively change our results in our MPA simulations. In Appendix A.8 we parameterize our model for black rockfish in the California Current and demonstrate the qualitative similarities of the results.

**A.1 Determining stability** Let  $n_t(x)$  be the number of adults at position  $x$  at time  $t$ , let  $k(x)$  be a dispersal kernel describing the probability of a larva traveling a distance  $x$ , let  $f(n)$  be the recruitment function describing the number of offspring that settle and survive in juvenile population of size  $n$ , let  $R_0$  be the intrinsic growth rate of the population, and let  $g(n)$  be the harvesting function describing the number of adults harvested from a population of size  $n$ . In the absence of harvesting, the integrodifference model describing the population over time is given by

$$n_{t+1}(x) = \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 f(n_t(y)) dy \quad (1)$$

as described in Zhou and Kot [2011]. With the addition of harvesting, the model becomes

$$n_{t+1}(x) = \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 g(f(n_t(y))) dy. \quad (2)$$

In evaluating persistence, we apply the methods of Zhou and Kot [2011] to the new model, Equation 2. A traveling pulse is a solution such that population size relative to location within the patch (rather than absolute position) is constant over time, i.e.

$$n^*(\bar{x}_t) \equiv n^*(x - ct) = n_t(x),$$

where  $\bar{x}_t \equiv x - ct$  gives position relative to the patch.

The integrodifference equation (2) gives us an expression for  $n^*$ :

$$\begin{aligned}
n^*(\bar{x}_{t+1}) &= n_{t+1}(x) \\
&= \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 g(f(n_t(y))) dy \\
&= \int_{-L/2+ct}^{L/2+ct} k(x-\bar{y}_t-ct) R_0 g(f(n^*(\bar{y}_t))) dy \\
&= \int_{-L/2+ct}^{L/2+ct} k(\bar{x}_t-\bar{y}_t) R_0 g(f(n^*(\bar{y}_t))) dy \\
\Rightarrow n^*(\bar{x}_t-c) &= \int_{-L/2+ct}^{L/2+ct} k(\bar{x}_t-\bar{y}_t) R_0 g(f(n^*(\bar{y}_t))) dy \\
\Rightarrow n^*(\bar{x}_t) &= \int_{-L/2}^{L/2} k(\bar{x}_t+c-\bar{y}_t) R_0 g(f(n^*(\bar{y}_t))) d\bar{y}_t \tag{3}
\end{aligned}$$

As long as  $f(0) = 0$ , there is a trivial solution to this problem where  $n^*(\bar{x}) \equiv 0$  for all  $\bar{x}$ , i.e., there is a trivial traveling pulse with no adults in it. If the trivial traveling pulse is unstable, even very small populations will persist or grow and avoid crashing back to the trivial pulse. To evaluate the stability of a traveling pulse, we introduce a small perturbation to the traveling pulse  $n^*(\bar{x})$  and see if this perturbation grows or shrinks over time:

$$\begin{aligned}
n_t(x) &= n^*(\bar{x}_t) + \xi_t(x) \\
\Rightarrow \xi_{t+1}(x) &= n_{t+1}(x) - n^*(\bar{x}_{t+1}) \\
&= n_{t+1}(x) - n^*(\bar{x}_t-c) \\
&= \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 g(f(n_t(y))) dy - \int_{-L/2}^{L/2} k(\bar{x}_t-\bar{y}_t) R_0 g(f(n^*(\bar{y}_t))) d\bar{y}_t \text{ using (3)} \\
&= \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 g(f(n_t(y))) dy - \int_{-L/2+ct}^{L/2+ct} k(x-ct-(y-ct)) R_0 g(f(n^*(\bar{y}_t))) d\bar{y}_t \\
&= \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 g(f(n_t(y))) dy - \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 g(f(n^*(\bar{y}_t))) d\bar{y}_t \\
&= \int_{-L/2+ct}^{L/2+ct} k(x-y) (R_0 g(f(n_t(y))) - R_0 g(f(n^*(\bar{y}_t)))) dy \\
&= \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 (g(f(n_t(y))) - g(f(n^*(\bar{y}_t)))) dy \\
\Rightarrow \xi_{t+1}(x) &= \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 g'(f(n^*(\bar{y}_t))) f'(n^*(\bar{y}_t)) (n_t(y) - n^*(\bar{y}_t)) dy \\
&\text{by linearizing around the traveling pulse} \\
\Rightarrow \xi_{t+1}(x) &= \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 g'(f(n^*(\bar{y}_t))) f'(n^*(\bar{y}_t)) \xi_t(y) dy \\
\Rightarrow \xi_{t+1}(x) &= \int_{-L/2+ct}^{L/2+ct} k(x-y) R_0 g'(0) f'(0) \xi_t(y) dy \text{ if } n^*(\bar{x}) = 0 \text{ and } f(0) = 0 \tag{4}
\end{aligned}$$

If we assume  $\xi_t(x) = \lambda^t u(x-ct)$  for some  $\lambda \in \mathbb{R}$  and  $u : [-L/2, L/2] \rightarrow \mathbb{R}$ , then the

perturbation grows in time if and only if  $\lambda > 1$ . Using Equation (4), we can rewrite  $\xi_{t+1}(x)$ ,

$$\begin{aligned}\lambda u(x - ct - c) &= R_0 g'(0) f'(0) \int_{-L/2+ct}^{L/2+ct} k(x - y) u(y - ct) dy \\ \Rightarrow \lambda u(\bar{x}) &= R_0 g'(0) f'(0) \int_{-L/2}^{L/2} k(\bar{x} + c - \bar{y}) u(\bar{y}) d\bar{y}\end{aligned}$$

Define the integral operator

$$\psi_f(u)(x) = R_0 g'(0) f'(0) \int_{-L/2}^{L/2} k(x + c - y) u(y) dy.$$

Then the perturbation to the traveling pulse will satisfy

$$\psi_f(u)(x) = \lambda u(x) \quad (5)$$

$\lambda$  and  $u$  are thus an eigenvalue and eigenfunction of the functional operator  $\psi_f$ . The trivial traveling pulse is unstable when the dominant eigenvalue of  $\psi_f$  is greater than 1.

The biomass in the equilibrium traveling wave depends on the specific functional forms of the harvesting function  $g(n)$  and the recruitment function  $f(n)$ . However, the persistence of the population only depends on  $R_0$ ,  $g'(0)$  and  $f'(0)$ . In this paper, we only considered a proportional harvesting function, i.e. the amount of adults harvested obeyed  $g(n) = (1 - h)n$ . For this function,  $g'(0) = 1 - h$ . For the recruitment function we considered,  $f'(0) = 1$ .

**A.2 Separable dispersal kernels** It is not immediately obvious that the operator  $\psi$  will have any eigenfunctions. However, Jentzsch's theorem guarantees that there is an eigenfunction  $u$ , provided that the kernel  $k$  satisfies some properties [Zhou and Kot, 2011]. Finding the eigenfunctions and eigenvalues is in general a hard problem to solve. It becomes easier if the kernel  $k$  is separable, i.e., there are functions  $a_n, b_n$  such that  $k(x - y) = \sum_{n=1}^{\infty} a_n(x) b_n(y)$ . In that case, (5) becomes

$$\begin{aligned}\lambda u(x) &= R_0 g'(0) f'(0) \sum_{n=1}^{\infty} \left( a_n(x) \int_{-L/2}^{L/2} b_n(y - c) u(y) dy \right) \\ \Rightarrow \lambda \int_{-L/2}^{L/2} b_k(x - c) u(x) dx &= R_0 g'(0) f'(0) \sum_{n=1}^{\infty} \left( \int_{-L/2}^{L/2} b_n(x - c) u(x) dx \right) \left( \int_{-L/2}^{L/2} a_n(y) b_k(y - c) dy \right) \\ &\quad \text{for any } k \\ \Rightarrow \lambda d_k &= R_0 g'(0) f'(0) \sum_{n=1}^{\infty} A_{nk} d_n\end{aligned} \quad (6)$$

where

$$A_{nk} = \int_{-L/2}^{L/2} a_n(x) b_k(x - c) dx \text{ and } d_k = \int_{-L/2}^{L/2} b_k(x - c) u(x) dx$$

Finding the eigenvalues of (5) then reduces to finding the eigenvalues of the matrix comprised of entires  $(A_{nk})_{n,k=1}^{\infty}$ .

To find the equilibrium biomass, we rewrite (3) using the separable kernel as in Latore et al. [1998]:

$$\begin{aligned}
n^*(x) &= \int_{-L/2}^{L/2} k(x+c-y) R_0 g(f(n^*(y))) dy \\
&= \int_{-L/2}^{L/2} \left( \sum_{n=1}^{\infty} a_n(x) b_n(y-c) \right) R_0 g(f(n^*(y))) dy \\
&= \sum_{n=1}^{\infty} a_n(x) \int_{-L/2}^{L/2} b_n(y-c) R_0 g(f(n^*(y))) dy
\end{aligned}$$

If we define  $m_n = \int_{-L/2}^{L/2} b_n(y-c) R_0 g(f(n^*(y))) dy$  then we find that

$$\begin{aligned}
n^*(x) &= \sum_{n=1}^{\infty} m_n a_n(x) \text{ and} \\
m_n &= \int_{-L/2}^{L/2} b_n(y-c) R_0 g \left( f \left( \sum_{n=1}^{\infty} m_n a_n(y) \right) \right) dy
\end{aligned} \tag{7}$$

The equations (7) allows us to find the  $m_n$  numerically and we then find the total equilibrium biomass by integrating  $n^*(x)$  over space.

**A.3 Gaussian dispersal kernel** The Gaussian dispersal kernel is given by

$$k(x-y) = \frac{1}{2\sqrt{D\pi}} e^{-\frac{(x-y)^2}{4D}},$$

where  $D$  is one half the variance of the kernel. This is a separable kernel with  $a_n(x) = b_n(x) = \frac{1}{\sqrt{2n! \sqrt{D\pi}}} e^{-x^2/4D} \left( \frac{x}{\sqrt{2D}} \right)^n$  [Latore et al., 1998].

As a first approximation to  $k$  we ignore all but the  $0^{th}$  terms for  $a_n$  and  $b_n$  so that Equation (6) becomes

$$\begin{aligned}
\lambda d_0(c) &= R_0(1-h) A_{00}(c) d_0(c) \\
\Rightarrow \lambda &= R_0(1-h) A_{00}(c) \\
\text{where } A_{00}(c) &= 2\sqrt{2} \exp \left( \frac{-c^2}{8D} \right) \left[ \operatorname{erf} \left( \frac{L-c}{2\sqrt{2D}} \right) - \operatorname{erf} \left( \frac{-L-c}{2\sqrt{2D}} \right) \right]
\end{aligned}$$

where  $\operatorname{erf}$  is the error function. The critical rate of environmental shift  $c^*$  and the critical harvesting rate  $h^*$  are those values of  $c$  and  $h$ , respectively, that make  $\lambda = 1$ .

**A.4 Sinusoidal dispersal kernel** The sinusoidal dispersal kernel is given by

$$k(x-y) = \begin{cases} \frac{w}{2} \cos(w(x-y)) & , \quad |x-y| \leq \frac{\pi}{2w} \\ 0 & , \quad |x-y| > \frac{\pi}{2w} \end{cases}$$

where  $L$  is the length of the patch and we assume  $\frac{\pi}{2w} > L, c < \frac{\pi}{2w} - L$ .

In this case,  $k(x - y) = \frac{w}{2} \cos(wx) \cos(w(y - c)) + \frac{w}{2} \sin(wx) \sin(w(y - c))$  so that  $A_{ij}$  and  $d_i$  can be found for  $i, j = 1, 2$  and (6) reduces to

$$\lambda^2 - \left( \frac{R_0(1-h)wL}{2} \cos(wc) \right) \lambda + \frac{R_0^2(1-h)^2}{16} (w^2L^2 - \sin^2(wL)) = 0.$$

If we solve for  $\lambda$ , we find

$$\lambda = (1-h)R_0 \left[ \frac{wL \cos(wc)}{4} + \frac{1}{4} \sqrt{\sin^2(wL) - w^2L^2 \sin^2(wc)} \right].$$

Zhou and Kot [2011] solve for the critical speed,  $c^*$ , at which the population will be driven extinct:

$$c^* = c^*(R_0) = \frac{1}{w} \cos^{-1} \left[ \frac{16 + R_0^2(1-h)^2(w^2L^2 - \sin^2(wL))}{8R_0(1-h)wL} \right].$$

In our model, we can additionally solve for the critical harvesting rate,  $h^*$ , at which the population will be driven extinct:

$$h^* = 1 - \frac{1}{R_0} \cdot \frac{4wL}{w^2L^2 - \sin^2(wL)} \left[ \cos(wc) - \sqrt{\cos^2(wc) - 1 + \frac{\sin^2(wL)}{w^2L^2}} \right]$$

### A.5 Approximate critical harvesting proportions

We will use the following Taylor series to make approximations of the critical harvesting proportions under the two dispersal kernels:

$$\begin{aligned} \cos(x) &= 1 - \frac{x^2}{2} \\ \cos^2(x) &= 1 - x^2 \\ \sin^2(x) &= x^2 - \frac{x^4}{3} \\ \operatorname{erf}(x) &= \frac{2}{\sqrt{\pi}} \left( x - \frac{x^3}{3} \right) \\ \exp(x) &= 1 + x + \frac{x^2}{2} \end{aligned}$$

For the Gaussian kernel we found

$$h^* = 1 - \frac{2\sqrt{2} \exp\left(\frac{c^2}{8D}\right)}{R_0 \left[ \operatorname{erf}\left(\frac{L-c}{2\sqrt{2D}}\right) - \operatorname{erf}\left(\frac{-L-c}{2\sqrt{2D}}\right) \right]} \quad (8)$$

Using the Taylor series and the fact that  $D = \frac{\sigma^2}{2}$  where  $\sigma^2$  is the variance of the exponential kernel,

$$\begin{aligned} h^* &\sim 1 - \frac{\sqrt{2\pi} \left( 1 + \frac{c^2}{8D} + \frac{c^4}{128D^2} \right)}{R_0 \sqrt{\pi} \left[ \frac{L-c}{2\sqrt{2D}} - \frac{(L-c)^3}{3(2\sqrt{2D})^3} - \frac{-L-c}{2\sqrt{2D}} + \frac{(-L-c)^3}{3(2\sqrt{2D})^3} \right]} \\ &= 1 - \frac{1}{R_0} \cdot \frac{3\sqrt{2\pi}}{8L} \frac{(32\sigma^4 + 8c^2\sigma^2 + c^4)}{\sigma(12\sigma^2 - (L^2 + 3c^2))} \end{aligned}$$

For the sinusoidal kernel we found

$$h^* = 1 - \frac{1}{R_0} \cdot \frac{4wL}{w^2L^2 - \sin^2(wL)} \left[ \cos(wc) - \sqrt{\cos^2(wc) - 1 + \frac{\sin^2(wL)}{w^2L^2}} \right] \quad (9)$$

Using the Taylor series and the fact that  $w = \frac{\sqrt{\frac{\pi^2}{4} - 2}}{\sigma}$  where  $\sigma^2$  is the variance of the sinusoidal kernel,

$$\begin{aligned} h^* &\sim 1 - \frac{1}{R_0} \cdot \frac{12wL}{w^4L^4} \left[ 1 - \frac{w^2c^2}{2} - \sqrt{1 - w^2c^2 - \frac{w^2L^2}{3}} \right] \\ &= 1 - \frac{1}{R_0} \cdot \frac{4\sqrt{3}}{L^3(\pi^2 - 8)^{3/2}} \cdot \sigma \left[ 8\sqrt{3}\sigma^2 - (\pi^2 - 8)\sqrt{3}c^2 - 4\sigma\sqrt{12\sigma^2 - (\pi^2 - 8)(3c^2 + L^2)} \right] \end{aligned}$$

In the case of both kernels, the critical harvesting proportion can be approximated by a function that looks like

$$h^* \sim 1 - \frac{1}{R_0} \cdot p(L)q(\sigma^2, c^2, L^2 + 3c^2) \quad (10)$$

where  $p(L)$  is a decreasing function of the length of the viable patch  $L$ .

### A.6 Protected Area fluctuations

After the simulations come to equilibrium, the fluctuations in total biomass per generation fluctuate more in reserves that are larger and spaced farther apart than simulations in which the reserves that are smaller and more closely spaced. The large reserves have a slightly larger average population, however large reserves here can induce fluctuations of biomass even in deterministic simulations. Thus we expect if reproduction was stochastic, large reserves spaced far apart would be more likely to result in extinction of the population than more closely spaced, smaller reserves. We find the same effect regardless of whether or not effort remains constant.




Figure 1: Total population biomass is on the y axis, and generation is on the x axis. These simulations were run with climate velocity = 0.1, and a proportional harvest rate = 0.02.

**A.7 Effort reallocation** Effort reallocation does not affect our results.



**A.8 California Current Black Rockfish Parameterization** We parameterize our model for black rockfish in the California Current, with MPAs of spacing and width that is qualitatively similar to the MLP Act reserves, and with a maximum climate velocity equal to that observed empirically. The parameters are as follows

We find qualitatively similar results with this parameterization to what is presented in the main text.




Figure 2: The equilibrium biomass of the population as a function of the climate velocity on the x-axis and the harvesting rate on the y-axis under alternative management strategies as parameterized for black rockfish. (a) The equilibrium biomass for simulations with constant harvest rates. (b) Equilibrium biomass for simulations with threshold management. For threshold management, the maximum threshold below which no harvesting is allowed is set to be the largest population size observed at a given time step before harvesting. For a less severe threshold, we use a proportion of this maximum threshold, so that a lower proportion gives a lower threshold and allows for more harvesting. We show this proportion on the y-axis. (c) Equilibrium biomass for simulations with protected areas where harvesting pressure outside reserves unchanged. (d) Equilibrium biomass for simulations with protected areas with harvesting pressure reallocated outside reserves. These results are from a simulation with a Laplacian dispersal kernel with parameters  $L = 1$ ,  $R_0 = 2.83$ ,  $K = 1$ , and  $b = 73$ .

## References

- J. Latore, P. Gould, and A. M. Mortimer. Spatial dynamics and critical patch size of annual plant populations. *Journal of Theoretical Biology*, 190(3):277–285, 1998.
- Ying Zhou and Mark Kot. Discrete-time growth-dispersal models with shifting species ranges. *Theoretical Ecology*, 4(1):13–25, 2 2011. ISSN 1874-1738. doi: 10.1007/s12080-010-0071-3.