bfield

Returns $\dot{s} \equiv B^s/B^{\zeta}$ and $\dot{\theta} \equiv B^{\theta}/B^{\zeta}$.

[called by: pp00ab.]

contents

1	bfie	ld
	1.1	equations of field line flow
	1.2	representation of magnetic field

1.1 equations of field line flow

1. The equations for the fieldlines are normalized to the toroidal field, i.e.

$$\dot{s} \equiv \frac{B^s}{B^{\zeta}}, \qquad \dot{\theta} \equiv \frac{B^{\theta}}{B^{\zeta}}. \tag{1}$$

1.2 representation of magnetic field

1. The components of the vector potential, $\mathbf{A} = A_{\theta} \nabla + A_{\zeta} \nabla \zeta$, are

$$A_{\theta}(s,\theta,\zeta) = \sum_{i,l} A_{\theta,e,i,l} \, \overline{T}_{l,i}(s) \cos \alpha_i + \sum_{i,l} A_{\theta,o,i,l} \, \overline{T}_{l,i}(s) \sin \alpha_i, \tag{2}$$

$$A_{\zeta}(s,\theta,\zeta) = \sum_{i,l} A_{\zeta,e,i,l} \, \overline{T}_{l,i}(s) \cos \alpha_i + \sum_{i,l} A_{\zeta,o,i,l} \, \overline{T}_{l,i}(s) \sin \alpha_i, \tag{3}$$

where $\overline{T}_{l,i}(s) \equiv \overline{s}^{m_i/2} T_l(s)$, $T_l(s)$ is the Chebyshev polynomial, and $\alpha_j \equiv m_j \theta - n_j \zeta$. The regularity factor, $\overline{s}^{m_i/2}$, where $\overline{s} \equiv (1+s)/2$, is only included if there is a coordinate singularity in the domain (i.e. only in the innermost volume, and only in cylindrical and toroidal geometry.)

2. The magnetic field, $\sqrt{g} \mathbf{B} = \sqrt{g} B^s \mathbf{e}_s + \sqrt{g} B^\theta \mathbf{e}_\theta + \sqrt{g} B^\zeta \mathbf{e}_\zeta$, is

$$\sqrt{g} \mathbf{B} = \mathbf{e}_{s} \sum_{i,l} \left[(-m_{i} A_{\zeta,e,i,l} - n_{i} A_{\theta,e,i,l}) \overline{T}_{l,i} \sin \alpha_{i} + (+m_{i} A_{\zeta,o,i,l} + n_{i} A_{\theta,o,i,l}) \overline{T}_{l,i} \cos \alpha_{i} \right]
+ \mathbf{e}_{\theta} \sum_{i,l} \left[(-m_{i} A_{\zeta,e,i,l}) \overline{T}'_{l,i} \cos \alpha_{i} + (-m_{i} A_{\zeta,o,i,l} + n_{i} A_{\theta,o,i,l}) \overline{T}'_{l,i} \sin \alpha_{i} \right]
+ \mathbf{e}_{\zeta} \sum_{i,l} \left[(-A_{\theta,e,i,l}) \overline{T}'_{l,i} \cos \alpha_{i} + (-A_{\theta,o,i,l}) \overline{T}'_{l,i} \sin \alpha_{i} \right]$$
(4)

3. In Eqn.(1), the coordinate Jacobian, \sqrt{g} , cancels. No coordinate metric information is required to construct the fieldline equations from the magnetic vector potential.

bfield.h last modified on; SPEC subroutines;