

The Use of Antibiotics in the Empirical Treatment of Acute Childhood Gastroenteritis in a Tertiary Hospital

Short Communication

* David, E. E¹, Igwenyi, I. O², Iroha, I. R³, Eghosa, L. E⁴, Ezeilo, U. R⁵, Ogbanshi, M. E¹⁶, David, C. N³

^{1,5,6,7}Department of Biochemistry, Alex Ekwueme Federal University, Ndufu-Alike, Ikwo, Nigeria

²Department of Biochemistry, Ebonyi State University, Ebonyi State, Nigeria

³Department of Microbiology, Ebonyi State University, Ebonyi State, Nigeria

⁴Department of Pediatrics, St Patrick's Hospital, Mile 4 Abakaliki, Ebonyi State, Nigeria

*Correspondence: david.ebuka@funai.edu.ng

Abstract

Back ground: The routine empirical use of antibiotics for infectious diarrhea in children in developing countries is associated with the risk of increasing antibiotics resistance. This work studied the use of antibiotics in the empirical treatment of acute childhood gastroenteritis. **Methodology:** The medical folders of children diagnosed with acute gastroenteritis and hospitalized at the pediatric ward of St Patrick's Hospital, Mile 4, Abakaliki, Ebonyi State, Nigeria were used to collect the antibiotics treatment regime and socio-demographic data of each child. **Results:** The result showed that most of the children were treated without antibiotics, 27% [13] out of the 48 medical folders examined. Ceftriaxone was seen as the most singly used antibiotics at 17% [8], followed by ciprofloxacin at 15% [7]. Male children were recorded to be more affected by gastroenteritis 68.8% [33] than the females 31.3% [15]. **Conclusion:** The widespread use of beta-lactam antibiotics in the empirical treatment of childhood gastroenteritis may lead to the emergence of extended spectrum beta lactamase (ESBL) producing pathogens.

Keywords: Acute ,Antibiotics , Childhood,Empirical-treatment, Gastroenteritis

1. Introduction

Gastroenteritis is an inflammation of the lining of the intestine caused by a virus, bacteria or parasites. It causes watery diarrhea, pain or abdominal cramp, nausea, vomiting and sometimes fever (Hartman *et al.*, 2019). Worldwide, about 3-5 billion cases of acute gastroenteritis occur yearly with nearly 2 million deaths in children below 5 years. This makes diarrhea disease the fifth leading cause of death in children, and the eighth leading cause of death among all ages (Troeger *et al.*, 2018). Nigeria is one of the two countries accounting for 42% of global deaths attributable to gastroenteritis in children less than five years (Arowolo *et al.*, 2019). Due to the high mortality rate of gastroenteritis in children, treatment decisions are mostly made on clinical

outcome, while the choice of antimicrobial drugs is empirically made, based on narrow drugs that cover most of the possible pathogens (Diniz-Santos *et al.*, 2016). The necessity for this empirical treatment method is because, the cost-effective means of identifying these pathogens involve conventional stool culture and antimicrobial susceptibility testing of isolated organisms which usually takes not less than 72 hours. The use of ciprofloxacin (fluoroquinolone) was recommended by WHO, as the first-line drugs in the treatment of bloody diarrhea (WHO, 2005). Empirically, cephalosporins, especially, cefixime and ceftriaxone are considered the best treatment for acute childhood gastroenteritis (Lieberman, 2003). However, reports have confirmed that ciprofloxacin are equally safe to be used in pediatrics, with some restrictions, like in severe cases (Bruzzeze *et al.*, 2018).

Unless in severe and in bloody diarrhea cases, antibiotics are not prescribed in the treatment of acute childhood diarrhea disease (Qureshi *et al.*, 2021). Routine use of antibiotics for infectious diarrhea in children is associated with the risk of increasing antibiotics resistance in developing countries (Diniz-Santos *et al.*, 2016; Ayukekbong *et al.*, 2017; Britto *et al.*, 2018). Without considerations to the WHO guidelines, ciprofloxacin (fluoroquinolone) and cephalosporins (ceftriaxone and cefepime) have been observed to be routinely used by physicians in the treatment of this disease in children in Nigeria (11-13)(Ekwochi *et al.*, 2013; Efunshile *et al.*, 2019; David *et al.*, 2021). Unlike in developed nations, intravenous re-hydrations are commonly used (Burstein *et al.*, 2022; Stanyevic *et al.*, 2022; Weghorst *et al.*, 2022). This indiscriminate use of antibiotics has resulted in the emergence of both multi-drug and cephalosporins (extended spectrum beta-lactamase, ESBL) resistant enteric pathogens (Wallace *et al.*, 2020). Center for Disease Control has categorized ESBL-producing *enterobacteriaceae* as an “urgent threat” (CDC, 2019). This work studied the use of antibiotics in the empirical treatment of acute childhood gastroenteritis in Mile 4 Hospital, Abakaliki, Nigeria.

2. Methodology


The study recruited medical folders of 48 children between the ages of 0-5 years who were diagnosed with gastroenteritis from January to March, 2020. These months covered the window period for the yearly outbreak of gastroenteritis in Abakaliki, Ebonyi State, Nigeria. The stools were collected from the pediatrics unit of the hospital. Ethical clearance (Ref.No: RE/M4H/48/19) was obtained from Ethical and Research Committee of the hospital, after which informed consent was obtained from the parents/guardians/attendants of the children. Socio-demographic characteristics like age, gender, living area, clinical symptoms, co-morbidity were collected for each child. Trends in the use of antibiotics, such as types of antibiotics and their overall effectiveness were also collected for each child.

3. Results

The pattern of empirical treatment of children with acute gastroenteritis in Mile 4 Hospital is shown in figure 1. A greater percentage of the children was treated without antibiotics, 13 [27%]. Ceftriaxone was seen as the most singly used antibiotics at 8 [17%], followed by ciprofloxacin at 7 [15%]. Cefepime and azithromycin represented the least used single antibiotics at 2 [4%] each.

The Use of Antibiotics in the Empirical Treatment of Acute Childhood Gastroenteritis

The most common double combination of antibiotics was the use of ciprofloxacin and ceftriaxone, 9 [19%], followed by ceftriaxone and cefepime at 4 [8%]. Triple combination of the use of ciprofloxacin, ceftriaxone and cefepime was 1 [2%], which was also the same with the double use of ciprofloxacin and cefepime.



#significantly different when compared to the No Antibiotics group@

CIP=Ciprofloxacin; CRO=Ceftriaxone; CPM=Cefepime; AZT=Azithromycin

Figure 1: Antibiotics used in empirical treatment of childhood gastroenteritis (n=48)

Table 1 shows the gender, age and treatment duration of children with acute gastroenteritis. The mean age of the children was 10.5 months [10.5±7.2]. Male children were recorded to be more affected by gastroenteritis 33 [68.8%] than the females 15 [31.3%]. The mean number of days spent at the hospital was 105 hrs [4.9±1.8]. Children treated with antibiotics were observed to spend higher number of days, 5.4±1.7 at the hospital than those not treated with antibiotics 3.5±1.5.

Table 1: Gender, age and treatment duration (n=48)

Male	Female	Mean age (mo)	Hospital duration (hr)		
			Mean	Antibiotics	No antibiotics
33 [68.8%]	15 [31.3%]	10.5±7.2	105±1.8	124±1.7 ^b	77±1.5 ^c

Significantly higher than c (p=0.0005).

4. Discussions

This study revealed that a greater percentage [27%] of the children was treated without antibiotics, while ceftriaxone was seen as the most singly used antibiotics at 17%, followed by ciprofloxacin at 15%. Male children were recorded to be more affected by gastroenteritis [68.8%] than the females [31.3%]. In this health facility, while higher number of children was treated without antibiotics, the guidelines established by WHO on the use of antibiotics were not fully followed. Ciprofloxacin (fluoroquinolone) was only recommended by WHO as the first-line drugs in the treatment of bloody diarrhea (WHO, 2005). The high mortality rate of gastroenteritis in children has led to treatment decisions mostly made on clinical outcome, hence, empirical treatments with antimicrobial drugs are based on narrow drugs that cover most of the possible pathogens (Diniz-Santos *et al.*, 2016). While the hospital rightly use cefixime and ceftriaxone which are considered the best treatment of acute childhood gastroenteritis (Lieberman, 2003), they equally used cefepime, a fourth generation cephalosporin which was not recommended. Routine use of antibiotics for infectious diarrhea in children is associated with the risk of increasing antibiotics resistance in developing countries (Diniz-Santos *et al.*, 2016). In a multi-stage random sampling study involving different health facilities in Rivers State, Nigeria, antibiotics were prescribed in 78.6% of the children with acute gastroenteritis. Unlike our result, metronidazole was prescribed in 50.9% of the cases, while ceftriaxone was used in 0.2% (Ayukekbong *et al.*, 2017). In another tertiary institution in Abakaliki Nigeria and Accra Ghana, 88.9% and 95% of the children were treated with antibiotics respectively (Ekwochi *et al.*, 2013). In an observational study of 210 children with acute gastroenteritis in Enugu, Nigeria, the use of antibiotics was observed to start from the caregivers, even before hospital admission, where 46.7% of the children were given un-prescribed antibiotics (Efunshile *et al.*, 2019).

Conclusion

The most used of ceftriaxone, a beta-lactam antibiotic in the empirical treatment of childhood gastroenteritis in this hospital facility may lead to the emergence of extended spectrum beta lactamase (ESBL) producing pathogens. Beta-lactamases break down beta lactam drugs and confer the ability to resist these drugs to pathogens.

References

- Arowolo, K.O., Ayolabi, C.I., Lapinski, B., Santos, J.S and Raboni, S.M (2019). Epidemiology of enteric viruses in children with gastroenteritis in Ogun State, Nigeria. *Journal of Medical Virology*, 91 (6), 1022–9.
- Ayukekbong, J.A., Ntemgwa, M and Atabe, A.N (2017). The threat of antimicrobial resistance in developing countries: causes and control strategies. *Antimicrobial Resistance & Infection Control*, 6 (1), 1–8.
- Britto, C.D., Wong, V.K., Dougan, G and Pollard, A.J (2018). A systematic review of antimicrobial resistance in *Salmonella enterica* serovar Typhi, the etiological agent of typhoid. *PLoS Neglected Tropical Diseases*, 12 (10), e0006779.

- Bruzzese, E., Giannattasio, A and Guarino, A (2018). Antibiotic treatment of acute gastroenteritis in children. *F1000Research*, 7.
- Burstein, B., Rogers, S., Klassen, T.P and Freedman, S.B (2022). Trends in Management of Children With Acute Gastroenteritis in US Emergency Departments. *JAMA Network Open*, 5(5), e2211201--e2211201.
- Center for Disease Control and Prevention (2019). Antibiotic resistance threats in the United States, 2019. US Department of Health and Human Services.
- David, E.E., Igwenyi, I.O., Iroha, I.R., David, C.N., Mbah, P.C and Okpala, O.F (2021). Trends in Empirical Treatment of Hospitalized Children with Acute Gastroenteritis in Nigeria. *Recent Advances in Anti-Infective Drug Discovery Former Recent Patents Anti-Infective Drug Discov*, 16 (3), 237–44.
- Diniz-Santos, D.R., Silva, L.R and Silva, N (2006). Antibiotics for the empirical treatment of acute infectious diarrhea in children. *Brazilian Journal of Infectious Disease*, 10 (3), 217–27.
- Efunshile, A.M., Ezeanosike, O., Nwangwu, C.C., König, B., Jokelainen, P and Robertson, L.J (2019). Apparent overuse of antibiotics in the management of watery diarrhoea in children in Abakaliki, Nigeria. *BMC Infectious Disease*, 19 (1), 1–7.
- Ekwochi, U., Chinawa, J.M., Obi, I., Obu, H.A and Agwu, S (2013). Use and/or misuse of antibiotics in management of diarrhea among children in Enugu, Southeast Nigeria. *Journal of Tropical Pediatrics*, 59(4), 314–6.
- Hartman, S., Brown, E., Loomis, E and Russell, H.A (2019). Gastroenteritis in children. *Am Fam Physician*, 99 (3), 159–65.
- Lieberman, J.M (2003). Appropriate antibiotic use and why it is important: the challenges of bacterial resistance. *Pediatrics Infectious Disease Journal*, 22 (12), 1143–51.
- Qureshi, S., Resham, S., Hashmi, M., Naveed, A.B., Haq, Z and Ali, S.A (2021). A retrospective review on antibiotic use in acute watery diarrhea in children in a tertiary care hospital of Karachi, Pakistan. *PLoS One*, 16 (7), e0253712.
- Stanyevic, B., Sepich, M., Biondi, S., Baroncelli, G.I., Peroni, D and Di Cicco, M (2022). The evolving epidemiology of acute gastroenteritis in hospitalized children in Italy. *European Journal of Pediatrics*, 181 (1), 349–58.
- Troeger, C., Blacker, B.F., Khalil, I.A., Rao, P.C., Cao, S and Zimsen, S.R.M (2018). Estimates of the global, regional, and national morbidity, mortality, and aetiologies of diarrhoea in 195 countries: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Infect Dis*, 18 (11), 1211–28.

Wallace, M.J., Fishbein, S.R.S and Dantas, G (2020). Antimicrobial resistance in enteric bacteria: current state and next-generation solutions. *Gut Microbes*, 12 (1), 1799654.

Weghorst, A.A.H., Bonvanie, I.J., Holtman, G.A., de Boer, M.R and Berger, M.Y (2022). Course of uncomplicated acute gastroenteritis in children presenting to out-of-hours primary care. *BMC Primary Care*, 23, (1), 1–8.

World Health Organization (2005). Guidelines for the control of shigellosis, including epidemics due to *Shigella dysenteriae* type 1. World Health organisation.
<https://apps.who.int/iris/handle/10665/43252>

Funding

This research was funded by International Center for Genetic engineering and Molecular Biology (ICGEB) under Arturo Falaschi ICGEB Fellowship with funding number S_NGA20-01-1. The funders had no role in study design, data collection, and analysis; in the decision to publish; or in the preparation of the manuscript.

Ethical issues

The present research work was permitted by the Ebonyi State University, University Human Ethical Committee and Ethical and Research Committee of Alex, Ekwueme Federal Teaching Hospital, Abakaliki. Nigeria

Conflict of interests

No conflict of interests are declared by authors.

Author's contributions

Conceptualization and design of the work : David, E.E., Igwenyi, I.O .; data acquisition, David, E.E., Eghosa, L. E., David, C.N³.; analysis and interpretation of data, David E.E., Ezeilo, U. R ., original draft preparation :David E.E.; manuscript review and editing: David E.E, Ogbanshi, M. E; supervision: Igwenyi, I.E., Iroha, I.R; funding acquisition: David, E.E. All authors read and agreed to the version to be published and to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Int. J.Bio. Sc. Mol. Res. April 2023 Vol.1(2) 85-90.