The endpoint slope of the first zero of $E_{a,2}(-t^a)$

Principia Math In Progress

21 September 2026

Task 1: MathDB 361625. Chaudhary, Diethelm and Hashemishahraki [1] conjecture

$$Z_2(a) = \pi + (2-a) + o(2-a) \qquad (a \uparrow 2), \tag{1}$$

where $Z_2(a)$ is the smallest positive zero of

$$G(a,t) = E_{a,2}(-t^a) = \sum_{k\geq 0} \frac{(-1)^k t^{ak}}{\Gamma(ak+2)}.$$

The coefficient in (1) is stated as exactly 1, both in current preprint Conjecture 2(a)(i) and journal-final Conjecture 2(b)(i). Here is its exact correction.

Theorem 1. *For a sufficiently close to 2, $G(a,\cdot)$ has a smallest positive zero, and*

$$\boxed{Z_2(a) = \pi + c(2-a) + O((2-a)^2), \qquad c = \frac{\pi\,\mathrm{Ci}(\pi) + \mathrm{Si}(\pi)}{2}.}$$

Moreover $1041/1000 < c < 1043/1000$, so $c \neq 1$. Consequently (1) is false.

Our conventions are

$$\mathrm{Si}(x) = \int_0^x \frac{\sin u}{u}\,du, \qquad \mathrm{Ci}(x) = \gamma + \log x + \int_0^x \frac{\cos u - 1}{u}\,du \quad (x>0),$$

where γ is Euler's constant. Numerically, $c = 1.041685811636407754238075729 5\ldots$; the rational enclosure below, rather than these digits, proves the separation from 1. The source's conjecture about its threshold near 1.599115206, its collision expansion there, and global monotonicity is not proved or refuted individually by this endpoint result. Their conjunction with (1), as recorded in MathDB, is refuted.

The first zero is a smooth local branch

Put $F(a,t) = tG(a,t)$. At $a = 2$ we have

$$F(2,t) = \sin t, \qquad F(2,\pi) = 0, \qquad F_t(2,\pi) = -1.$$

The defining series and its first two parameter and t derivatives converge locally uniformly near $(2,\pi)$. For example, on a compact rectangle with $3/2 \leq a \leq 5/2$ and t bounded away from zero, $\Gamma(ak+2) \geq (k+1)!$, while derivatives add only polynomial factors in k and $\log(k+2)$. A convergent

factorial majorant applies. Thus the implicit function theorem gives a C^2 branch $t = z(a)$, with $z(2) = \pi$, and

$$z'(2) = -\frac{F_a(2,\pi)}{F_t(2,\pi)} = F_a(2,\pi). \tag{2}$$

This branch is the *first* positive zero. Choose a small fixed $\eta > 0$ so that $\cos t < -1/2$ on $[\pi - \eta, \pi + \eta]$. For a close enough to 2, uniform continuity of F_t gives $F_t(a,t) < -1/4$ on this interval. Hence it contains exactly one zero, namely $z(a)$. On $[0, \pi - \eta]$, the series for G converges uniformly for a near 2, including $t = 0$. Its limit $G(2,t) = \sin t/t$ (value 1 at zero) has a positive minimum there. It follows that $G(a,t) > 0$ throughout that earlier interval. There are therefore no positive zeros before $z(a)$.

An exact expression for the derivative

Let $w(t) = F_a(2,t)$ and $H_m = \sum_{j=1}^{m} 1/j$. Differentiating the series and using $\psi(2k+2) = H_{2k+1} - \gamma$ gives

$$w(t) = \sum_{k\geq 1} \frac{(-1)^k k t^{2k+1}}{(2k+1)!}(\log t + \gamma - H_{2k+1}). \tag{3}$$

Equivalently,

$$w(t) = -\frac{1}{2}\int_0^t (\log(t-u) + \gamma) u \sin u \, du. \tag{4}$$

To verify this identity without an inversion formula, use

$$\int_0^t u^m (\log(t-u) + \gamma)\, du = \frac{t^{m+1}}{m+1}(\log t + \gamma - H_{m+1}) \quad (m \geq 0).$$

After $u = tv$, the identity follows from $\int_0^1 v^m \log(1-v)\, dv = -H_{m+1}/(m+1)$, itself obtained by integrating the logarithm series. Expand $u \sin u$ in (4) and integrate termwise, justified by absolute convergence against the integrable logarithmic singularity. The coefficient identity

$$\frac{1}{2(2k-1)!(2k+1)} = \frac{k}{(2k+1)!}$$

then gives precisely (3).

By (2), the desired coefficient is $c = -w(\pi)$. Changing variables $v = \pi - u$ in (4) yields

$$c = \frac{1}{2}\int_0^\pi (\log v + \gamma)(\pi - v) \sin v \, dv. \tag{5}$$

The following integrations by parts evaluate it:

$$\int_0^\pi \log v \sin v \, dv = \log \pi + \mathrm{Ci}(\pi) - \gamma,$$
$$\int_0^\pi v \log v \sin v \, dv = \pi \log \pi - \mathrm{Si}(\pi),$$
$$\int_0^\pi (\pi - v) \sin v \, dv = \pi.$$

For the first identity the primitive is $-\log v \cos v + \mathrm{Ci}(v)$, with limit γ at zero. For the second it is $-v \log v \cos v + (\log v + 1)\sin v - \mathrm{Si}(v)$, with limit zero at zero. Substitution in (5) cancels both γ and $\log \pi$, giving the stated formula. Taylor's theorem for the C^2 branch gives the $O((2-a)^2)$ remainder.

A rational certificate that $c \neq 1$

Here are fully specified finite rational bounds, reproduced by the accompanying verifier. No floating-point special-function evaluation is used for this certificate. First enclose

$$p_- := \frac{31415926535}{10^{10}} < \pi < p_+ := \frac{31415926536}{10^{10}}.$$

These inequalities follow from Machin's identity $\pi = 16\arctan(1/5) - 4\arctan(1/239)$: in each arctangent use the first 12 terms of its alternating series and its next term as the remainder bound.

For a positive rational $x > 1$, put $y = (x-1)/(x+1)$ and

$$A(x) = 2\sum_{j=0}^{49} \frac{y^{2j+1}}{2j+1}, \qquad B(x) = A(x) + \frac{2y^{101}}{101(1-y^2)}.$$

The positive series for $\log x$ gives $A(x) < \log x < B(x)$. Let $N = 2048 = 2^{11}$. The elementary harmonic-sum bounds give

$$H_N - 11B(2) - \frac{1}{N} < \gamma < H_N - 11A(2).$$

Indeed, $H_N - \log N - \gamma$ is the positive series $\sum_{j\geq N}(\log(1+1/j) - 1/(j+1))$, which is less than $\sum_{j\geq N}(1/j - 1/(j+1)) = 1/N$. Also $A(p_-) < \log\pi < B(p_+)$.

Finally use

$$\mathrm{Ci}(\pi) = \gamma + \log\pi + \sum_{k=1}^{10} \frac{(-1)^k \pi^{2k}}{2k(2k)!} + R_C, \qquad -\frac{p_+^{22}}{22\cdot 22!} < R_C < 0,$$

$$\mathrm{Si}(\pi) = \sum_{k=0}^{10} \frac{(-1)^k \pi^{2k+1}}{(2k+1)(2k+1)!} + R_S, \qquad -\frac{p_+^{23}}{23\cdot 23!} < R_S < 0.$$

The term magnitudes decrease since $\pi^2 < 10$; direct ratios of successive terms verify this for both series. Evaluate each positive power in $[p_-^j, p_+^j]$, add intervals with the displayed signs, and multiply intervals using all four endpoint products. These specified rational operations give an enclosing interval $[c_-, c_+]$ satisfying the exact integer comparisons

$$\frac{1041}{1000} < c_- < c_+ < \frac{1043}{1000}.$$

The verifier stores both exact rational endpoints. For orientation only, they are approximately 1.041302284865325 and 1.042069276032071. Thus $c > 1$ rigorously, finishing the disproof.

Verification, attribution, and follow-ups

The implicit-function argument and parameter differentiation are standard methods. Earlier perturbation work on fractional oscillators, including [2], is relevant background; these methods are not claimed as new. The target result here is the exact first-zero slope and its rigorous separation from the source's coefficient. This endpoint result for second parameter 2 is distinct from the order-one logarithmic asymptotic for second parameter 1 in MathDB 361626. All source and priority checks are recorded separately.

The verifier checks the rational certificate, derivative primitives, and series coefficient identity exactly. Supplemental high-precision computations compare (3), (4), and the closed formula, and check the slope with zeros at orders approaching 2. Those supplemental computations do not establish first-zero minimality. Review is by the authoring agent only.

Task 2: Determine the quadratic coefficient of $(2-a)^2$ in $Z_2(a)$ and give a rigorous enclosure for it.

Task 3: Derive the corresponding endpoint expansion for every fixed positive zero approaching $n\pi$, with its dependence on n explicit.

Task 4: Prove or refute strict decrease of $Z_2(a)$ throughout the whole interval on which the source conjectures its existence.

Task 5: Rigorously locate the first double-zero threshold and prove or refute the proposed square-root collision law there.

References

[1] R. Chaudhary, K. Diethelm and S. Hashemishahraki, On the separation of solutions to fractional differential equations of order $\alpha \in (1,2)$, *Applied Numerical Mathematics* **203** (2024), 84–96, doi:10.1016/j.apnum.2024.05.020. Current preprint: arXiv:2401.14771v3.

[2] V. E. Tarasov and G. M. Zaslavsky, Dynamics with low-level fractionality, *Physica A* **368** (2006), 399–415, doi:10.1016/j.physa.2005.12.015. Inspected preprint: arXiv:physics/0511138v1, Sections 1–3.2 (through preprint p. 10).