Uniqueness in the mirror Markov tree

Principia Math In Progress

22 September 2026

MathDB 373162. Task 1: a proof of Conjecture 6.3 of [1]. Fix a mirror Markov tree with the initial segment and signed mutations of [1, Definitions 4.1–4.2]. Its triples are regarded as vertices of that tree, with the source's ordering by evaluation at $q = 1$.

Theorem 1. *Two distinct triples in this tree cannot have the same polynomial as their maximum-degree entry. In fact, the degree and the value at $q = 1$ of that entry already determine the triple.*

The proof uses the established fixed-sum ordering theorem for classical Markov numbers [2]. It does not assume classical Markov uniqueness, or positivity of the coefficients of the mirror factors.

1. Degree on the squared tree

Index the ordinary Farey tree in $[0, 1]$ by reduced fractions a/b. The endpoint labels are $0/1$ and $1/1$, and the first mediant is $1/2$. Write $m_{a/b}$ for its classical Markov label, with $m_{0/1} = 1$, $m_{1/1} = 2$, $m_{1/2} = 5$. These are labels at positions: we do not presume that different positions have different integer labels. Use the identical indexing for the symmetric squared deformation $F_{a/b}(q)$ and its mirror factor $f_{a/b}(q)$. The source establishes

$$F_{a/b}(q) = f_{a/b}(q) f_{a/b}(q^{-1}), \qquad f_{a/b}(1) = m_{a/b}, \tag{1}$$

and that every $F_{a/b}$ is a symmetric Laurent polynomial with nonnegative integer coefficients [1, Theorem 2.2, Section 3.1, Theorem 4.4 and Notation 4.5]. For a nonzero Laurent polynomial g let $\delta(g)$ be its largest exponent. For an ordinary polynomial $f \neq 0$, its exponent width is $w(f) = \deg f - \operatorname{val}_q f$. The endpoint terms give

$$\delta\big(f(q) f(q^{-1})\big) = w(f). \tag{2}$$

There is exactly one product contributing to the largest exponent in (2).

Lemma 2. *For every reduced $a/b \in [0, 1]$,*

$$\delta(F_{a/b}) = w(f_{a/b}) = a + b - 1. \tag{3}$$

At every noninitial squared triple the largest degree is the sum of the other two degrees plus one, and hence is strictly larger than both.

Proof. The first two triples are $(1, 1, 1)$ and $(1, 1, q + 2 + q^{-1})$. The latter has degrees $(0, 0, 1)$. Suppose a triple (A, B, C), ordered by its classical labels, has degrees (r, s, t) with $t = r + s + 1$ and $r, s \geq 0$. Mutation replacing A gives

$$AA' = B^2 + (q + q^{-1})BC + C^2. \tag{4}$$

Since $t \geq s+1$, the numerator has largest exponent $\max\{2s, s+t+1, 2t\} = 2t$. Its coefficients are nonnegative, so a tie cannot cancel. Therefore

$$\delta(A') = 2t - r = s + t + 1 > t.$$

Replacing B likewise produces degree $r + t + 1 > t$. Induction proves the asserted degree invariant on every branch. In particular the labels $0/1$, $1/1$, $1/2$ have degrees $0, 1, 2$. For a new Farey mediant $(a+c)/(b+d)$, the recurrence just proved is

$$\delta(F_{(a+c)/(b+d)}) = \delta(F_{a/b}) + \delta(F_{c/d}) + 1.$$

The proposed expression $a + b - 1$ satisfies precisely this recurrence and the endpoint values. Equation (2) finishes the proof. □

2. The normalization has nonzero constant term

This step identifies ordinary polynomial degree with the width in (3). It uses only the source's squared positivity and polynomiality. The initial mirror polynomials 1, $1 + q$, and $1 + 2q + 2q^2$ have nonzero constant term, as do their reversed choices. Assume this for the three entries of a triple. Write $d(q)$ for the entry being replaced, $h(q)$ for the other nonmaximum entry, and $c(q)$ for the maximum entry. Set $H = h(q)h(q^{-1})$ and $C = c(q)c(q^{-1})$. By the induction hypothesis and Lemma 2,

$$t := \deg c = \delta(C), \qquad \delta(H) \leq t - 1.$$

For either permitted sign $\epsilon \in \{-1, 1\}$ the source's mutation has the form

$$p(q) = \frac{q^t\left(q^\epsilon H + C\right)}{\widetilde{d}(q)}, \qquad \widetilde{d}(q) = q^{\deg d} d(q^{-1}). \tag{5}$$

The numerator is an ordinary polynomial: its lowest possible exponents are $t + \epsilon - \delta(H) \geq 0$ and $t - \delta(C) = 0$. Its constant coefficient is positive, since C contributes its positive endpoint coefficient and H cannot cancel it. The denominator has nonzero constant coefficient, namely the leading coefficient of d. The source's Theorem 4.4 says that p is a polynomial. Evaluating the polynomial identity in (5) at zero now gives $p(0) \neq 0$. This proves the induction. Consequently

$$\deg f_{a/b} = a + b - 1, \tag{6}$$

and the mediant entry is the unique maximum-degree polynomial of its triple. The argument applies to every fixed permitted initial choice.

3. Fixed-sum ordering separates all positions

We use the following published theorem of Lagisquet, Pelantová, Tavenas and Vuillon [2, Theorem 2(3), Proposition 22]: if $0 < i < a < b$, $\gcd(a, b) = 1$, and $\gcd(a - i, b + i) = 1$, then

$$m_{a/b} < m_{(a-i)/(b+i)}. \tag{7}$$

Thus distinct reduced fractions in $(0, 1)$ having the same numerator plus denominator have different Markov labels. Indeed, given two such fractions, choose a to be the larger numerator and let i be their positive numerator difference. The other numerator $a - i$ is positive, and equality of the

sums makes its denominator $b+i$; all hypotheses of (7) follow. The cited theorem is proved for an extension to noncoprime pairs as well, so it does not require intermediate fractions in a comparison chain to be reduced.

Suppose two interior positions give polynomials with equal degree and equal value at $q=1$. Equation (6) gives equal sums $a+b$; equation (1) gives equal classical Markov labels. By (7) their reduced fractions are equal. The ordinary Farey construction has exactly one vertex whose mediant is any given reduced fraction in $(0,1)$, so the two triples in the fixed mirror tree coincide. The exceptional initial triples $(1,1,1)$ and $(1,1,1+q)$ have maximum degrees 0 and 1, whereas every interior triple has maximum degree at least 2. They cause no collision with one another or with an interior triple. This proves Theorem 1.

The same reasoning proves injectivity of $a/b \mapsto F_{a/b}(q)$: equal squared polynomials give equal degrees and $F_{a/b}(1) = m_{a/b}^2$; positive classical labels then agree. This is a consequence of the same proof, not a second problem credit. In particular the proof does not infer the still separate classical uniqueness assertion by discarding the degree information.

Verification and scope

The accompanying exact verifier checks the squared mutation, Farey degree formula and independent Christoffel-matrix labels on a finite binary subtree; both initial mirror choices and their signed mutations; and the fixed-sum order for a bounded set of coprime pairs. These are supplementary checks. The inductions and the published theorem establish the unrestricted conclusion. Review is by the authoring agent only.

Proposed follow-up Tasks 2–5

These are research directions following the completed Task 1.

Task 2. Construct an algorithm recovering the Farey fraction from $(\deg f, f(1))$ by the fixed-sum ordering, and give bit-complexity bounds.

Task 3. Determine whether the degree and two endpoint coefficients already identify every mirror polynomial in a fixed tree.

Task 4. Derive a closed Christoffel-word formula for the leading and constant coefficients of $f_{a/b}$, including their dependence on the initial choice.

Task 5. For the squared recurrence with $q+q^{-1}$ replaced by a symmetric Laurent polynomial, characterize the substitutions for which a degree statistic and an integer specialization still separate all Farey positions.

References

[1] L. Bittmann, P. Jouteur, E. Kantarcı Oğuz, M. Molander and E. Yıldırım, *A mirror deformation of Markov numbers*, arXiv:2602.14802v1, 2026.

[2] C. Lagisquet, E. Pelantová, S. Tavenas and L. Vuillon, *On the Markov numbers: fixed numerator, denominator, and sum conjectures*, Advances in Applied Mathematics **130** (2021), 102227, doi:10.1016/j.aam.2021.102227. Theorem 2 and Proposition 22 in the complete preprint arXiv:2010.10335v1.