Hilbert coefficients of the square of the maximal ideal

Principia Math In Progress

22 September 2026

MathDB 360642. Full intended formula proved; zero new campaign credit. The coefficient identity is already a direct specialization of classical Gegenbauer formulas. We record that prior implication explicitly and supply elementary algebraic and combinatorial proofs. Neither the alternative proof nor the source's index correction earns new credit.

Task 1: the complete intended statement

Let $S_n = k[x_1, \dots, x_n]$, where k is any field and $n \geq 1$, and put $I = (x_1, \dots, x_n)^2$. Use the source's defining convention

$$\ell(S_n/I^s) = \sum_{j=0}^{n} (-1)^j e_j(I) \binom{s+n-j-1}{n-j} \qquad (s \gg 0). \tag{1}$$

Theorem 1. *For every $n \geq 1$,*

$$e_j(I) = \begin{cases} 2^{n-2j}\binom{n-j}{j}, & 0 \leq j \leq \lfloor n/2 \rfloor, \\ 0, & \lfloor n/2 \rfloor < j \leq n. \end{cases} \tag{2}$$

In particular the claimed nonzero coefficients are strictly positive. The corresponding identity, valid for every integer $s \geq 1$, is

$$\binom{2s+n-1}{n} = \sum_{j=0}^{\lfloor n/2 \rfloor} (-1)^j 2^{n-2j} \binom{n-j}{j} \binom{s+n-j-1}{n-j}. \tag{3}$$

The current preprint's Conjecture 3 and the published Conjecture 3.1 state the intended coefficient formula (2), but omit the final minus one in the upper entry of the last binomial in (3). Their own example $n = 4, s = 5$ uses the corrected convention. The uncorrected identity already fails at $n = 1$, giving $2s = 2s + 2$. This typographical issue is separate from proving (2) for every dimension.

An elementary proof in every dimension

The monomials of total degree less than $2s$ form a basis of S_n/I^s, so stars and bars gives its length as the left side of (3). In formal power series, for a fixed positive integer s,

$$(1-t)^{-2s} = (1-2t+t^2)^{-s} = \sum_{r \geq 0} \binom{s+r-1}{r} (2t-t^2)^r. \tag{4}$$

All substitutions are valid formally because $2t-t^2$ has zero constant term. Its r-th power contributes to degree n only when $r = n - j$ and $0 \le j \le \lfloor n/2 \rfloor$; that coefficient is $(-1)^j 2^{n-2j}\binom{n-j}{j}$. The coefficient of t^n on the left of (4) is $\binom{2s+n-1}{n}$, proving (3).

For fixed n, the polynomials $\binom{s+n-j-1}{n-j}$, $0 \le j \le n$, have distinct degrees $n-j$ and nonzero leading coefficients. They form a basis of the polynomials of degree at most n. Comparing (1) and (3) therefore gives every coefficient in (2), including the vanishing ones. This argument concerns integer dimensions and lengths and is valid independently of the characteristic of k.

A combinatorial proof by cancellation

Consider s ordered, possibly empty words made from tiles A, B, D. Tiles A, B each have size one and positive sign; D has size two and negative sign. Give a tuple of words total size n and sign $(-1)^{\#D}$. When it has j tiles D, it has $r = n - j$ tiles altogether. Choose the j domino locations, color the other tiles A or B, and distribute the ordered tile string among the s words by a weak composition of r. Its signed count is precisely the j-th summand of (3).

Scan the words in order and the tiles within each word from left to right. Find the first occurrence of either a tile D or an adjacent pair AB within one word. Replace

$$D \longleftrightarrow AB. \tag{5}$$

If no such occurrence exists, leave the tuple fixed. The replacement preserves size and flips the sign. It is an involution: the unchanged prefix has no earlier occurrence, and expanding D starts with A, so no new AB starts just before the replacement. Conversely contracting AB to D creates no earlier occurrence. The same place is consequently selected on applying (5) again. Word boundaries are never crossed.

Every fixed tuple has no domino and no AB in any word. Thus each word is uniquely $B^{b_i}A^{a_i}$, with $a_i, b_i \ge 0$. The fixed tuples are in bijection with the weak compositions

$$a_1 + b_1 + \cdots + a_s + b_s = n.$$

They all have positive sign, and their number is $\binom{n+2s-1}{n}$. Cancellation proves (3) for every n, s, and the polynomial-basis argument above completes (2).

The classical identity and counting decision

This result is already implied by the two classical identities

$$C_n^{(s)}(x) = \sum_{j=0}^{\lfloor n/2 \rfloor} \frac{(-1)^j (s)_{n-j}}{j!(n-2j)!}(2x)^{n-2j}, \qquad C_n^{(s)}(1) = \frac{(2s)_n}{n!}. \tag{6}$$

Here $(a)_r$ is the rising factorial. The first formula is NIST DLMF 18.5.10; the second follows immediately from 18.5.9 at $x = 1$, where the hypergeometric argument is zero. Since

$$\frac{(s)_{n-j}}{j!(n-2j)!} = \binom{n-j}{j}\binom{s+n-j-1}{n-j},$$

their specialization is exactly (3). DLMF 18.12.4 also gives the generating function behind (4). The relevant equations and parameter specializations were inspected directly. The full DLMF chapter and its cited books were not read.

The complete current and published problem sources still call this a conjecture. Nevertheless, the existing full identity above already implies every intended coefficient, so we conservatively exclude this package from new campaign credit. No claim of a new binomial identity or historical priority is made for either elementary proof.

Tasks 2–5: four proposed follow-ups

Task 2: Extend the explicit coefficient extraction to every ideal $(x_1, \ldots, x_n)^d$, with $d \geq 3$.

Task 3: Find a cancellation model for those higher-power coefficient formulas, with fixed objects explaining their signs.

Task 4: Derive exact Hilbert series for powers of ideals generated by all squares and the edges of an arbitrary finite graph.

Task 5: Determine which such graph ideals have nonnegative Hilbert coefficients in every dimension.

These are proposed mathematical directions, not additional solutions.

Sources and verification

R. Fröberg, *Hilbert coefficients of quadratic algebras*, arXiv:2311.02860v1, 6 November 2023, Conjecture 3, p. 5; all seven pages read. Published as *Journal of Experimental Mathematics* 1(2) (2025), 315–321, DOI 10.56994/JXM.001.002.007, Conjecture 3.1, p. 319; all seven article pages and both front-matter pages read.

NIST Digital Library of Mathematical Functions: 18.5.9, 18.5.10, and 18.12.4. Exact equations and their displayed context checked 22 September 2026.

Supplementary exact checks independently recover the Hilbert coefficients from lengths, enumerate the cancellation involution on small word tuples, and test the displayed index correction. Finite checks supplement the proofs; they do not prove the infinite claims. Review by the authoring agent only, without independent peer review or a proof assistant.