Unimodality for hedgehog trees with delays one and two

A complete proof for MathDB 360955; source audit pending

Principia Math In Progress

18 September 2026

Abstract

We prove the unimodality conjecture for the plucking polynomial of a star whose leaf delays belong to $\{1, 2\}$. The source's factorization reduces it to multiplying a polynomial of degree at most m by $[m]_q!$. One long quantum-integer factor gives unimodality, and each remaining factor preserves it by an elementary moving-sum argument. The proof extends to arbitrary nonnegative real coefficients in the initial polynomial.

Publication status. This note gives a full proof of the mathematical assertion in Conjecture 4.1 of the current arXiv preprint [1]. A published 2026 chapter exists [2], but its full text could not be retrieved for comparison. Accordingly, this result has **zero new campaign credit pending that source audit**. The preprint was read completely; novelty relative to the final chapter is not claimed.

Task 1: the original problem and its full solution

A finite nonnegative sequence is *unimodal* if it is weakly increasing up to some index and weakly decreasing thereafter. Initial or final zeros are permitted, as is the zero sequence. A polynomial is unimodal when its coefficient sequence is. Put

$$[r]_q = 1 + q + \cdots + q^{r-1}, \qquad [r]_q! = \prod_{j=1}^{r} [j]_q, \qquad [0]_q! = 1.$$

Let T be a plane star with n leaves. Label the leaves from right to left by $v_0, \ldots, v_{n-1}$ and give each a delay $f(v_i) \in \{1, 2\}$. Only delay-one leaves can be plucked at the first step. Plucking v_i contributes q^i, removes that leaf, and replaces every surviving delay by $\max(1, f(v) - 1)$. Thus all remaining delays are one, and the remaining star has plucking polynomial $[n-1]_q!$. The source's Proposition 2.5 is precisely the resulting formula

$$Q(T, f) = A(q)[n-1]_q!, \qquad A(q) = \sum_{i=0}^{n-1} \varepsilon_i q^i, \quad \varepsilon_i = \begin{cases} 1 & f(v_i) = 1, \\ 0 & f(v_i) = 2. \end{cases} \tag{1}$$

An empty set of eligible leaves gives $Q = 0$. The conjecture asks whether (1) is always unimodal. We credit (1) and the binary-coefficient case of the next lemma to Propositions 2.5 and 2.6 of [1].

Lemma 1. *For $m \geq 1$ and $A(q) = \sum_{j=0}^{m} a_j q^j$ with $a_j \geq 0$, the polynomial $A(q)[m]_q$ is unimodal.*

Proof. Write $A(q)[m]_q = \sum_{i=0}^{2m-1} b_i q^i$, allowing zero coefficients at the ends. Its coefficients satisfy

$$b_i = \begin{cases} a_0 + \cdots + a_i, & 0 \le i \le m-1, \\ a_{i-m+1} + \cdots + a_m, & m \le i \le 2m-1. \end{cases}$$

Hence $b_0 \le \cdots \le b_{m-1}$ and $b_m \ge \cdots \ge b_{2m-1}$. The single comparison between b_{m-1} and b_m can have either sign: in the first case the peak is at m, and in the other it is at $m-1$. More explicitly, $b_m - b_{m-1} = a_m - a_0$. This also covers $m = 1$. □

Lemma 2 (Moving sums)**.** *If $B(q)$ has a finite nonnegative unimodal coefficient sequence, then $B(q)[r]_q$ is unimodal for every integer $r \ge 1$.*

Proof. Extend the coefficients b_i by zero to every integer index, and choose a peak index p. For $B(q)[r]_q = \sum_i c_i q^i$, telescoping gives

$$c_i - c_{i-1} = b_i - b_{i-r}. \tag{2}$$

When $i \le p$, the right side is nonnegative, since both indices are in the increasing part of b. When $i \ge p + r$, it is nonpositive, since both are in the decreasing part. For $p \le i \le p + r$, the term b_i is nonincreasing in i and the term b_{i-r} is nondecreasing. Their difference is therefore nonincreasing throughout this middle interval. Thus the differences in (2) can change sign only from nonnegative to nonpositive. This is unimodality of (c_i). The all-zero sequence is immediate. □

Theorem 1. *If $m \ge 1$ and A is a polynomial of degree at most m with nonnegative real coefficients, then $A(q)[m]_q!$ is unimodal. Consequently $Q(T, f)$ is unimodal for every star and every delay function with values in $\{1, 2\}$.*

Proof. Apply Lemma 1 to $A(q)[m]_q$, padding the coefficients of A by zeros when necessary. Then multiply successively by $[m-1]_q, [m-2]_q, \ldots, [1]_q$, applying Lemma 2 each time. The product is $A(q)[m]_q!$. For a star with $n \ge 2$, take $m = n - 1$ in (1). For $n = 1$, the polynomial is 1 or 0 according to the delay. For the empty star it is 1 by the initial condition. □

Scope and attribution. The argument allows nonsymmetric delay patterns and arbitrary numbers of leaves. It does not require the delay function to be anti-unimodal. The first factor and the initial unimodality comparison are already in the source. The moving-sum lemma is an elementary preservation property, reproduced here in full; we do not claim it as a new general theorem. Applying it to the remaining quantum-integer factors completes the particular conjecture as stated in the preprint. The source's other delay families and general rooted trees are not included in this result.

Task 2: delays taking three values

Determine whether every star with leaf delays in $\{1, 2, 3\}$ has a unimodal plucking polynomial. After the first pluck some delays can still be two, so the common remaining factor in (1) no longer follows immediately. Seek a decomposition whose summands have compatible coefficient peaks, or an explicit counterexample.

Task 3: strict increase and the peak interval

For a nonempty eligible set $S \subseteq \{0, \dots, n-1\}$, determine the complete interval of maximal coefficients of $[n-1]_q! \sum_{i \in S} q^i$. Characterize the sets S for which the coefficients, after deleting end zeros, increase strictly and then decrease strictly, allowing at most a two-term peak.

Task 4: the largest universally smoothed degree

For each m, determine the largest $d(m)$ for which $A(q)[m]_q!$ is unimodal for every nonnegative polynomial A of degree at most $d(m)$. Theorem 1 proves $d(m) \geq m$. Study sharp obstructions from polynomials supported at two widely separated exponents and the growth of $d(m)$ with m.

Task 5: extending the argument beyond stars

For a general plane rooted tree with delays one and two, the first-step summands have different remaining tree shapes. Find structural hypotheses that force these summands to have compatible modes or a common sufficiently long quantum-integer factor. Establish such a criterion for an explicit family of trees with two levels of branching.

Verification. The accompanying script uses exact integers to compare the recursive definition with (1), check coefficient unimodality on finite exhaustive families, verify moving sums, and reproduce two already-known nonunimodal examples outside the delay-one-or-two hypothesis. The all-size proof above is independent of those checks. Review was by the authoring agent only. Tasks 2–5 are proposed questions, not claimed results.

References

[1] D. Ibarra, A. Landry, G. Montoya-Vega, and J. H. Przytycki, *Exploring unimodality of the plucking polynomial with delay function*, arXiv:2311.18059v1 (29 November 2023), 13 pages. Complete preprint read.

[2] The same authors, same title, Contemporary Mathematics **834** (2026), 95–108, doi:10.1090/conm/834/16686. Publication metadata verified; full chapter not retrieved or read.