



The Influence of Random Effects on the Unconditional Hazard Rate and Survival Functions

Yasuhiro Omori; Richard A. Johnson

Biometrika, Vol. 80, No. 4. (Dec., 1993), pp. 910-914.

Stable URL:

<http://links.jstor.org/sici?sici=0006-3444%28199312%2980%3A4%3C910%3ATIOREO%3E2.0.CO%3B2-M>

Biometrika is currently published by Biometrika Trust.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/bio.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact support@jstor.org.

The influence of random effects on the unconditional hazard rate and survival functions

BY YASUHIRO OMORI

Faculty of Law and Economics, Chiba University, Chiba 263, Japan

AND RICHARD A. JOHNSON

Department of Statistics, University of Wisconsin, Madison, Wisconsin 53703, U.S.A.

SUMMARY

The consequences of introducing random effects have been a major issue in models for survival analysis. A general result is derived to establish the underestimation of the unconditional hazard rate when a random effect is present. Also, the survival time is shown to be underestimated. No distributional assumption regarding the random effect is required.

Some key words: Hazard rate; Random effects; Underestimation.

1. INTRODUCTION

The ‘random effects survival model’ has been a major focus in the recent literature (Heckman & Singer, 1984; Littlewood, 1984; Clayton & Cuzick, 1985; Heckman, Robb & Walker, 1990; Honoré, 1990). When a hazard rate function is estimated, it is conventional practice to assume that observations are taken from a homogeneous group. If they are based on several heterogeneous subgroups, we should estimate their hazard rate functions separately. However, if we could assume that the hazard rate function from one group is parallel to those of other groups, we can deal with all observations at the same time by taking account of covariate variables by setting

$$\lambda_N(t) = \lambda_0(t) \exp(\beta'x),$$

where $\lambda_0(t)$ is a specified hazard rate, x is a vector of fixed covariates, and the subscript N is attached to emphasize that this is a nominal hazard rate, i.e. without heterogeneity. This helps to describe the differences among subgroups. But, if we want to describe all differences exactly, we have to use too many covariates. This is not realistic since the number of observations available is limited. Thus we choose several important covariates and regard the remaining covariates as random.

In order to introduce heterogeneity into the population via a random effect, we let the hazard rate for an individual be $V\lambda_N(t)$ or

$$V\lambda_0(t) \exp(\beta'x),$$

where V is a random variable such that $\text{pr}(V \geq 0) = 1$, $E(V) = 1$ and $\text{pr}(V = 1) < 1$. Conditional on $V = v$, an individual hazard rate is $\lambda_C(t|v) = v\lambda_N(t)$. The corresponding cumulative distribution functions are:

$$F_C(t|v) = 1 - \exp\{-v\Lambda_N(t)\}, \quad F_N(t) = 1 - \exp\{-\Lambda_N(t)\}, \quad F_U(t) = 1 - E[\exp\{-V\Lambda_N(t)\}], \quad (1.1)$$

where $\Lambda_N(t) = \int \lambda_N(s) ds$ with the integral over the range $(0, t)$, and the subscripts C , N , U denote the conditional, nominal, and unconditional distributions, respectively. The expectations always exist since $V \geq 0$ and $\Lambda_N(t) \geq 0$. Using $F_U(t)$ and $F_C(t|v)$, by taking an expected value of $f_C(t|V)$,

the unconditional hazard rate is

$$\lambda_U(t) = \lambda_N(t) \frac{E[V \exp \{-V\Lambda_N(t)\}]}{E[\exp \{-V\Lambda_N(t)\}]} \quad (1.2)$$

Lancaster (1979) originated the random effects model in the analysis of unemployment data. He used a Weibull hazard rate function and found that the maximum likelihood estimate $\hat{\delta}$ in the hazard function $\lambda_0(t) = \delta t^{\delta-1}$ increases with an increase of the number of covariates. He showed that, if V has a gamma distribution with mean 1 and variance c , then the unconditional hazard rate is

$$\lambda_U(t) = \{1 - F_U(t)\}^c \lambda_N(t) < \lambda_N(t).$$

If the heterogeneity modelled by V is ignored, the hazard rate function is underestimated. Using this model, Lancaster (1985) proved the inconsistency of maximum likelihood estimators when heterogeneity is ignored. Littlewood (1984) mentioned that mixtures of increasing failure rate distributions do not necessarily have an increasing failure rate even though mixtures of decreasing failure rate distributions have a decreasing failure rate.

A gamma heterogeneity is often used because it is easy to manipulate. Different distributions for heterogeneity such as an inverse Gaussian distribution also need to be considered, but they are not discussed in the literature. Heckman & Singer (1984) used some simulation results to suggest that estimators for the structural parameters, β , are very sensitive to the assumption of a specific heterogeneity distribution. Their suggestion started a discussion on the robustness of the estimators under heterogeneity. Nonparametric estimation was used to solve this problem (Heckman & Singer, 1984; Clayton & Cuzick, 1985; Honoré, 1990). In the next section, we generalize Lancaster (1979).

2. EFFECTS OF HETEROGENEITY ON THE UNCONDITIONAL HAZARD RATE AND SURVIVAL FUNCTIONS

If the heterogeneity modelled by V exists and is ignored, the hazard rate function is underestimated regardless of the heterogeneity distribution. We summarize our results in the following theorem, corollary and lemma. Proofs are given in the Appendix.

THEOREM 2.1. *Let T be a nonnegative random variable with probability density function $f_U(t)$. Suppose that $\text{pr}(V \geq 0) = 1$, $E(V) = 1$ and $\text{pr}(V = 1) < 1$. Then*

$$\lambda_U(t) - \lambda_N(t) = \lambda_N(t) \{E(V|T > t) - 1\}, \quad (2.1)$$

$$\frac{d}{dt} E(V|T > t) = -\text{var}(V|T > t), \quad \lambda_N(t) > 0, \quad (2.2)$$

and so $E(V|T > t)$ is nonincreasing in $t > 0$, with $E(V|T > 0) = E(V) = 1$.

Equations (2.1) and (2.2) show that the unconditional hazard rate $\lambda_U(t)$ is less than the nominal hazard rate $\lambda_N(t)$ at every point $t > 0$ where $\lambda_N(t) > 0$. They also imply that the difference between two hazard rates goes to 0 as $\text{var}(V)$ goes to 0. If $t_0 = \inf\{t: \lambda_N(t) > 0\}$ then the conditional expected value goes to 1 as t decreases to t_0 . Consequently, $\lambda_U(t)$ is a decreasing multiple of $\lambda_N(t)$. Also, if $F_N(t) = 1$ for some finite value t , then $f_N(t)$ and hence $\lambda_N(t)$ are 0 for that t and all larger values.

COROLLARY 2.1. *Let T be a nonnegative random variable with a probability density function $f_U(t)$. Suppose that $\text{pr}(V \geq 0) = 1$, $E(V) = 1$ and $\text{pr}(V = 1) < 1$. Then the unconditional survival time is stochastically larger than the nominal survival time. That is,*

$$F_U(t) < F_N(t) \quad \text{for } t > 0 \text{ and } \lambda_N(t) > 0, \quad (2.3)$$

and $F_U(t) \leq F_N(t)$ otherwise.

LEMMA 2.1. Let T be a nonnegative random variable with a probability density function $f_V(t)$. Suppose that $\text{pr}(V \geq 0) = 1$, $E(V) = 1$ and $\text{pr}(V = 1) < 1$. If $E\{\exp(uV)\}$ is finite for u in the neighbourhood of 0, then it has a series expansion

$$\lambda_V(t) = \lambda_N(t) \sum_{j=1}^{\infty} \frac{\kappa_j}{(j-1)!} \{-\Lambda_N(t)\}^{j-1}, \quad (2.4)$$

in the neighbourhood of 0, where κ_j is the j th cumulant of V .

Equation (2.4) shows that a large value of $\kappa_2 = \text{var}(V)$ contributes to the underestimation of hazard rates.

Example. Assume that V has a gamma distribution with mean 1 and variance c . Then

$$\lambda_V(t) = \lambda_N(t) \{1 + c\Lambda_N(t)\}^{-1}.$$

For example, consider an increasing failure rate, $\lambda_N(t) = 2t$. Then the mixture of these increasing failure rate distributions has an increasing rate for $t < c^{-\frac{1}{2}}$ and a decreasing failure rate for $t > c^{-\frac{1}{2}}$.




Fig. 1. Comparison of unconditional hazard rates, $\lambda_V(t)$, for different random effects given variance; solid lines, inverse Gaussian; dotted lines, gamma.




Fig. 2. Comparison of hazard rates, $\lambda_V(t)$, for different random effects given random effects; solid lines, $\text{var}(V) = 0.1$; dotted lines, $\text{var}(V) = 1.0$.

Similarly, if V has an inverse Gaussian distribution with mean 1 and variance c , then

$$\lambda_V(t) = \lambda_N(t) \{1 + 2c\Lambda_N(t)\}^{-\frac{1}{2}}.$$

For example, if $\lambda_N(t) = 2t$, the mixture of these increasing failure rate distributions has an increasing failure rate again, but $\lambda_V(t) < (2/c)^{\frac{1}{2}}$. Figures 1 and 2 give examples for the gamma and inverse Gaussian distributions with $\lambda_N(t) = 1$ and $\lambda_N(t) = 2t$ where the effect is more serious for larger t values. Also, the effect of heterogeneity is more serious for the gamma distributions.

ACKNOWLEDGEMENT

The authors would like to thank the referee for helpful comments.

APPENDIX

Proofs

Proof of Theorem 2.1. Let $I(T > t)$ be the indicator function of the event $T > t$. Then

$$E\{VI(T > t)\} = E[VE\{I(T > t) | V = v\}] = E[V \exp\{-V\Lambda_N(t)\}],$$

$$E\{I(T > t)\} = E[E\{I(T > t) | V = v\}] = E[\exp\{-V\Lambda_N(t)\}].$$

Hence, by (1.2),

$$\begin{aligned}\lambda_V(t) - \lambda_N(t) &= \lambda_N(t) \left(\frac{E[V \exp \{-V\Lambda_N(t)\}]}{E[\exp \{-V\Lambda_N(t)\}]} - 1 \right) \\ &= \lambda_N(t) \left[\frac{E\{VI(T>t)\}}{E\{I(T>t)\}} - 1 \right] = \lambda_N(t) \{E(V|T>t) - 1\}.\end{aligned}$$

For all t where $\Lambda_N(t) > 0$, $\exp \{-v\Lambda_N(t)\}$, $v \exp \{-v\Lambda_N(t)\}$ and $v^2 \exp \{-v\Lambda_N(t)\}$ are uniformly bounded. By the dominated convergence theorem,

$$\frac{d}{dt} E(V|T>t) = - \frac{E[V^2 \exp \{-V\Lambda_N(t)\}]}{E[\exp \{-V\Lambda_N(t)\}]} + \frac{E^2[V \exp \{-V\Lambda_N(t)\}]}{E^2[\exp \{-V\Lambda_N(t)\}]} = -\text{var}(V|T>t).$$

The result follows. \square

Proof of Corollary 2.1. By Theorem 2.1, $\lambda_V(t) < \lambda_N(t)$ for $t > 0$ and $\Lambda_N(t) > 0$. Hence, for any such t ,

$$F_V(t) = 1 - \exp \left\{ - \int_0^t \lambda_V(s) ds \right\} < 1 - \exp \left\{ - \int_0^t \lambda_N(s) ds \right\} = F_N(t). \quad \square$$

Proof of Lemma 2.1. When $E\{\exp(uV)\}$ is finite for all u in a neighbourhood of 0,

$$\psi(u) \equiv \log E\{\exp(uV)\} = \sum_{j=1}^{\infty} \frac{\kappa_j}{j!} u^j.$$

Since

$$\log E[\exp \{-V\Lambda_N(t)\}] = \psi\{-\Lambda_N(t)\},$$

by the dominated convergence theorem,

$$\lambda_V(t) = \lambda_N(t) \frac{E[V \exp \{-V\Lambda_N(t)\}]}{E[\exp \{-V\Lambda_N(t)\}]} = \lambda_N(t) \psi' \{-\Lambda_N(t)\} = \lambda_N(t) \sum_{j=1}^{\infty} \frac{\kappa_j}{(j-1)!} \{-\Lambda_N(t)\}^{j-1},$$

in the neighbourhood of 0 and (2.4) follows. \square

REFERENCES

- CLAYTON, D. & CUZICK, J. (1985). Multivariate generalization of the proportional hazard model. *J. R. Statist. Soc. A* **148**, 82–117.
- HECKMAN, J. J., ROBB, R. & WALKER, J. R. (1990). Testing the mixture of exponentials hypothesis and estimating the mixing distribution by the method of moments. *J. Am. Statist. Assoc.* **85**, 582–90.
- HECKMAN, J. J. & SINGER, B. (1984). A method for minimizing the impact of distributional assumptions in econometric models for a duration data. *Econometrica* **52**, 271–320.
- HONORÉ, B. E. (1990). Simple estimation of a duration model with unobserved heterogeneity. *Econometrica* **58**, 453–73.
- LANCASTER, T. (1979). Econometric methods for duration of unemployment. *Econometrica* **47**, 939–55.
- LANCASTER, T. (1985). Generalized residuals and heterogeneous duration models with application to the Weibull model. *J. Economet.* **28**, 155–69.
- LITTLEWOOD, B. (1984). Subjective probability and the DFR-mixture closure theorem. *Commun. Statist. A* **13**, 859–63.

[Received December 1992. Revised May 1993]

LINKED CITATIONS

- Page 1 of 2 -



You have printed the following article:

The Influence of Random Effects on the Unconditional Hazard Rate and Survival Functions

Yasuhiro Omori; Richard A. Johnson

Biometrika, Vol. 80, No. 4. (Dec., 1993), pp. 910-914.

Stable URL:

<http://links.jstor.org/sici?sici=0006-3444%28199312%2980%3A4%3C910%3ATIOREO%3E2.0.CO%3B2-M>

This article references the following linked citations. If you are trying to access articles from an off-campus location, you may be required to first logon via your library web site to access JSTOR. Please visit your library's website or contact a librarian to learn about options for remote access to JSTOR.

References

Multivariate Generalizations of the Proportional Hazards Model

David Clayton; Jack Cuzick

Journal of the Royal Statistical Society. Series A (General), Vol. 148, No. 2. (1985), pp. 82-117.

Stable URL:

<http://links.jstor.org/sici?sici=0035-9238%281985%29148%3A2%3C82%3AMGOTPH%3E2.0.CO%3B2-Y>

Testing the Mixture of Exponentials Hypothesis and Estimating the Mixing Distribution by the Methods of Moments

James J. Heckman; Richard Robb; James R. Walker

Journal of the American Statistical Association, Vol. 85, No. 410. (Jun., 1990), pp. 582-589.

Stable URL:

<http://links.jstor.org/sici?sici=0162-1459%28199006%2985%3A410%3C582%3ATTMOEH%3E2.0.CO%3B2-U>

A Method for Minimizing the Impact of Distributional Assumptions in Econometric Models for Duration Data

J. Heckman; B. Singer

Econometrica, Vol. 52, No. 2. (Mar., 1984), pp. 271-320.

Stable URL:

<http://links.jstor.org/sici?sici=0012-9682%28198403%2952%3A2%3C271%3AAMFMTI%3E2.0.CO%3B2-U>

LINKED CITATIONS

- Page 2 of 2 -



Simple Estimation of a Duration Model with Unobserved Heterogeneity

Bo E. Honoré

Econometrica, Vol. 58, No. 2. (Mar., 1990), pp. 453-473.

Stable URL:

<http://links.jstor.org/sici?sici=0012-9682%28199003%2958%3A2%3C453%3ASEOADM%3E2.0.CO%3B2-0>

Econometric Methods for the Duration of Unemployment

Tony Lancaster

Econometrica, Vol. 47, No. 4. (Jul., 1979), pp. 939-956.

Stable URL:

<http://links.jstor.org/sici?sici=0012-9682%28197907%2947%3A4%3C939%3AEMFTDO%3E2.0.CO%3B2-7>