

[Courseware \(/courses/MITx/15.071x/1T2014/courseware/\)](/courses/MITx/15.071x/1T2014/courseware/)[Course Info \(/courses/MITx/15.071x/1T2014/info/\)](/courses/MITx/15.071x/1T2014/info/)[Discussion \(/courses/MITx/15.071x/1T2014/discussion/forum/\)](/courses/MITx/15.071x/1T2014/discussion/forum/)[Progress \(/courses/MITx/15.071x/1T2014/progress/\)](/courses/MITx/15.071x/1T2014/progress/)[Syllabus \(/courses/MITx/15.071x/1T2014/4264e68418f34d839cf0b33a5da644b2/\)](/courses/MITx/15.071x/1T2014/4264e68418f34d839cf0b33a5da644b2/)[Schedule \(/courses/MITx/15.071x/1T2014/2891f8bf120945b9aa12e6601739c3e6/\)](/courses/MITx/15.071x/1T2014/2891f8bf120945b9aa12e6601739c3e6/)

Help

CLIMATE CHANGE

There have been many studies documenting that the average global temperature has been increasing over the last century. The consequences of a continued rise in global temperature will be dire. Rising sea levels and an increased frequency of extreme weather events will affect billions of people.

In this problem, we will attempt to study the relationship between average global temperature and several other factors.

The file `climate_change.csv` (`/c4x/MITx/15.071x/asset/climate_change.csv`) contains climate data from May 1983 to December 2008. The available variables include:

- *Year*: the observation year.
- *Month*: the observation month.
- *Temp*: the difference in degrees Celsius between the average global temperature in that period and a reference value. This data comes from the Climatic Research Unit at the University of East Anglia (<http://www.cru.uea.ac.uk/cru/data/temperature/>).
- *CO2, N2O, CH4, CFC.11, CFC.12*: atmospheric concentrations of carbon dioxide (CO₂), nitrous oxide (N₂O), methane (CH₄), trichlorofluoromethane (CCl₃F; commonly referred to as CFC-11) and dichlorodifluoromethane (CCl₂F₂; commonly referred to as CFC-12), respectively. This data comes from the ESRL/NOAA Global Monitoring Division (<http://www.esrl.noaa.gov/gmd/ccgg/data-products.html>).
 - CO₂, N₂O and CH₄ are expressed in ppmv (parts per million by volume -- i.e., 397 ppmv of CO₂ means that CO₂ constitutes 397 millionths of the total volume of the atmosphere)
 - CFC.11 and CFC.12 are expressed in ppbv (parts per billion by volume).
- *Aerosols*: the mean stratospheric aerosol optical depth at 550 nm. This variable is linked to volcanoes, as volcanic eruptions result in new particles being added to the atmosphere, which affect how much of the sun's energy is reflected back into space. This data is from the Godard Institute for Space Studies at NASA (<http://data.giss.nasa.gov/modelforce/strataer/>).
- *TSI*: the total solar irradiance (TSI) in W/m² (the rate at which the sun's energy is deposited per unit area). Due to sunspots and other solar phenomena, the amount of energy that is given off by the sun varies substantially with time. This data is from the SOLARIS-HEPPA project website (<http://solarisheppa.geomar.de/solarisheppa/cmip5>).
- *MEI*: multivariate El Nino Southern Oscillation index (MEI), a measure of the strength of the El Nino/La Nina-Southern Oscillation (http://en.wikipedia.org/wiki/El_nino) (a weather effect in the Pacific Ocean that affects global temperatures). This data comes from the ESRL/NOAA Physical Sciences Division (<http://www.esrl.noaa.gov/psd/enso/mei/table.html>).

PROBLEM 1.1 - CREATING OUR FIRST MODEL (1/1 point)

We are interested in how changes in these variables affect future temperatures, as well as how well these variables explain temperature changes so far. To do this, first read the dataset `climate_change.csv` into R.

Then, split the data into a *training set*, consisting of all the observations up to and including 2006, and a *testing set* consisting of the remaining years (hint: use `subset`). A training set refers to the data that will be used to build the model (this is the data we give to the `lm()` function), and a testing set refers to the data we will use to test our predictive ability.

Next, build a linear regression model using all of the independent variables (except *Year* and *Month*) to predict the dependent variable *Temp*. Use the training set to build the model.

Enter the model R^2 :

Answer: 0.75

EXPLANATION

First, read in the data and split it using the `subset` command:

```
climate = read.csv("climate_change.csv")
```

```
train = subset(climate, Year <= 2006)
```

```
test = subset(climate, Year > 2006)
```

Then, create the model using the command:

```
climatelm = lm(Temp ~ MEI + CO2 + CH4 + N2O + CFC.11 + CFC.12 + TSI + Aerosols, data=train)
```







Lastly, look at the model using `summary(climatelm)`. The Multiple R-squared value is 0.7509.

Hide Answer

You have used 1 of 3 submissions

PROBLEM 1.2 - CREATING OUR FIRST MODEL (1/1 point)

Which variables are significant in the model? We will consider a variable significant only if the p-value is below 0.05.

- ☒ MEI 
- ☒ CO2 
- ☐ CH4
- ☐ N2O
- ☒ CFC.11 
- ☒ CFC.12 
- ☒ TSI 
- ☒ Aerosols 

EXPLANATION

If you look at the model we created in the previous problem using `summary(climate_lm)`, all of the variables have at least one star except for CH4 and N2O. So MEI, CO2, CFC.11, CFC.12, TSI, and Aerosols are all significant.


Hide Answer

You have used 1 of 2 submissions

PROBLEM 2.1 - UNDERSTANDING THE MODEL (1/1 point)

Current scientific opinion is that nitrous oxide and CFC-11 are greenhouse gases: gases that are able to trap heat from the sun and contribute to the heating of the Earth. However, the regression coefficients of both the N2O and CFC-11 variables are **negative**, indicating that increasing atmospheric concentrations of either of these two compounds is associated with lower global temperatures.

Which of the following is the *simplest* correct explanation for this contradiction?

- ☐ Climate scientists are wrong that N2O and CFC-11 are greenhouse gases - this regression analysis constitutes part of a disproof.
- ☐ There is not enough data, so the regression coefficients being estimated are not accurate.
- ☒ All of the gas concentration variables reflect human development - N2O and CFC.11 are correlated with other variables in the data set. 

EXPLANATION

The linear correlation of N2O and CFC.11 with other variables in the data set is quite large. The first explanation does not seem correct, as the warming effect of nitrous oxide and CFC-11 are well documented, and our regression analysis is not enough to disprove it. The second explanation is unlikely, as we have estimated eight coefficients and the intercept from 284 observations.

Hide Answer

You have used 1 of 1 submissions

PROBLEM 2.2 - UNDERSTANDING THE MODEL (2/2 points)

Compute the correlations between all the variables in the training set. Which of the following independent variables is N2O highly correlated with (absolute correlation greater than 0.7)?

- ☐ MEI
- ☒ CO2 ✓
- ☒ CH4 ✓
- ☐ CFC.11
- ☒ CFC.12 ✓
- ☐ Aerosols
- ☐ TSI

Which of the following independent variables is CFC.11 highly correlated with?

- ☐ MEI
- ☐ CO2
- ☒ CH4 ✓
- ☐ N2O
- ☒ CFC.12 ✓
- ☐ Aerosols
- ☐ TSI

EXPLANATION

You can calculate all correlations at once using `cor(train)` where `train` is the name of the training data set.

[Hide Answer](#)

You have used 1 of 2 submissions

PROBLEM 3 - SIMPLIFYING THE MODEL (2/2 points)

Given that the correlations are so high, let us focus on the N2O variable and build a model with only MEI, TSI, Aerosols and N2O. Remember to use the training set to build the model.

Enter the coefficient of N2O in this reduced model:

Answer: 0.02532

(How does this compare to the coefficient in the previous model with all of the variables?)

Enter the model R^2 :

0.7261

Answer: 0.7261

EXPLANATION

We can create this simplified model with the command:

```
LinReg = lm(Temp ~ MEI + N2O + TSI + Aerosols, data=train)
```

You can get the coefficient for N2O and the model R-squared by typing `summary(LinReg)`.

We have observed that, for this problem, when we remove many variables the sign of N2O flips. The model has not lost a lot of explanatory power (the model R^2 is 0.7261 compared to 0.7509 previously) despite removing many variables. As discussed in lecture, this type of behavior is typical when building a model where many of the independent variables are highly correlated with each other. In this particular problem many of the variables (CO₂, CH₄, N₂O, CFC.11 and CFC.12) are highly correlated, since they are all driven by human industrial development.

Hide Answer

You have used 1 of 3 submissions

PROBLEM 4 - AUTOMATICALLY BUILDING THE MODEL (2/2 points)

We have many variables in this problem, and as we have seen above, dropping some from the model does not decrease model quality. R provides a function, `step` (<http://stat.ethz.ch/R-manual/R-devel/library/stats/html/step.html>), that will automate the procedure of trying different combinations of variables to find a good compromise of model simplicity and R^2 . This trade-off is formalized by the Akaike information criterion (http://en.wikipedia.org/wiki/Akaike_information_criterion) (AIC) - it can be informally thought of as the quality of the model with a penalty for the number of variables in the model.


The `step` function has one argument - the name of the initial model. It returns a simplified model. Use the `step` function in R to derive a new model, with the full model as the initial model (HINT: If your initial full model was called "climateLM", you could create a new model with the `step` function by typing `step(climateLM)`. Be sure to save your new model to a variable name so that you can look at the summary. For more information about the `step` function, type `?step` in your R console.)

Enter the R^2 value of the model produced by the `step` function:

0.7508

Answer: 0.7508

Which of the following variable(s) were eliminated from the full model by the step function?

- ☐ MEI
- ☐ CO2
- ☒ CH4 
- ☐ N2O
- ☐ CFC.11
- ☐ CFC.12
- ☐ TSI
- ☐ Aerosols

EXPLANATION

You can create a model using the step function by typing:

```
StepModel = step(climateLM)
```

where "climateLM" is the name of the full model.

If you look at the summary of the model with `summary(StepModel)`, you can see that the R-squared value is 0.75, and only CH4 was removed.

It is interesting to note that the step function does not address the collinearity of the variables, except that adding highly correlated variables will not improve the R^2 significantly. The consequence of this is that the step function will not necessarily produce a very interpretable model - just a model that has balanced quality and simplicity for a particular weighting of quality and simplicity (AIC).

Hide Answer

You have used 1 of 4 submissions

PROBLEM 5 - TESTING ON UNSEEN DATA (1 point possible)

We have developed an understanding of how well we can fit a linear regression to the training data, but does the model quality hold when applied to unseen data?

Using the model produced from the step function, calculate temperature predictions for the testing data set, using the predict function.

Enter the testing set R^2 :

0.6286051

Answer: 0.6286051

EXPLANATION

The R code to calculate the R-squared can be written as follows (your variable names may be different):

```
tempPredict = predict(climateStep, newdata = test)
```

```
SSE = sum((tempPredict - test$Temp)^2)
```

```
SST = sum( (mean(train$Temp) - test$Temp)^2)
```

```
R2 = 1 - SSE/SST
```

[Hide Answer](#)

You have used 3 of 3 submissions

Please remember not to ask for or post complete answers to homework questions in this discussion forum.

[Show Discussion](#)
[New Post](#)


About (<https://www.edx.org/about-us>) Jobs (<https://www.edx.org/jobs>)
 Press (<https://www.edx.org/press>) FAQ (<https://www.edx.org/student-faq>)
 Contact (<https://www.edx.org/contact>)



EdX is a non-profit created by founding partners Harvard and MIT whose mission is to bring the best of higher education to students of all ages anywhere in the world, wherever there is Internet access. EdX's free online MOOCs are interactive and subjects include computer science, public health, and artificial intelligence.



(<http://www.meetup.com/edX-Global-Community/>)



(<http://www.facebook.com/EdxOnline>)



(<https://twitter.com/edXOnline>)



(<https://plus.google.com/108235383044095082735/posts>)



(<http://youtube.com/user/edxonline>)

© 2014 edX, some rights reserved.

Terms of Service and Honor Code -
 Privacy Policy (<https://www.edx.org/edx-privacy-policy>)