CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 1 OF 12]:

[GYN-CERV_03-P01-C0001] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IA1 [Microscopic Stromal Invasion < 3 mm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P01-C0002] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IA2 [Stromal Invasion 3 mm to < 5 mm] | Modality: Systemic Immunotherapy & Chemotherapy
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P01-C0003] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IB1 [Invasive Carcinoma >= 5 mm to < 2 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P01-C0004] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IB2 [Invasive Carcinoma >= 2 cm to < 4 cm] | Modality: Systemic Immunotherapy & Chem...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P01-C0005] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IB3 [Invasive Carcinoma >= 4 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P01-C0006] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIA1 [Vaginal Involvement Upper 2/3, < 4 cm] | Modality: Systemic Immunotherapy & Ch...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P01-C0007] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IIA2 [Vaginal Involvement Upper 2/3, >= 4 cm] | Modality: Concurrent Chemoradiation ...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P01-C0008] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIB [Parametrial Extension without Pelvic Wall] | Modality: Systemic Immunotherapy &...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P01-C0009] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IIIA [Lower Third Vagina Involvement] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P01-C0010] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIIB [Pelvic Sidewall Extension or Hydronephrosis] | Modality: Systemic Immunotherap...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 2 OF 12]:

[GYN-CERV_03-P02-C0023] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIIA [Lower Third Vagina Involvement] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P02-C0024] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIIB [Pelvic Sidewall Extension or Hydronephrosis] | Modality: Systemic Immunotherap...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P02-C0025] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IIIC1 [Pelvic Lymph Node Metastasis] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P02-C0026] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIIC2 [Para-Aortic Lymph Node Metastasis] | Modality: Systemic Immunotherapy & Chemo...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P02-C0027] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IVA [Mucosal Invasion of Bladder or Rectum] | Modality: Concurrent Chemoradiation [C...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P02-C0028] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IVB [Distant Hematogenous or Peritoneal Metastasis] | Modality: Systemic Immunothera...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P02-C0029] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IA1 [Microscopic Stromal Invasion < 3 mm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P02-C0030] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IA2 [Stromal Invasion 3 mm to < 5 mm] | Modality: Systemic Immunotherapy & Chemotherapy
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P02-C0031] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IB1 [Invasive Carcinoma >= 5 mm to < 2 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P02-C0032] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IB2 [Invasive Carcinoma >= 2 cm to < 4 cm] | Modality: Systemic Immunotherapy & Chem...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 3 OF 12]:

[GYN-CERV_03-P03-C0045] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IB1 [Invasive Carcinoma >= 5 mm to < 2 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P03-C0046] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IB2 [Invasive Carcinoma >= 2 cm to < 4 cm] | Modality: Systemic Immunotherapy & Chem...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P03-C0047] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IB3 [Invasive Carcinoma >= 4 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P03-C0048] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIA1 [Vaginal Involvement Upper 2/3, < 4 cm] | Modality: Systemic Immunotherapy & Ch...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P03-C0049] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IIA2 [Vaginal Involvement Upper 2/3, >= 4 cm] | Modality: Concurrent Chemoradiation ...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P03-C0050] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIB [Parametrial Extension without Pelvic Wall] | Modality: Systemic Immunotherapy &...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P03-C0051] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IIIA [Lower Third Vagina Involvement] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P03-C0052] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIIB [Pelvic Sidewall Extension or Hydronephrosis] | Modality: Systemic Immunotherap...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P03-C0053] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIIC1 [Pelvic Lymph Node Metastasis] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P03-C0054] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIIC2 [Para-Aortic Lymph Node Metastasis] | Modality: Systemic Immunotherapy & Chemo...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 4 OF 12]:

[GYN-CERV_03-P04-C0067] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IIIC1 [Pelvic Lymph Node Metastasis] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P04-C0068] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIIC2 [Para-Aortic Lymph Node Metastasis] | Modality: Systemic Immunotherapy & Chemo...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P04-C0069] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IVA [Mucosal Invasion of Bladder or Rectum] | Modality: Concurrent Chemoradiation [C...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P04-C0070] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IVB [Distant Hematogenous or Peritoneal Metastasis] | Modality: Systemic Immunothera...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P04-C0071] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IA1 [Microscopic Stromal Invasion < 3 mm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P04-C0072] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IA2 [Stromal Invasion 3 mm to < 5 mm] | Modality: Systemic Immunotherapy & Chemotherapy
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P04-C0073] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IB1 [Invasive Carcinoma >= 5 mm to < 2 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P04-C0074] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IB2 [Invasive Carcinoma >= 2 cm to < 4 cm] | Modality: Systemic Immunotherapy & Chem...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P04-C0075] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IB3 [Invasive Carcinoma >= 4 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P04-C0076] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIA1 [Vaginal Involvement Upper 2/3, < 4 cm] | Modality: Systemic Immunotherapy & Ch...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 5 OF 12]:

[GYN-CERV_03-P05-C0089] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IB3 [Invasive Carcinoma >= 4 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P05-C0090] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIA1 [Vaginal Involvement Upper 2/3, < 4 cm] | Modality: Systemic Immunotherapy & Ch...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P05-C0091] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IIA2 [Vaginal Involvement Upper 2/3, >= 4 cm] | Modality: Concurrent Chemoradiation ...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P05-C0092] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIB [Parametrial Extension without Pelvic Wall] | Modality: Systemic Immunotherapy &...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P05-C0093] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IIIA [Lower Third Vagina Involvement] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P05-C0094] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIIB [Pelvic Sidewall Extension or Hydronephrosis] | Modality: Systemic Immunotherap...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P05-C0095] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIIC1 [Pelvic Lymph Node Metastasis] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P05-C0096] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIIC2 [Para-Aortic Lymph Node Metastasis] | Modality: Systemic Immunotherapy & Chemo...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P05-C0097] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IVA [Mucosal Invasion of Bladder or Rectum] | Modality: Concurrent Chemoradiation [C...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P05-C0098] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IVB [Distant Hematogenous or Peritoneal Metastasis] | Modality: Systemic Immunothera...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 6 OF 12]:

[GYN-CERV_03-P06-C0111] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IVA [Mucosal Invasion of Bladder or Rectum] | Modality: Concurrent Chemoradiation [C...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P06-C0112] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IVB [Distant Hematogenous or Peritoneal Metastasis] | Modality: Systemic Immunothera...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P06-C0113] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IA1 [Microscopic Stromal Invasion < 3 mm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P06-C0114] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IA2 [Stromal Invasion 3 mm to < 5 mm] | Modality: Systemic Immunotherapy & Chemotherapy
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P06-C0115] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IB1 [Invasive Carcinoma >= 5 mm to < 2 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P06-C0116] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IB2 [Invasive Carcinoma >= 2 cm to < 4 cm] | Modality: Systemic Immunotherapy & Chem...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P06-C0117] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IB3 [Invasive Carcinoma >= 4 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P06-C0118] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIA1 [Vaginal Involvement Upper 2/3, < 4 cm] | Modality: Systemic Immunotherapy & Ch...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P06-C0119] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIA2 [Vaginal Involvement Upper 2/3, >= 4 cm] | Modality: Concurrent Chemoradiation ...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P06-C0120] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIB [Parametrial Extension without Pelvic Wall] | Modality: Systemic Immunotherapy &...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 7 OF 12]:

[GYN-CERV_03-P07-C0133] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IIA2 [Vaginal Involvement Upper 2/3, >= 4 cm] | Modality: Concurrent Chemoradiation ...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P07-C0134] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIB [Parametrial Extension without Pelvic Wall] | Modality: Systemic Immunotherapy &...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P07-C0135] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IIIA [Lower Third Vagina Involvement] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P07-C0136] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIIB [Pelvic Sidewall Extension or Hydronephrosis] | Modality: Systemic Immunotherap...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P07-C0137] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIIC1 [Pelvic Lymph Node Metastasis] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P07-C0138] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIIC2 [Para-Aortic Lymph Node Metastasis] | Modality: Systemic Immunotherapy & Chemo...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P07-C0139] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IVA [Mucosal Invasion of Bladder or Rectum] | Modality: Concurrent Chemoradiation [C...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P07-C0140] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IVB [Distant Hematogenous or Peritoneal Metastasis] | Modality: Systemic Immunothera...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P07-C0141] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IA1 [Microscopic Stromal Invasion < 3 mm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P07-C0142] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IA2 [Stromal Invasion 3 mm to < 5 mm] | Modality: Systemic Immunotherapy & Chemotherapy
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 8 OF 12]:

[GYN-CERV_03-P08-C0155] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IA1 [Microscopic Stromal Invasion < 3 mm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P08-C0156] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IA2 [Stromal Invasion 3 mm to < 5 mm] | Modality: Systemic Immunotherapy & Chemotherapy
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P08-C0157] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IB1 [Invasive Carcinoma >= 5 mm to < 2 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P08-C0158] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IB2 [Invasive Carcinoma >= 2 cm to < 4 cm] | Modality: Systemic Immunotherapy & Chem...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P08-C0159] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IB3 [Invasive Carcinoma >= 4 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P08-C0160] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIA1 [Vaginal Involvement Upper 2/3, < 4 cm] | Modality: Systemic Immunotherapy & Ch...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P08-C0161] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIA2 [Vaginal Involvement Upper 2/3, >= 4 cm] | Modality: Concurrent Chemoradiation ...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P08-C0162] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIB [Parametrial Extension without Pelvic Wall] | Modality: Systemic Immunotherapy &...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P08-C0163] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IIIA [Lower Third Vagina Involvement] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P08-C0164] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIIB [Pelvic Sidewall Extension or Hydronephrosis] | Modality: Systemic Immunotherap...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 9 OF 12]:

[GYN-CERV_03-P09-C0177] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IIIA [Lower Third Vagina Involvement] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P09-C0178] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIIB [Pelvic Sidewall Extension or Hydronephrosis] | Modality: Systemic Immunotherap...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P09-C0179] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIIC1 [Pelvic Lymph Node Metastasis] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P09-C0180] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIIC2 [Para-Aortic Lymph Node Metastasis] | Modality: Systemic Immunotherapy & Chemo...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P09-C0181] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IVA [Mucosal Invasion of Bladder or Rectum] | Modality: Concurrent Chemoradiation [C...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P09-C0182] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IVB [Distant Hematogenous or Peritoneal Metastasis] | Modality: Systemic Immunothera...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P09-C0183] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IA1 [Microscopic Stromal Invasion < 3 mm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P09-C0184] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IA2 [Stromal Invasion 3 mm to < 5 mm] | Modality: Systemic Immunotherapy & Chemotherapy
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P09-C0185] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IB1 [Invasive Carcinoma >= 5 mm to < 2 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P09-C0186] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IB2 [Invasive Carcinoma >= 2 cm to < 4 cm] | Modality: Systemic Immunotherapy & Chem...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 10 OF 12]:

[GYN-CERV_03-P10-C0199] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IB1 [Invasive Carcinoma >= 5 mm to < 2 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P10-C0200] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IB2 [Invasive Carcinoma >= 2 cm to < 4 cm] | Modality: Systemic Immunotherapy & Chem...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P10-C0201] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IB3 [Invasive Carcinoma >= 4 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P10-C0202] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIA1 [Vaginal Involvement Upper 2/3, < 4 cm] | Modality: Systemic Immunotherapy & Ch...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P10-C0203] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIA2 [Vaginal Involvement Upper 2/3, >= 4 cm] | Modality: Concurrent Chemoradiation ...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P10-C0204] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIB [Parametrial Extension without Pelvic Wall] | Modality: Systemic Immunotherapy &...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P10-C0205] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IIIA [Lower Third Vagina Involvement] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P10-C0206] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIIB [Pelvic Sidewall Extension or Hydronephrosis] | Modality: Systemic Immunotherap...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P10-C0207] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IIIC1 [Pelvic Lymph Node Metastasis] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P10-C0208] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIIC2 [Para-Aortic Lymph Node Metastasis] | Modality: Systemic Immunotherapy & Chemo...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 11 OF 12]:

[GYN-CERV_03-P11-C0221] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIIC1 [Pelvic Lymph Node Metastasis] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P11-C0222] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIIC2 [Para-Aortic Lymph Node Metastasis] | Modality: Systemic Immunotherapy & Chemo...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P11-C0223] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IVA [Mucosal Invasion of Bladder or Rectum] | Modality: Concurrent Chemoradiation [C...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P11-C0224] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IVB [Distant Hematogenous or Peritoneal Metastasis] | Modality: Systemic Immunothera...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P11-C0225] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IA1 [Microscopic Stromal Invasion < 3 mm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P11-C0226] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IA2 [Stromal Invasion 3 mm to < 5 mm] | Modality: Systemic Immunotherapy & Chemotherapy
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P11-C0227] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IB1 [Invasive Carcinoma >= 5 mm to < 2 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P11-C0228] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IB2 [Invasive Carcinoma >= 2 cm to < 4 cm] | Modality: Systemic Immunotherapy & Chem...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P11-C0229] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IB3 [Invasive Carcinoma >= 4 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P11-C0230] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIA1 [Vaginal Involvement Upper 2/3, < 4 cm] | Modality: Systemic Immunotherapy & Ch...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: FIGO Cancer Report: Staging and Treatment of Cervical Carcinoma [2018/2021 Update]
ISSUING BODY: International Federation of Gynecology and Obstetrics [FIGO]
CANCER DOMAIN: Cervical Cancer | ANATOMICAL SITE: Cervix Uteri & Regional Lymphatics

SECTION PROTOCOLS [PAGE 12 OF 12]:

[GYN-CERV_03-P12-C0243] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IB3 [Invasive Carcinoma >= 4 cm] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P12-C0244] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIA1 [Vaginal Involvement Upper 2/3, < 4 cm] | Modality: Systemic Immunotherapy & Ch...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P12-C0245] Targeted Second-Line Therapy with Antibody-Drug Conjugates: Concurrent Chemora...
Stage: FIGO Stage IIA2 [Vaginal Involvement Upper 2/3, >= 4 cm] | Modality: Concurrent Chemoradiation ...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for pharmacolo...

[GYN-CERV_03-P12-C0246] Patient Education and Symptom Recognition: Systemic Immunotherapy & Chemotherapy
Stage: FIGO Stage IIB [Parametrial Extension without Pelvic Wall] | Modality: Systemic Immunotherapy &...
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for patient ed...

[GYN-CERV_03-P12-C0247] Screening and HPV Cotesting Guidelines: Concurrent Chemoradiation [CCRT]
Stage: FIGO Stage IIIA [Lower Third Vagina Involvement] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for diagnostic...

[GYN-CERV_03-P12-C0248] Surgical Management of Early-Stage Invasive Carcinoma: Systemic Immunotherapy ...
Stage: FIGO Stage IIIB [Pelvic Sidewall Extension or Hydronephrosis] | Modality: Systemic Immunotherap...
Regimen: Pembrolizumab [200 mg IV every 3 weeks [or 400 mg IV every 6 weeks] for PD-L1 CPS >= 1]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for surgical o...

[GYN-CERV_03-P12-C0249] Definitive Chemoradiation in Locally Advanced Disease: Concurrent Chemoradiati...
Stage: FIGO Stage IIIC1 [Pelvic Lymph Node Metastasis] | Modality: Concurrent Chemoradiation [CCRT]
Regimen: Tisotumab Vedotin [2.0 mg/kg IV every 3 weeks for recurrent/metastatic disease post-chemother...
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...

[GYN-CERV_03-P12-C0250] First-Line Systemic Therapy and Immunotherapy Integration: Systemic Immunother...
Stage: FIGO Stage IIIC2 [Para-Aortic Lymph Node Metastasis] | Modality: Systemic Immunotherapy & Chemo...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiation [max 70 mg/dose]]
Summary: International Federation of Gynecology and Obstetrics [FIGO] recommendation for treatment ...