CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 1 OF 16]:

[GYN-OVAR_02-P01-C0001] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Primary Debu...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P01-C0002] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Ma...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P01-C0003] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Primary Debulking ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P01-C0004] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Mainte...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P01-C0005] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Primary Debulkin...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P01-C0006] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Maintenance PARP Inhibitor ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P01-C0007] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Primary Debulkin...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P01-C0008] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Maintenance PARP Inhi...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P01-C0009] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Primary Debulking...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P01-C0010] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Maintenance PARP Inhibi...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 2 OF 16]:

[GYN-OVAR_02-P02-C0023] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Primary Debulking Surge...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P02-C0024] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P02-C0025] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Primary Debulking Surgery ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P02-C0026] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Ma...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P02-C0027] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Primary Debu...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P02-C0028] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Ma...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P02-C0029] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Primary Debulking ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P02-C0030] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Mainte...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P02-C0031] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Primary Debulkin...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P02-C0032] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Maintenance PARP Inhibitor ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 3 OF 16]:

[GYN-OVAR_02-P03-C0045] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Primary Debulking Surgery [...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P03-C0046] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Maintenance PARP...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P03-C0047] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Primary Debulking Sur...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P03-C0048] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Maintenance PARP ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P03-C0049] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Primary Debulking Surge...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P03-C0050] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P03-C0051] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Primary Debulking Surgery ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P03-C0052] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Ma...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P03-C0053] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Primary Debu...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P03-C0054] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Ma...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 4 OF 16]:

[GYN-OVAR_02-P04-C0067] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Pr...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P04-C0068] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Maintenance PARP I...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P04-C0069] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Primar...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P04-C0070] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Maintenance PARP...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P04-C0071] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Primary Debulking Surgery [...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P04-C0072] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Maintenance PARP...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P04-C0073] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Primary Debulking Sur...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P04-C0074] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Maintenance PARP ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P04-C0075] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Primary Debulking Surge...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P04-C0076] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 5 OF 16]:

[GYN-OVAR_02-P05-C0089] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P05-C0090] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Maintenance PARP Inhibitor...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P05-C0091] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Pr...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P05-C0092] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Maintenance ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P05-C0093] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Pr...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P05-C0094] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Maintenance PARP I...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P05-C0095] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Primar...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P05-C0096] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Maintenance PARP...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P05-C0097] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Primary Debulking Surgery [...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P05-C0098] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Maintenance PARP...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 6 OF 16]:

[GYN-OVAR_02-P06-C0111] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Primary Debulkin...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P06-C0112] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Maintenance PARP Inhi...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P06-C0113] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Primary Debulking...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P06-C0114] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Maintenance PARP Inhibi...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P06-C0115] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P06-C0116] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Maintenance PARP Inhibitor...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P06-C0117] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Pr...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P06-C0118] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Maintenance ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P06-C0119] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Pr...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P06-C0120] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Maintenance PARP I...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 7 OF 16]:

[GYN-OVAR_02-P07-C0133] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Primary Debulking ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P07-C0134] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Mainte...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P07-C0135] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Primary Debulkin...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P07-C0136] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Maintenance PARP Inhibitor ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P07-C0137] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Primary Debulkin...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P07-C0138] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Maintenance PARP Inhi...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P07-C0139] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Primary Debulking...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P07-C0140] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Maintenance PARP Inhibi...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P07-C0141] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P07-C0142] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Maintenance PARP Inhibitor...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 8 OF 16]:

[GYN-OVAR_02-P08-C0155] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Primary Debulking Surgery ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P08-C0156] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Ma...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P08-C0157] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Primary Debu...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P08-C0158] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Ma...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P08-C0159] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Primary Debulking ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P08-C0160] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Mainte...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P08-C0161] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Primary Debulkin...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P08-C0162] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Maintenance PARP Inhibitor ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P08-C0163] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Primary Debulkin...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P08-C0164] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Maintenance PARP Inhi...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 9 OF 16]:

[GYN-OVAR_02-P09-C0177] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Primary Debulking Sur...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P09-C0178] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Maintenance PARP ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P09-C0179] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Primary Debulking Surge...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P09-C0180] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P09-C0181] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Primary Debulking Surgery ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P09-C0182] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Ma...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P09-C0183] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Primary Debu...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P09-C0184] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Ma...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P09-C0185] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Primary Debulking ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P09-C0186] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Mainte...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 10 OF 16]:

[GYN-OVAR_02-P10-C0199] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Primar...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P10-C0200] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Maintenance PARP...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P10-C0201] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Primary Debulking Surgery [...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P10-C0202] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Maintenance PARP...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P10-C0203] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Primary Debulking Sur...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P10-C0204] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Maintenance PARP ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P10-C0205] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Primary Debulking Surge...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P10-C0206] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P10-C0207] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Primary Debulking Surgery ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P10-C0208] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Ma...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 11 OF 16]:

[GYN-OVAR_02-P11-C0221] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Pr...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P11-C0222] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Maintenance ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P11-C0223] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Pr...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P11-C0224] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Maintenance PARP I...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P11-C0225] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Primar...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P11-C0226] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Maintenance PARP...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P11-C0227] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Primary Debulking Surgery [...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P11-C0228] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Maintenance PARP...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P11-C0229] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Primary Debulking Sur...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P11-C0230] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Maintenance PARP ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 12 OF 16]:

[GYN-OVAR_02-P12-C0243] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Primary Debulking...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P12-C0244] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Maintenance PARP Inhibi...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P12-C0245] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P12-C0246] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Maintenance PARP Inhibitor...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P12-C0247] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Pr...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P12-C0248] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Maintenance ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P12-C0249] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Pr...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P12-C0250] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Maintenance PARP I...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P12-C0251] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Primar...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P12-C0252] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Maintenance PARP...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 13 OF 16]:

[GYN-OVAR_02-P13-C0265] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Primary Debulkin...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P13-C0266] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Maintenance PARP Inhibitor ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P13-C0267] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Primary Debulkin...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P13-C0268] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Maintenance PARP Inhi...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P13-C0269] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Primary Debulking...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P13-C0270] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Maintenance PARP Inhibi...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P13-C0271] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P13-C0272] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Maintenance PARP Inhibitor...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P13-C0273] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Pr...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P13-C0274] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Maintenance ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 14 OF 16]:

[GYN-OVAR_02-P14-C0287] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Primary Debu...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P14-C0288] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Ma...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P14-C0289] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Primary Debulking ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P14-C0290] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Mainte...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P14-C0291] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Primary Debulkin...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P14-C0292] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Maintenance PARP Inhibitor ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P14-C0293] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Primary Debulkin...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P14-C0294] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Maintenance PARP Inhi...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P14-C0295] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Primary Debulking...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P14-C0296] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Maintenance PARP Inhibi...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 15 OF 16]:

[GYN-OVAR_02-P15-C0309] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIIB [Macroscopic Peritoneal Metastases <= 2 cm] | Modality: Primary Debulking Surge...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P15-C0310] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIIC [Macroscopic Peritoneal Metastases > 2 cm or Capsular Liver/Spleen] | Modality:...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P15-C0311] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IVA [Pleural Effusion with Positive Cytology] | Modality: Primary Debulking Surgery ...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P15-C0312] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IVB [Parenchymal Liver or Spleen Metastases / Extra-Abdominal Organs] | Modality: Ma...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-OVAR_02-P15-C0313] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IA [Tumor Limited to One Ovary, Capsule Intact, No Ascites] | Modality: Primary Debu...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P15-C0314] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IB [Tumor Involves Both Ovaries, Capsule Intact, No Tumor on Surface] | Modality: Ma...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P15-C0315] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IC1 [Surgical Capsule Rupture / Intraoperative Spill] | Modality: Primary Debulking ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P15-C0316] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IC2 [Capsule Ruptured Before Surgery or Tumor on Ovarian Surface] | Modality: Mainte...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...

[GYN-OVAR_02-P15-C0317] Novel Antibody-Drug Conjugates in Platinum-Resistant Relapse: Primary Debulkin...
Stage: FIGO Stage IC3 [Malignant Cells in Ascites or Peritoneal Washings] | Modality: Primary Debulkin...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for targeted therapeutics & re...

[GYN-OVAR_02-P15-C0318] Symptom Triad and Patient Education in Ovarian Cancer: Maintenance PARP Inhibi...
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Maintenance PARP Inhibitor ...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Epithelial Ovarian & Primary Peritoneal Carci...
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Ovarian Cancer | ANATOMICAL SITE: Ovaries & Peritoneal Cavity

SECTION PROTOCOLS [PAGE 16 OF 16]:

[GYN-OVAR_02-P16-C0331] Surgical Cytoreduction and R0 Resection Goals: Primary Debulking Surgery [PDS]
Stage: FIGO Stage IIA [Extension to Uterus or Fallopian Tubes] | Modality: Primary Debulking Surgery [...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for surgical oncology in Ovari...

[GYN-OVAR_02-P16-C0332] First-Line Platinum-Based Chemotherapy and Regimens: Maintenance PARP Inhibito...
Stage: FIGO Stage IIB [Extension to Other Pelvic Intraperitoneal Tissues] | Modality: Maintenance PARP...
Regimen: Niraparib [200 mg [weight < 77 kg or platelets < 150k] or 300 mg orally daily for up to 3 years]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-OVAR_02-P16-C0333] PARP Inhibitor Maintenance in BRCA and HRD-Positive Tumors: Primary Debulking ...
Stage: FIGO Stage IIIA1 [Retroperitoneal Lymph Node Metastasis Only] | Modality: Primary Debulking Sur...
Regimen: Mirvetuximab Soravtansine [6 mg/kg [adjusted ideal body weight] IV q3w for FRalpha-positive p...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for pharmacology & precision t...

[GYN-OVAR_02-P16-C0334] Neoadjuvant Chemotherapy and Interval Debulking Criteria: Maintenance PARP Inh...
Stage: FIGO Stage IIIA2 [Microscopic Extrapelvic Peritoneal Involvement] | Modality: Maintenance PARP ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for combined multimodal oncolo...