CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 1 OF 10]:

[GYN-VAG_01-P01-C0001] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P01-C0002] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P01-C0003] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P01-C0004] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P01-C0005] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P01-C0006] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P01-C0007] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P01-C0008] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P01-C0009] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P01-C0010] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 2 OF 10]:

[GYN-VAG_01-P02-C0023] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P02-C0024] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P02-C0025] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P02-C0026] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P02-C0027] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P02-C0028] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P02-C0029] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P02-C0030] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P02-C0031] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P02-C0032] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 3 OF 10]:

[GYN-VAG_01-P03-C0045] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P03-C0046] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P03-C0047] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P03-C0048] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P03-C0049] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P03-C0050] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P03-C0051] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P03-C0052] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P03-C0053] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P03-C0054] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 4 OF 10]:

[GYN-VAG_01-P04-C0067] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P04-C0068] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P04-C0069] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P04-C0070] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P04-C0071] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P04-C0072] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P04-C0073] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P04-C0074] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P04-C0075] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P04-C0076] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 5 OF 10]:

[GYN-VAG_01-P05-C0089] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P05-C0090] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P05-C0091] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P05-C0092] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P05-C0093] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P05-C0094] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P05-C0095] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P05-C0096] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P05-C0097] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P05-C0098] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 6 OF 10]:

[GYN-VAG_01-P06-C0111] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P06-C0112] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P06-C0113] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P06-C0114] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P06-C0115] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P06-C0116] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P06-C0117] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P06-C0118] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P06-C0119] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P06-C0120] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 7 OF 10]:

[GYN-VAG_01-P07-C0133] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P07-C0134] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P07-C0135] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P07-C0136] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P07-C0137] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P07-C0138] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P07-C0139] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P07-C0140] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P07-C0141] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P07-C0142] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 8 OF 10]:

[GYN-VAG_01-P08-C0155] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P08-C0156] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P08-C0157] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P08-C0158] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P08-C0159] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P08-C0160] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P08-C0161] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P08-C0162] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P08-C0163] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P08-C0164] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 9 OF 10]:

[GYN-VAG_01-P09-C0177] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P09-C0178] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P09-C0179] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P09-C0180] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P09-C0181] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

[GYN-VAG_01-P09-C0182] Definitive Chemoradiation and Interstitial Brachytherapy Delivery: Radical Surg...
Stage: FIGO Stage II [Carcinoma Has Involved Subvaginal Tissue without Pelvic Wall Extension] | Modali...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5 IV + Paclitaxel 175 mg/m2 IV q21d for metastatic ...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [radiation oncol...

[GYN-VAG_01-P09-C0183] Lymphatic Drainage Pathways: Upper vs Lower Vagina: Interstitial Needle Brachyt...
Stage: FIGO Stage III [Carcinoma Has Extended onto the Pelvic Sidewall or Nodal Metastasis] | Modality...
Regimen: Mitomycin C [10 mg/m2 IV day 1 and 29 combined with 5-FU in select squamous protocols]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for anatomical considerations ...

[GYN-VAG_01-P09-C0184] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P09-C0185] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...

[GYN-VAG_01-P09-C0186] Etiology and High-Risk HPV in Vaginal Neoplasia: Definitive External Beam Radio...
Stage: FIGO Stage I [Carcinoma Strictly Confined to the Vaginal Mucosa and Wall] | Modality: Definitiv...
Regimen: Cisplatin [40 mg/m2 IV weekly concurrent with radiotherapy for radiosensitization]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for etiology & pathology in Va...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: NCCN Clinical Practice Guidelines in Oncology: Vaginal Cancer Management Protocols
ISSUING BODY: National Comprehensive Cancer Network [NCCN]
CANCER DOMAIN: Vaginal Cancer | ANATOMICAL SITE: Vagina [Vaginal Canal]

SECTION PROTOCOLS [PAGE 10 OF 10]:

[GYN-VAG_01-P10-C0199] Systemic Protocols in Advanced and Recurrent Disease: Systemic Chemotherapy for...
Stage: FIGO Stage IVA [Tumor Invades Mucosa of Bladder or Rectum or Extends Beyond True Pelvis] | Moda...
Regimen: Fluorouracil [5-FU] [1,000 mg/m2/day continuous IV infusion on days 1-4 and 29-32 with radiat...
Summary: National Comprehensive Cancer Network [NCCN] recommendation for treatment [medical oncolog...

[GYN-VAG_01-P10-C0200] Patient Education and Symptom Presentation: Concurrent Radiosensitizing Chemoth...
Stage: FIGO Stage IVB [Distant Metastasis to Lungs, Bone, Liver, or Supraclavicular Nodes] | Modality:...
Regimen: Pembrolizumab [200 mg IV q3w for recurrent/metastatic PD-L1 positive or MSI-H tumors]
Summary: National Comprehensive Cancer Network [NCCN] recommendation for patient education & sympto...