
CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 1 OF 10]:

[GYN-FALL_04-P01-C0001] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Comprehensive Surgi...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P01-C0002] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P01-C0003] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Cytoreducti...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P01-C0004] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P01-C0005] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: First-Line...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P01-C0006] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P01-C0007] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P01-C0008] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Cytoreductive Surgery
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P01-C0009] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P01-C0010] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: First-Line ...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 2 OF 10]:

[GYN-FALL_04-P02-C0023] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P02-C0024] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Risk-Reduci...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P02-C0025] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P02-C0026] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: Comprehens...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P02-C0027] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P02-C0028] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P02-C0029] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Risk-Reducing Salpi...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P02-C0030] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P02-C0031] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Comprehensi...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P02-C0032] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 3 OF 10]:

[GYN-FALL_04-P03-C0045] Clinical Presentation and Latzko's Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: First-Line ...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P03-C0046] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P03-C0047] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: Targeted P...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P03-C0048] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P03-C0049] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P03-C0050] Clinical Presentation and Latzko's Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: First-Line Platinum...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P03-C0051] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P03-C0052] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Targeted PA...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P03-C0053] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P03-C0054] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: Risk-Reduc...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 4 OF 10]:

[GYN-FALL_04-P04-C0067] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P04-C0068] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: Cytoreduct...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P04-C0069] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P04-C0070] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P04-C0071] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Comprehensive Surgi...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P04-C0072] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P04-C0073] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Cytoreducti...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P04-C0074] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P04-C0075] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: First-Line...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P04-C0076] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 5 OF 10]:

[GYN-FALL_04-P05-C0089] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: Risk-Reduc...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P05-C0090] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P05-C0091] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P05-C0092] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Targeted PARP Inhib...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P05-C0093] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P05-C0094] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Risk-Reduci...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P05-C0095] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P05-C0096] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: Comprehens...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P05-C0097] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P05-C0098] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 6 OF 10]:

[GYN-FALL_04-P06-C0111] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P06-C0112] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P06-C0113] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Cytoreductive Surgery
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P06-C0114] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P06-C0115] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: First-Line ...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P06-C0116] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P06-C0117] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: Targeted P...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P06-C0118] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P06-C0119] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P06-C0120] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: First-Line Platinum...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 7 OF 10]:

[GYN-FALL_04-P07-C0133] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P07-C0134] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Risk-Reducing Salpi...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P07-C0135] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P07-C0136] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Comprehensi...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P07-C0137] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P07-C0138] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: Cytoreduct...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P07-C0139] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P07-C0140] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P07-C0141] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Comprehensive Surgi...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P07-C0142] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 8 OF 10]:

[GYN-FALL_04-P08-C0155] Clinical Presentation and Latzko's Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: First-Line Platinum...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P08-C0156] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P08-C0157] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Targeted PA...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P08-C0158] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P08-C0159] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: Risk-Reduc...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P08-C0160] Clinical Presentation and Latzko's Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P08-C0161] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P08-C0162] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Targeted PARP Inhib...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P08-C0163] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P08-C0164] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Risk-Reduci...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 9 OF 10]:

[GYN-FALL_04-P09-C0177] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P09-C0178] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Cytoreducti...
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P09-C0179] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P09-C0180] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage II [Extension to Uterus, Ovaries, or Other Pelvic Structures] | Modality: First-Line...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P09-C0181] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage III [Microscopic or Macroscopic Peritoneal Implants Outside Pelvis / Nodal Spread] |...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

[GYN-FALL_04-P09-C0182] Surgical Management and Complete Macroscopic Resection: Targeted PARP Inhibito...
Stage: FIGO Stage IV [Distant Parenchymal Metastases or Pleural Effusion with Malignant Cytology] | Mo...
Regimen: Niraparib [200-300 mg orally daily maintenance in platinum-sensitive primary or recurrent tub...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for surgical onc...

[GYN-FALL_04-P09-C0183] Systemic Platinum Chemotherapy and PARP Inhibition: Cytoreductive Surgery
Stage: FIGO Stage I [Carcinoma Limited to One or Both Fallopian Tubes] | Modality: Cytoreductive Surgery
Regimen: Gemcitabine + Carboplatin [Gemcitabine 1,000 mg/m2 days 1,8 + Carboplatin AUC 4 day 1 q21d fo...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for treatment [m...

[GYN-FALL_04-P09-C0184] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IA [Limited to One Tube, No Penetration of Serosal Surface, No Ascites] | Modality: ...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P09-C0185] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: First-Line ...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...

[GYN-FALL_04-P09-C0186] The Fallopian Tube as the Origin of High-Grade Serous Carcinoma: Comprehensive...
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Carboplatin + Paclitaxel [Carboplatin AUC 5-6 IV + Paclitaxel 175 mg/m2 IV over 3 hours every...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for etiology & c...

CLINICAL PRACTICE GUIDELINE & REFERENCE MANUAL
DOCUMENT: ACOG Practice Bulletin No. 182: Hereditary Breast and Ovarian Cancer Syndrome & Opportunisti...
ISSUING BODY: American College of Obstetricians and Gynecologists [ACOG]
CANCER DOMAIN: Fallopian Tube Cancer | ANATOMICAL SITE: Fallopian Tubes & Ovaries [BRCA1/2 Carriers]

SECTION PROTOCOLS [PAGE 10 OF 10]:

[GYN-FALL_04-P10-C0199] Risk-Reducing Salpingo-Oophorectomy and the SEE-FIM Protocol: Risk-Reducing Sa...
Stage: FIGO Stage IB [Involves Both Tubes, No Serosal Penetration, No Ascites] | Modality: Risk-Reduci...
Regimen: Olaparib [300 mg orally twice daily maintenance for up to 2 years in BRCA-mutated tubal cancer]
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for preventive s...

[GYN-FALL_04-P10-C0200] Clinical Presentation and Latzko’s Triad: First-Line Platinum-Based Chemotherapy
Stage: FIGO Stage IC [Serosal Penetration, Capsule Rupture, or Positive Peritoneal Washings] | Modalit...
Regimen: Bevacizumab [15 mg/kg IV every 3 weeks combined with carboplatin/paclitaxel and continued as ...
Summary: American College of Obstetricians and Gynecologists [ACOG] recommendation for patient educ...