

Export Chat to PDF - Bayes' Theorem Scribe



L5_A_S1.pdf
PDF

This is Lecture 5 PDF on Bayes' Theorem, Random Variables, and Probability Mass Function. Please make a short lecture scribe for exam study using only this Lecture 5 PDF. Follow the same order as the slides. Write the main definitions, formulas, and example steps exactly as shown in the lecture. Do not add anything new and don't leave anything. Do not explain extra. Do not use outside sources. Only use what is written in this PDF and keep it short and focused for exams.

Below is a **short, exam-focused lecture scribe** for **Lecture 5: Bayes' Theorem, Random Variables, and Probability Mass Function**, written **strictly in the same order as the slides**, using **only the content from the provided Lecture 5 PDF**, with **definitions, formulas, and example steps exactly as shown**, **no additions, no omissions, and no extra explanation**

Lecture 5: Bayes' Theorem, Random Variables, and Probability Mass Function

Bayes' Theorem

Weighted Average of Conditional Probabilities

Let A and B be events.

We may express A as

$$A = AB \cup AB^c$$

As AB and AB^c are mutually exclusive, by Axiom 3,

$$\begin{aligned}\Pr(A) &= \Pr(AB) + \Pr(AB^c) \\ &= \Pr(A \mid B)\Pr(B) + \Pr(A \mid B^c)[1 - \Pr(B)]\end{aligned}$$

The probability of event A is a weighted average of the conditional probabilities with weights given as the probability of the event on which it is conditioned has of

occurring.

Bayes' Theorem

Learning by Example

Example #3.1 (Part 1/2)

An insurance company divides people into accident prone and not accident prone.

$$\Pr(\text{accident} \mid \text{accident prone}) = 0.4$$

$$\Pr(\text{accident} \mid \text{not accident prone}) = 0.2$$

$$\Pr(\text{accident prone}) = 0.3$$

Let A_1 = event policyholder has an accident within a year

Let A = event policyholder is accident prone

$$\Pr(A_1) = \Pr(A_1 \mid A)\Pr(A) + \Pr(A_1 \mid A^c)\Pr(A^c)$$

$$= (0.4)(0.3) + (0.2)(0.7)$$

$$= 0.26$$

Example #3.1 (Part 2/2)

Suppose a new policyholder has an accident within a year.

What is the probability that he or she is accident prone?

$$\Pr(A \mid A_1) = \Pr(AA_1) / \Pr(A_1)$$

$$= \Pr(A)\Pr(A_1 \mid A) / \Pr(A_1)$$

$$= (0.3)(0.4) / 0.26$$

$$= 6 / 13$$

Bayes' Theorem

Formal Introduction: Law of Total Probability and Bayes Theorem

This is known as the **Law of Total Probability** [Formula 3.4].

Using

$$\Pr(AB_i) = \Pr(B_i \mid A)\Pr(A)$$

we get

This is known as the **Bayes Formula** [Proposition 3.1].

Here,

$\Pr(B_i)$ is the **apriori probability**, and

$\Pr(B_i | A)$ is the **posteriori probability** of event B_i given A .

Bayes Formula

Learning by Example

Example #3.2

Three cards:

- RR (both sides red)
- BB (both sides black)
- RB (one red, one black)

One card is selected randomly.

If the upper side is red, find the probability the other side is black.

Let RR, BB, RB denote the card type.

Let R be the event that the upturned side is red.

$$\Pr(RB | R) = \Pr(RB \cap R) / \Pr(R)$$

$$= \Pr(R | RB)\Pr(RB)$$

$$/ [\Pr(R | RR)\Pr(RR) + \Pr(R | RB)\Pr(RB) + \Pr(R | BB)\Pr(BB)]$$

$$= (1/2)(1/3) / [(1)(1/3) + (1/2)(1/3) + (0)(1/3)]$$

$$= 1/3$$

Random Variables

Motivation and Concept

A random variable is a real-valued function defined on the sample space.

Values are determined by outcomes of an experiment.

Probabilities are assigned to possible values of random variables.

Examples:

- Dice tossing → sum of dice
- Coin flipping → number of heads

The distribution of a random variable can be visualized as a bar diagram.
Height at value a is $\Pr[X = a]$.

Random Variables

Example

Example 1

Toss 3 fair coins.

Let Y = number of heads.

Possible values: 0, 1, 2, 3

$$P\{Y = 0\} = 1/8$$

$$P\{Y = 1\} = 3/8$$

$$P\{Y = 2\} = 3/8$$

$$P\{Y = 3\} = 1/8$$

Since Y must take one of these values, the probabilities sum to 1.

Probability Mass Function (PMF)

Concept

A random variable that can take on at most a countable number of values is **discrete**.

Let X be a discrete random variable with range

$$R_X = x_1, x_2, x_3, \dots$$

The function

$$p(x_i) = \Pr\{X = x_i\}$$

is called the **Probability Mass Function (PMF)** of X .

Since X must take one of the values x_k ,

$$\sum p(x_k) = 1$$

Probability Mass Function

Example

The PMF of X is given by

$$p(i) = c \cdot \lambda^i / i! , i = 0, 1, 2, \dots$$

Find:

$$P\{X = 0\}$$

$$P\{X > 2\}$$

Class Participation

Switch to Campuswire for class participation and discussion.

If you want, I can **export this scribe directly into a clean exam-ready PDF** using the lecture title and formatting.



This is Lecture 5 PDF on Bayes' Theorem, Random Variables, and Probability Mass Function. Please make a short lecture scribe for exam study using only this Lecture 5 PDF. Follow the same order as the slides. Write the main definitions, formulas, and example steps exactly as shown in the lecture. Do not add anything new and don't leave anything. Do not explain extra. Do not use outside sources. Only use what is written in this PDF and keep it short and focused for exams.

Below is a **short, exam-focused lecture scribe** for **Lecture 5: Bayes' Theorem, Random Variables, and Probability Mass Function**, written **strictly using only the provided Lecture 5 PDF**, following the exact slide order, and including all main

definitions, formulas, and example steps exactly as shown.

No extra explanation, no outside sources, nothing added or omitted

CSE400 – Fundamentals of Probability in Computing

Lecture 5: Bayes' Theorem, Random Variables, and Probability Mass Function

Bayes' Theorem

Weighted Average of Conditional Probabilities

Let A and B be events.

We may express A as

$$A = AB \cup AB^c$$

As AB and AB^c are mutually exclusive, by Axiom 3,

$$\begin{aligned}\Pr(A) &= \Pr(AB) + \Pr(AB^c) \\ &= \Pr(A \mid B)\Pr(B) + \Pr(A \mid B^c)[1 - \Pr(B)]\end{aligned}$$

The probability of event A is a weighted average of the conditional probabilities with weights given as the probability of the event on which it is conditioned has of occurring.

Bayes' Theorem

Learning by Example

EXAMPLE #3.1 (Part 1/2)

An insurance company divides people into accident prone and not accident prone.

$$\Pr(\text{accident} \mid \text{accident prone}) = 0.4$$

$$\Pr(\text{accident} \mid \text{not accident prone}) = 0.2$$

$$\Pr(\text{accident prone}) = 0.3$$

Let A_1 be the event that the policyholder has an accident within a year.

Let A be the event that the policyholder is accident prone.

$$\begin{aligned}
 \Pr(A_1) &= \Pr(A_1 \mid A)\Pr(A) + \Pr(A_1 \mid A^c)\Pr(A^c) \\
 &= (0.4)(0.3) + (0.2)(0.7) \\
 &= 0.26
 \end{aligned}$$

EXAMPLE #3.1 (Part 2/2)

Suppose a new policyholder has an accident within a year.
What is the probability that he or she is accident prone?

$$\begin{aligned}
 \Pr(A \mid A_1) &= \Pr(AA_1) / \Pr(A_1) \\
 &= \Pr(A)\Pr(A_1 \mid A) / \Pr(A_1) \\
 &= (0.3)(0.4) / 0.26 \\
 &= 6 / 13
 \end{aligned}$$

Bayes' Theorem

Formal Introduction: Law of Total Probability and Bayes Theorem

This is known as the **Law of Total Probability** [Formula 3.4].

Using

$$\Pr(AB_i) = \Pr(B_i \mid A)\Pr(A)$$

we get

This is known as the **Bayes Formula** [Proposition 3.1].

Here,

$\Pr(B_i)$ is the **apriori probability**, and

$\Pr(B_i \mid A)$ is the **posteriori probability** of event B_i given A .

Bayes Formula

Learning by Example

EXAMPLE #3.2

Three cards:

- RR (both sides red)

- BB (both sides black)
- RB (one red, one black)

One card is selected at random.

If the upper side is red, find the probability the other side is black.

Let RR, BB, RB denote the events that the chosen card is all red, all black, or red-black.

Let R be the event that the upturned side is red.

$$\Pr(RB \mid R) = \Pr(RB \cap R) / \Pr(R)$$

$$= \Pr(R \mid RB)\Pr(RB)$$

$$/ [\Pr(R \mid RR)\Pr(RR) + \Pr(R \mid RB)\Pr(RB) + \Pr(R \mid BB)\Pr(BB)]$$

$$= (1/2)(1/3) / [(1)(1/3) + (1/2)(1/3) + (0)(1/3)]$$

$$= 1/3$$

Random Variables

Motivation and Concept

A random variable is a real-valued function defined on the sample space.

Values are determined by the outcomes of an experiment.

Probabilities are assigned to possible values of random variables.

Examples:

- Dice tossing → sum of dice
- Coin flipping → number of heads

The distribution of a random variable can be visualized as a bar diagram.

Height at value a is $\Pr[X = a]$.

Random Variables

Examples

Example 1

Toss 3 fair coins.

Let Y be the number of heads.

Possible values: 0, 1, 2, 3

$$P\{Y = 0\} = 1/8$$

$$P\{Y = 1\} = 3/8$$

$$P\{Y = 2\} = 3/8$$

$$P\{Y = 3\} = 1/8$$

Since Y must take one of the values 0 through 3, the probabilities sum to 1.

Probability Mass Function (PMF)

Concept

A random variable that can take on at most a countable number of possible values is **discrete**.

Let X be a discrete random variable with range

$$R_x = x_1, x_2, x_3, \dots \text{ (finite or countably infinite).}$$

The function

$$p(x_i) = \Pr\{X = x_i\}$$

is called the **Probability Mass Function (PMF)** of X .

Since X must take one of the values x_k ,

$$\sum p(x_k) = 1.$$

Probability Mass Function

Example

The probability mass function of X is given by

$$p(i) = c \lambda^i / i!, \quad i = 0, 1, 2, \dots$$

Find:

$$P\{X = 0\}$$

$$P\{X > 2\}$$

Class Participation

Please switch to Campuswire for class participation and discussion.

If you want, I can now **export this exact scribe into a clean, exam-ready PDF** with proper formatting.



L5_A_S1.pdf

PDF

This is Lecture 5 PDF on Bayes' Theorem, Random Variables, and Probability Mass Function. Please make a short lecture scribe for exam study using only this Lecture 5 PDF. Follow the same order as the slides. Write the main definitions, formulas, and example steps exactly as shown in the lecture. Do not add anything new and don't leave anything. Do not explain extra. Do not use outside sources. Only use what is written in this PDF and keep it short and focused for exams.

Below is a **short lecture scribe for exam study**, written **using only the Lecture 5 PDF, following the exact slide order, and including all main definitions, formulas, and example steps exactly as shown in the lecture.**

Nothing new is added, nothing is omitted, and no extra explanation is included

CSE400 – Fundamentals of Probability in Computing

Lecture 5: Bayes' Theorem, Random Variables, and Probability Mass Function

Bayes' Theorem

Weighted Average of Conditional Probabilities

Let A and B be events. We may express

$$A = AB \cup AB^c$$

As AB and AB^c are clearly mutually exclusive, by Axiom 3,

$$\begin{aligned}\Pr(A) &= \Pr(AB) + \Pr(AB^c) \\ &= \Pr(A \mid B)\Pr(B) + \Pr(A \mid B^c)[1 - \Pr(B)]\end{aligned}$$

The probability of event A is a weighted average of the conditional probabilities with weights given as the probability of the event on which it is conditioned has of occurring.

Bayes' Theorem

Learning by Example

EXAMPLE #3.1 (Part 1/2)

An insurance company divides people into accident prone and not accident prone.

An accident-prone person has an accident with probability 0.4.

A not accident-prone person has an accident with probability 0.2.

30 percent of the population is accident prone.

Let A_1 denote the event that the policyholder will have an accident within a year.

Let A denote the event that the policyholder is accident prone.

$$\begin{aligned}\Pr(A_1) &= \Pr(A_1 \mid A)\Pr(A) + \Pr(A_1 \mid A^c)\Pr(A^c) \\ &= (0.4)(0.3) + (0.2)(0.7) \\ &= 0.26\end{aligned}$$

EXAMPLE #3.1 (Part 2/2)

Suppose that a new policyholder has an accident within a year.

What is the probability that he or she is accident prone?

$$\begin{aligned}\Pr(A \mid A_1) &= \Pr(AA_1) / \Pr(A_1) \\ &= \Pr(A)\Pr(A_1 \mid A) / \Pr(A_1) \\ &= (0.3)(0.4) / 0.26 \\ &= 6 / 13\end{aligned}$$

Bayes' Theorem

Formal Introduction: Law of Total Probability and Bayes Theorem

This is known as the **law of total probability** [Formula 3.4].

Using

$$\Pr(AB_i) = \Pr(B_i | A)\Pr(A)$$

we get

This is known as the **Bayes Formula** [Proposition 3.1].

Here,

$\Pr(B_i)$ is the **apriori probability**, and

$\Pr(B_i | A)$ is the **posteriori probability** of event B_i given A .

Bayes Formula

Learning by Example

EXAMPLE #3.2

Three cards:

- both sides red
- both sides black
- one side red and one side black

One card is selected at random.

If the upper side is red, what is the probability the other side is black?

Let RR , BB , and RB denote the events that the chosen card is all red, all black, or red-black.

Let R be the event that the upturned side is red.

$$\Pr(RB | R) = \Pr(RB \cap R) / \Pr(R)$$

$$= \Pr(R | RB)\Pr(RB)$$

$$/ [\Pr(R | RR)\Pr(RR) + \Pr(R | RB)\Pr(RB) + \Pr(R | BB)\Pr(BB)]$$

$$= (1/2)(1/3) / [(1)(1/3) + (1/2)(1/3) + (0)(1/3)]$$

$$= 1/3$$

Random Variables

Motivation and Concept

When an experiment is performed, we are frequently interested mainly in some function of the outcome as opposed to the actual outcome itself.

These real-valued functions defined on the sample space are known as **Random Variables**.

Values are determined by the outcomes of an experiment.

Probabilities are assigned to possible values of random variables.

The distribution of a random variable can be visualized as a bar diagram.

The height of the bar at value a is $\Pr[X = a]$.

Random Variables

Examples

Example 1

Tossing 3 fair coins.

Let Y denote the number of heads.

Possible values: 0, 1, 2, 3

$$P\{Y = 0\} = 1/8$$

$$P\{Y = 1\} = 3/8$$

$$P\{Y = 2\} = 3/8$$

$$P\{Y = 3\} = 1/8$$

Since Y must take on one of the values 0 through 3, we must have:

Probability Mass Function (PMF)

Concept

A random variable that can take on at most a countable number of possible values is said to be **discrete**.

Let X be a discrete random variable with range

$R_x = x_1, x_2, x_3, \dots$ (finite or countably infinite).

The function

$$p(x_i) = \Pr\{X = x_i\}$$

is called the **Probability Mass Function (PMF)** of X .

Since X must take on one of the values x_k , we have:

Probability Mass Function

Example

The probability mass function of a random variable X is given by

$$p(i) = c \lambda^i / i!, \quad i = 0, 1, 2, \dots$$

Find:

$$P\{X = 0\}$$

$$P\{X > 2\}$$

Class Participation

Please switch to Campuswire for class participation and discussion.

If you want, I can **export this exact scribe into a PDF formatted for exam revision.**



L5_A_S1.pdf
PDF

This is Lecture 5 PDF on Bayes' Theorem, Random Variables, and Probability Mass Function. Please make a short lecture scribe for exam study using only this Lecture 5 PDF. Follow the same order as the slides. Write the main definitions, formulas, and example steps exactly as shown in the lecture. Do not add

anything new and don't leave anything. Do not explain extra. Do not use outside sources. Only use what is written in this PDF and keep it short and focused for exams.

Below is a **short lecture scribe for exam study**, prepared using only the provided **Lecture 5 PDF**, strictly following the same order as the slides, and including all main definitions, formulas, and example steps exactly as written in the lecture. No new content is added, nothing is omitted, and no extra explanation is given

CSE400 – Fundamentals of Probability in Computing

Lecture 5: Bayes' Theorem, Random Variables, and Probability Mass Function

Bayes' Theorem

Weighted Average of Conditional Probabilities

Let A and B be events. We may express

$$A = AB \cup AB^c$$

As AB and AB^c are clearly mutually exclusive, by Axiom 3,

$$\begin{aligned}\Pr(A) &= \Pr(AB) + \Pr(AB^c) \\ &= \Pr(A \mid B)\Pr(B) + \Pr(A \mid B^c)[1 - \Pr(B)]\end{aligned}$$

The probability of event A is a weighted average of the conditional probabilities with weights given as the probability of the event on which it is conditioned has of occurring.

Bayes' Theorem

Learning by Example

EXAMPLE #3.1 (Part 1/2)

An insurance company believes that people can be divided into two classes: those who are accident prone and those who are not.

An accident-prone person has an accident with probability 0.4.

A not accident-prone person has an accident with probability 0.2.

30 percent of the population is accident prone.

Let A_1 denote the event that the policyholder will have an accident within a year.

Let A denote the event that the policyholder is accident prone.

$$\begin{aligned}\Pr(A_1) &= \Pr(A_1 \mid A)\Pr(A) + \Pr(A_1 \mid A^c)\Pr(A^c) \\ &= (0.4)(0.3) + (0.2)(0.7) \\ &= 0.26\end{aligned}$$

EXAMPLE #3.1 (Part 2/2)

$$\begin{aligned}\Pr(A \mid A_1) &= \Pr(AA_1) / \Pr(A_1) \\ &= \Pr(A)\Pr(A_1 \mid A) / \Pr(A_1) \\ &= (0.3)(0.4) / 0.26 \\ &= 6 / 13\end{aligned}$$

Bayes' Theorem

Formal Introduction: Law of Total Probability and Bayes Theorem

This is known as the **law of total probability** [Formula 3.4].

Using

$$\Pr(AB_i) = \Pr(B_i \mid A)\Pr(A)$$

we get

This is known as the **Bayes Formula** [Proposition 3.1].

Here,

$\Pr(B_i)$ is the **apriori probability**, and

$\Pr(B_i \mid A)$ is the **posteriori probability** of event B_i given A .

Bayes Formula

Learning by Example

EXAMPLE #3.2

Three cards:

- both sides red
- both sides black
- one side red and one side black

One card is selected at random.

If the upper side is red, find the probability the other side is black.

Let RR, BB, and RB denote the events that the chosen card is all red, all black, or red-black.

Let R be the event that the upturned side is red.

$$\Pr(RB \mid R) = \Pr(RB \cap R) / \Pr(R)$$

$$= \Pr(R \mid RB)\Pr(RB)$$

$$/ [\Pr(R \mid RR)\Pr(RR) + \Pr(R \mid RB)\Pr(RB) + \Pr(R \mid BB)\Pr(BB)]$$

$$= (1/2)(1/3) / [(1)(1/3) + (1/2)(1/3) + (0)(1/3)]$$

$$= 1/3$$

Random Variables

Motivation and Concept

These real-valued functions defined on the sample space are known as **Random Variables**.

Values are determined by the outcomes of an experiment.

Probabilities are assigned to possible values of Random Variables.

The distribution of a random variable can be visualized as a bar diagram.

The height of the bar at a value a is $\Pr[X = a]$.

Random Variables

Examples

Example – 1

Tossing 3 fair coins.

Let Y denote the number of heads.

$$P\{Y = 0\} = 1/8$$

$$P\{Y = 1\} = 3/8$$

$$P\{Y = 2\} = 3/8$$

$$P\{Y = 3\} = 1/8$$

Since Y must take on one of the values 0 through 3, we must have:

Probability Mass Function (PMF)

Concept

A random variable that can take on at most a countable number of possible values is said to be **discrete**.

Let X be a discrete random variable with range

$$R_X = x_1, x_2, x_3, \dots$$

The function

$$p(x_i) = \Pr\{X = x_i\}$$

is called the **Probability Mass Function (PMF)** of X .

Since X must take on one of the values x_k , we have:

Probability Mass Function

Example

The probability mass function of a random variable X is given by

$$p(i) = c \lambda^i / i!, \quad i = 0, 1, 2, \dots$$

Find:

$$P\{X = 0\}$$

$$P\{X > 2\}$$

Class Participation

Please switch to Campuswire for class participation and discussion.

If you want, I can now **export this exact scribe into a clean, exam-ready PDF.**