Deal or No Deal: A Stratified Survey of Simulated Winnings

STAT4102 Applied Sampling Mini-Project

Priyan Sathabridge (2457076)

October 2026

Abstract

This study evaluates a stratified probability survey of a fixed population of 10,000 simulated *Deal or No Deal* contestants. The primary objective is to estimate mean winnings in US dollars with a 95% confidence-interval half-width no greater than 5% of its baseline value. Region-based Neyman allocation is compared with simple random sampling without replacement (SRSWOR). Allowing for 90% response, 2,670 contestants were selected and 2,391 responded. The weighted mean was \$67,453 (95% confidence interval \$64,111–\$70,795), compared with the true mean of \$67,430; its interval met the precision target. Across 1,000 repetitions per design, stratification reduced the mean's empirical variance by 32%, with coverage compatible with 95%. However, it increased the variance of the secondary proportion below opening expected value by 67%. The results demonstrate the value of stratification and weighting for the primary outcome, while showing that allocation gains are outcome-specific and that planned precision does not guarantee the precision of every realised sample.

1. Introduction and literature review

Deal or No Deal presents contestants with a sequence of choices between a guaranteed banker offer and an uncertain prize. Although the expected value of a known prize board is readily calculated, actual winnings also depend on offers and contestants' decisions. Mean winnings therefore provide a useful summary for a broadcaster's prize-budget planning, while the proportion below the opening expected value describes how frequently payouts fall below that benchmark.

This study designs and evaluates a probability survey of **10,000 simulated contestants** from simulated UK, US and Australian versions of the game. It asks: what are mean winnings and the proportion of contestants below opening expected value; does region stratification improve precision relative to simple random sampling without replacement (SRSWOR); and do the estimators show small bias and approximately 95% interval coverage over repeated sampling? All conclusions concern the constructed finite population rather than actual television contestants.

The primary parameter is

$$\bar{Y} = N^{-1} \sum_{i=1}^{N} y_i, \qquad N = 10{,}000,$$

where y_i is contestant i's winnings in US dollars (USD). The secondary parameter is $P = N^{-1} \sum_i I(y_i < EV_{h(i)})$, where $I(\cdot)$ is the indicator function, $h(i)$ is contestant i's region and $EV_{h(i)}$ is the opening expected value (EV) of that region's board.

Post et al. (2008) study *Deal or No Deal* as a setting for risky choice with substantial stakes. Their work motivates the behavioural setting, but the offer curve and decision parameters used here are

assumptions rather than fitted estimates from that study. The present question concerns population-level payouts and survey precision, rather than the estimation of individual preferences.

Neyman (1934) establishes the allocation principle underlying the primary design: with equal unit costs, samples are allocated in proportion to stratum size and within-stratum standard deviation. Cochran (1977) and Lohr (2010) explain the corresponding finite-population variance and weighting framework. This suggests that region may be useful because prize scales differ, but an allocation optimised for winnings need not also be efficient for a binary outcome.

Repeated sampling provides a way to assess those theoretical expectations. Following Morris, White and Crowther (2019), this study compares bias, empirical and estimated standard errors, interval coverage and Monte Carlo uncertainty. Together, the realised survey and simulation assess both the immediate precision gain and its consistency across samples.

2. Methodology

2.1 Population and sampling design

The target population contains 10,000 simulated contestants: 4,000 UK, 4,000 US and 2,000 Australian. The sampling frame is a register with one record per person, including a unique identifier, region, age, gender and opening board value. It covers the whole target population, so target and sampled populations coincide. Each contestant is both the sampling unit and the observation unit reporting winnings and game details. In practice, incomplete registers could exclude eligible contestants.

Each game randomly assigns the regional board's prizes to boxes, keeps one box for the contestant, and opens the others in random order. UK games use 22 prizes (British pounds, GBP 0.01–250,000), opening 5 boxes and then 3 in each of five rounds. US and Australian games use 26 prizes and opening counts 6, 5, 4, 3, 2, 1, 1, 1, 1. The US maximum is USD 1 million; the Australian maximum is 2 million Australian dollars (AUD). The Australian board and its schedule are modelling assumptions, not a verified reconstruction of a particular series.

After round t of a game with T rounds, the offer is f_t times the mean unopened prize, where

$$f_t = \min\{[0.15 + 0.85((t-1)/(T-1))^{1.5}]e^{\epsilon_t}, 1.10\}, \quad \epsilon_t \sim N(0, 0.10^2)$$

is independent random variation between games. This form reproduces the qualitative pattern reported by Post et al. (2008): the banker opens with a small fraction of the expected value, which keeps contestants playing, and raises the fraction towards the expected value in the final rounds. The value 0.15 sets a low opening offer; the base fraction reaches exactly 1 at $t = T$; the power 1.5 makes the curve convex, so offers stay low for several rounds and climb steeply near the end; the noise term lets offers for the same position differ between games by roughly $\pm 20\%$; and the cap of 1.10 allows occasional offers slightly above the expected value. These constants are stylised rather than estimated. They shape the simulated population only: once the population is fixed, its true mean and variances are known, so the sampling results remain valid whatever constants generated it.

Each contestant then chooses between the offer and playing on for the unopened prizes. The certainty equivalent (CE) of that gamble is the guaranteed amount the contestant values exactly as much as the gamble itself. A risk-neutral contestant's CE equals the mean unopened prize; a risk-averse contestant's CE is lower, so they may accept an offer below the expected value. Preferences follow constant relative risk aversion (CRRA) utility, $u(c) = c^{1-\rho}/(1-\rho)$, or $\log c$ when $\rho = 1$, applied to wealth plus prize. This standard model of risky choice summarises risk attitude in one parameter ρ: $\rho = 0$ is risk neutral and larger values are more risk averse. Treating each unopened prize as equally

likely, the CE solves $u(w + CE) = m^{-1} \sum_j u(w + x_j)$, giving

$$CE = \left[m^{-1} \sum_{j=1}^{m} (w + x_j)^{1-\rho} \right]^{1/(1-\rho)} - w,$$

where $x_1, \ldots, x_m$ are the m unopened prizes and w is the contestant's wealth. At $\rho = 1$, the certainty equivalent is $\exp\{m^{-1} \sum_j \log(w + x_j)\} - w$. A contestant accepts the first offer that is at least equal to their CE; if every offer is refused, they receive the prize in their own box. This myopic rule ignores future offers.

Age is a rounded $N(40, 12^2)$ draw **clipped** to 18–75; gender is assigned Female/Male with equal probability. Risk aversion is $\max\{0, N(0.4 + 0.01(\text{age} - 40), 0.3^2)\}$. Wealth is lognormal with median USD 20,000 and log-SD 0.8, converted to local currency for decisions. Fixed modelling factors of 1.27 USD/GBP and 0.66 USD/AUD convert winnings back to USD. They are not dated exchange-rate observations. The population was generated in R with seed 20261019.

Table 1: Population structure. Mean and standard deviation (SD) are in USD; below EV is the percentage below the opening expected value.

Region	N	Share	Mean	SD	Below EV
AU	2,000	0.2	71,223	128,845	80.2
UK	4,000	0.4	28,297	40,194	72.4
US	4,000	0.4	104,666	147,045	74.9

The true values are $\bar{Y} = \$67{,}429.76$ and $P = 0.7498$. Winnings are right-skewed: the median is \$26,624 and the standard deviation (SD) is \$117,400. This arises from many modest payouts and relatively few large prizes.

Region is a natural stratifier because it fixes the prize board. Stratum means range from \$28,297 (UK) to \$104,666 (US), so a large part of the variation in winnings lies between regions; stratified sampling removes this between-stratum component from the variance of the estimator. The strata are therefore externally heterogeneous, although winnings remain highly variable within each region, and stratum SDs differ substantially. Neyman allocation concentrates sampling where variability is greatest, improving precision under equal unit costs. Regional registers also simplify selection and contact. Proportional allocation requires 2,990 respondents for the primary target, compared with 2,401 under continuous Neyman allocation. Cluster sampling could inflate variance if contestants in the same recording series have similar outcomes; no cost evidence justifies that trade-off here.

2.2 Sample size and precision

The target is a **95% confidence-interval half-width** of no more than 5% of the baseline mean: $e = 0.05\bar{Y} = \$3{,}371.49$. Let $z = 1.959964$, the 97.5th percentile of the standard normal distribution, and $V = (e/z)^2$, the largest acceptable variance of the estimator. Baseline means and stratum variances are treated as known prior information, as permitted by the brief. This is a planning assumption, not information obtained from the realised sample.

For SRSWOR, using the finite-population variance S^2 with denominator $N - 1$,

$$n_0 = (zS/e)^2, \qquad n = \frac{n_0}{1 + n_0/N},$$

where n_0 is the size required before the finite population correction. For stratified sampling, let N_h, n_h and S_h^2 be the population size, sample size and population variance of stratum h, with $W_h = N_h/N$. Then

$$\mathrm{Var}(\bar{y}_{st}) = \sum_h W_h^2(1 - n_h/N_h)S_h^2/n_h.$$

With equal unit costs, unconstrained Neyman allocation and its required size are

$$n_h = n\frac{N_h S_h}{\sum_h N_h S_h}, \qquad n = \frac{(\sum_h N_h S_h)^2}{N^2 V + \sum_h N_h S_h^2}.$$

The factors $1 - n_h/N_h$ are the finite population corrections. There are no clusters, so an intracluster correlation is not required. The planned design effect is the ratio of stratified to SRSWOR variance at the same respondent count; for the mean it is approximately 0.68.

Table 2: Neyman allocation, stratum variances and non-response allowance.

Region	Share	S_h^2 (USD^2)	Respondents	Selected
AU	0.2	16,601,039,692	615	684
UK	0.4	1,615,555,249	384	427
US	0.4	21,622,207,151	1,403	1,559

The continuous result rounds to 2,401 respondents; rounding **each stratum up** gives a feasible plan of **2,402**. This conservative integer allocation is not asserted to be the exact integer optimum. Assuming 90% independent response, selection counts are $\lceil n_h/0.9 \rceil$, totalling **2,670**. Inflation gives the desired response count in expectation, not a guarantee. The corresponding SRSWOR requirement is 3,178 respondents, so the stratified plan reduces respondent demand by about 24%.

For the secondary proportion, use $S_{ph}^2 = N_h P_h(1-P_h)/(N_h-1)$, where P_h is the proportion in stratum h. Its target is $0.05P = 0.03749$. An allocation separately optimised for the proportion needs 486 respondents (488 under SRSWOR); these are **not** the allocation used for the survey. The mean-based allocation still comfortably meets this secondary precision target. Holding this half-width fixed, the SRSWOR requirement is 241 at $P = 0.10$ or 0.90 but 640 at $P = 0.50$, because the variance $P(1-P)$ is largest at $P = 0.5$ and symmetric about it.

The unconstrained formula can imply $n_h > N_h$. The calculation caps such strata at a census and recalculates the optimum for the remaining strata. Census strata contribute zero sampling variance, while all population weights stay fixed. This constraint matters at tight precision targets, although no stratum reaches its population capacity in the final selection of 2,670 contestants.

The sensitivity analysis examines how the required sample size changes with the precision target and assumed baseline mean.




Figure 1: Required continuous sample size versus relative half-width. Stratified allocations respect each stratum's population capacity.

Figure 1 shows that the required size falls steeply as the permitted half-width widens, because n_0 is inversely proportional to e^2. The stratified curve lies below the SRSWOR curve at every margin, and the absolute saving is largest at tight margins, where both designs approach the population size and the finite population correction limits further growth.




Figure 2: Four assumed means with fixed stratum sizes and SDs. Tighter absolute margins require larger samples.

The quantities held fixed are $N = 10{,}000$, N_h, S_h, 95% confidence and the capacity-constrained Neyman rule. Only the assumed mean changes; each curve uses $e = qk\bar{Y}$, with $k = 0.25, 0.50, 0.75, 1$ and q the relative margin. Thus lower means imply smaller absolute margins, with SDs deliberately

unchanged. At $q = 5\%$, the required stratified sizes are **7,011, 5,064, 3,464, 2,401**, respectively. These are respondent planning sizes before non-response inflation. Curves show continuous optima; implementable allocations require integer rounding. A smaller assumed mean therefore raises the required sample sharply: at 25% of the baseline the stratified design needs 7,011 respondents (about 70% of N), against 8,817 under SRSWOR. As n approaches N, the finite population correction increasingly limits the required size, and the stratified design needs fewer respondents than SRSWOR at every assumed value.

2.3 Selection, weighting and data collection

The planned number of contestants is selected independently within each region using SRSWOR. Every population member has a known positive selection probability $\pi_i = n_h/N_h$, with design weight $d_i = 1/\pi_i$. Here n_h denotes selected contestants; the realised responding count is r_h.

Table 3: Selection probabilities and respondent weight adjustment.

Region	Selected	π_i	Design weight d_i	Respondents	Final weight w_i
AU	684	0.3420	2.924	609	3.284
UK	427	0.1067	9.368	377	10.610
US	1,559	0.3897	2.566	1,405	2.847

Independent Bernoulli response with probability 0.90 gives **2,391 responses (89.6%)**. Respondents receive $w_i = N_h/r_h$ and nonrespondents receive final weight zero. Final weights sum to each known regional count and to 10,000 overall. A UK respondent represents about 10.6 population members; a US respondent represents about 2.8. Equal weighting would overrepresent regions selected at higher rates.

Conditional on r_h, independent equal-probability response leaves an SRSWOR of r_h from region h. This justifies analysing respondents with weights N_h/r_h and respondent-level finite population corrections. It would not justify the same analysis if response depended on winnings within regions.

The population dataset contains all 10,000 individuals. The realised sample records all 2,670 selected individuals, including ID, stratum, inclusion probability, design weight, response flag, final weight, outcome and auxiliary variables. Outcomes are missing for nonrespondents. Latent risk aversion and wealth are simulation variables; the questionnaire's self-reported risk rating and wealth bands are only proposed proxies.

Contestants would receive an online questionnaire or telephone interview through the production company. The instrument in Appendix A records winnings in local currency, deal acceptance and round. With consent, production records would verify winnings. The opening expected value comes from the frame and determines the secondary indicator after currency conversion.

- **Coverage:** missing or outdated contestant records could exclude particular eras or regions. Reconcile registers and update contact details before selection.
- **Non-response:** winners with large payouts might respond differently. Follow-up and response comparisons on frame variables could help; weighting alone cannot remove unmeasured within-region selection bias.
- **Measurement:** recalled winnings can be rounded or confused with advertised prizes. Specify actual main-game payouts and verify records where permitted.

- **Processing and valuation:** currency miscoding changes outcomes. Check currency and amount ranges. Fixed conversion factors define this model's estimand but do not measure historical purchasing power.

3. Analysis

3.1 Realised estimates and design comparison

Design-based estimation accounts for independent individual selection, regional strata, final respondent weights and finite population corrections. Confidence intervals (CIs) use a normal approximation at the 95% level. The estimators are

$$\hat{\bar{Y}}_{st} = \sum_h W_h \bar{y}_h, \qquad \hat{V}(\hat{\bar{Y}}_{st}) = \sum_h W_h^2 (1 - r_h/N_h) s_h^2 / r_h.$$

Replacing y_i by $I(y_i < EV_{h(i)})$ estimates P and its variance. Here $\bar{y}_h$ and s_h^2 are the respondent mean and variance in stratum h. The coefficient of variation (CV) is the standard error (SE) divided by the estimate. The benchmark selects 2,670 individuals by SRSWOR, then simulates independent 90% response; 2,404 respond. With r respondents and respondent variance s^2, the weight is N/r and the estimated variance is $(1 - r/N)s^2/r$.

Table 4: Primary parameter: realised-sample estimates.

Measure	Stratified	SRSWOR
Estimate (USD)	67,453	69,418
Estimated variance	2,907,119	4,445,203
SE (USD)	1,705	2,108
95% CI	64,111 to 70,795	65,285 to 73,550
CV (%)	2.53	3.04
Survey design effect	0.654	1.000
CI half-width (USD)	3,342	4,132

The stratified interval covers the true mean and its half-width of $3,342 meets the $3,371 target. The benchmark half-width is $4,132 and misses the target. The naive stratified-sample mean is $84,806, an error of $17,376 (26%). It overstates the mean because US contestants, who have the highest winnings, make up 59% of respondents (1,405 of 2,391) but only 40% of the population; weighting reduces the realised error to $23. One sample's error is not itself an estimate of repeated-sampling bias.

Table 5: Secondary proportion; true value 0.7498.

Measure	Stratified	SRSWOR
Estimate	0.7528	0.7396
Estimated variance	0.0000962	0.0000609
SE	0.0098	0.0078
95% CI	0.7336 to 0.7720	0.7243 to 0.7549
CV (%)	1.30	1.05
Survey design effect	1.623	1.000

Both proportion intervals cover the truth and meet the 0.03749 target. The unweighted proportion is 0.7583, versus the weighted 0.7528. Because the proportion's stratum SDs are similar, its best allocation is close to proportional; the mean-driven allocation, which heavily undersamples the UK, therefore raises its variance.

The tabulated design effects compare each estimated variance with an estimated SRSWOR reference variance at the same respondent size. Ratios of variances from the two independently drawn realised samples are **0.654 for the mean**, essentially the same as the table, **and 1.579 for the proportion**. The latter differs from the table's 1.623 because the benchmark's respondent count and estimated variance differ from the reference variance used by the survey package.

3.2 Repeated-sampling performance

Selection, non-response, weighting and estimation are repeated **1,000 times per design** using the same fixed population. Every replicate retains estimates, SEs, interval limits and respondent counts.

Following Morris, White and Crowther (2019), performance is assessed by bias, relative bias, empirical SE, average estimated SE, root mean squared error (RMSE), coverage and interval width. For $B =$ 1,000 repetitions, true value θ, estimate $\hat{\theta}_b$ in repetition b and estimated coverage $\hat{c}$ (the share of intervals containing θ),

$$\widehat{\text{Bias}} = B^{-1}\sum_b \hat{\theta}_b - \theta, \quad \text{RMSE} = \sqrt{B^{-1}\sum_b (\hat{\theta}_b - \theta)^2}, \quad \text{MCSE}(\hat{c}) = \sqrt{\hat{c}(1-\hat{c})/B},$$

where MCSE is the Monte Carlo standard error. Relative bias is bias divided by θ. Empirical SE is the sample SD of estimates; the empirical design effect is the ratio of their variances, stratified over SRSWOR.

Table 6: Performance over 1,000 repetitions. pp = percentage points.

Measure	Mean: strat.	Mean: SRS	Prop.: strat.	Prop.: SRS
Average estimate	67,459.64	67,332.89	0.7496	0.7497
Bias	29.88	-96.87	-0.0002	-0.0001
Relative bias (%)	0.044	-0.144	-0.023	-0.008
Empirical SE	1,757.86	2,125.04	0.0100	0.0077
Average estimated SE	1,718.13	2,082.70	0.0098	0.0077
RMSE	1,757.23	2,126.19	0.0100	0.0077
Coverage (%)	94.9	94.4	94.3	95.2
Coverage MCSE (pp)	0.70	0.73	0.73	0.68
Average CI width	6,734.95	8,164.02	0.0385	0.0302
Empirical design effect	0.68	1.00	1.67	1.00
Meets bound (%)	52.1	0.0	100.0	100.0

All relative biases are below 0.15% in magnitude. The mean's bias MCSEs are approximately \$56 (stratified) and \$67 (SRSWOR); observed biases of \$30 and -\$97 are within two MCSEs of zero. Coverage ranges from 94.3% to 95.2%, with MCSEs around 0.7 percentage points, consistent with nominal 95% coverage at this simulation resolution.

Average estimated SEs for the mean are about 2% below empirical SEs. This is a descriptive comparison, not proof that the variance estimators are biased: average SE and average variance are different measures, and simulation estimates have Monte Carlo error.

Stratification reduces the mean's empirical variance by 32% and RMSE by 17%; its average interval is \$6,735 wide versus \$8,164 for SRSWOR. For the proportion, variance increases by 67%, demonstrating the trade-off from optimising allocation for winnings.




Figure 3: Distribution of realised primary-parameter interval half-widths over 1,000 repetitions. The dashed line marks the 5% target.

The stratified design meets the mean's precision bound in **52.1%** of repetitions; the same-sized SRSWOR design meets it in none. The sample was planned using baseline variances and expected response counts. The planning equation sets the expected variance exactly equal to the bound, with no margin. Realised response counts and, more importantly, estimated stratum variances fluctuate around their planning values, so the estimated half-width scatters around the target and roughly half of the samples fall on either side of it. Satisfying the planning equation therefore does not guarantee that every estimated interval meets the bound. The approximately one-half rate is specific to this experiment, not a universal property of sample-size formulas.

For a practical survey, planning for a tighter half-width, such as 4.5%, or a lower anticipated response rate would provide a buffer. Such choices should be evaluated against cost and desired assurance before collecting data. Both designs meet the secondary proportion's precision target in every repetition here.

4. Conclusion

For this finite population, mean winnings are about **\$67,400** and approximately **75%** win less than their board's opening expected value. Neyman allocation reduces the mean's required respondents by 24% and empirical variance by 32%, while sacrificing precision for the proportion. Weighting is essential. Bias and coverage results support the design under the stated assumptions; practical planning should allow a buffer beyond the nominal precision target.

The simulated outcome distribution depends on assumed prize boards, offer progression, wealth and risk aversion. The Australian high-stakes setting is intentionally stylised, and fixed currency factors restrict interpretation. The model does not estimate real contestants' behaviour or regional winnings. The age distribution is clipped at its limits rather than sampled from a truncated normal distribution.

Sampling and variance calculations also assume a complete frame, equal within-region response propensities and accurate measurements. Non-response weights are valid for the simulated mechanism but would not automatically remove outcome-dependent response bias in a real survey. The normal intervals appear adequate in this large-sample experiment; this does not establish their performance for much smaller samples or other populations.

References

Cochran, W. G. (1977). *Sampling techniques* (3rd ed.). Wiley.

Lohr, S. L. (2010). *Sampling: Design and analysis* (2nd ed.). Brooks/Cole.

Lumley, T. (2004). Analysis of complex survey samples. *Journal of Statistical Software, 9*(8), 1–19. https://doi.org/10.18637/jss.v009.i08

Morris, T. P., White, I. R., & Crowther, M. J. (2019). Using simulation studies to evaluate statistical methods. *Statistics in Medicine, 38*(11), 2074–2102. https://doi.org/10.1002/sim.8086

Neyman, J. (1934). On the two different aspects of the representative method: The method of stratified sampling and the method of purposive selection. *Journal of the Royal Statistical Society, 97*(4), 558–625.

Post, T., van den Assem, M. J., Baltussen, G., & Thaler, R. H. (2008). Deal or no deal? Decision making under risk in a large-payoff game show. *American Economic Review, 98*(1), 38–71. https://doi.org/10.1257/aer.98.1.38

Appendix A: Proposed questionnaire

Purpose and consent. This proposed academic survey concerns main-game winnings. Participation is voluntary; respondents may skip an item or stop. Responses would be used for aggregate analysis. A real deployment would first add approved consent, privacy, retention and researcher-contact details.

Confirm the contestant's ID and regional version against the register.

1. What amount did you actually receive from the main game, before tax or personal expenses? Record the amount and currency (GBP / USD / AUD). Exclude unrelated promotional awards. Allow "do not recall".
2. Did you accept a banker offer? (Yes / No / Do not recall.)
3. If yes, after which round? Record the round number or "do not recall".
4. At the time you played, how willing were you to take financial risks, from 0 (not willing) to 10 (very willing)? Optional.
5. At that time, what was your approximate household savings balance in the show's local currency? (Below 5,000 / 5,000–19,999 / 20,000–49,999 / 50,000 or more / Prefer not to say.) Optional.
6. May the production company confirm your reported winnings from its records? (Yes / No.)

Region, age, gender and opening expected value are frame variables. Items 4 and 5 are contextual proxies and are not used to estimate the primary mean. The exact CRRA coefficient and wealth used in the simulation would not be directly observed by this instrument. No real respondents were contacted for this project.

Appendix B: Files and reproducibility

Run `Rscript run_all.R` from the project folder. It executes scripts in isolated environments, checks the outputs, records `sessionInfo()` and renders this report. R packages required are `survey`, `ggplot2`, `scales`, `knitr` and `rmarkdown`; PDF rendering also requires Pandoc and XeLaTeX. Use `--no-render` to run just the analysis and checks.

All code, data, outputs and the README are available in the project's GitHub repository: https://github.com/Priyan-Sathabridge/Applied-Sampling-Project.

File	Role / seed
`01_simulate_population.R`	Generate population; 20261019
`02_sample_size.R`	Planning and capacity-constrained sensitivity
`03_draw_sample.R`	Stratified selection and response; 20261020
`04_estimation.R`	Survey estimates and benchmark; 20261021
`05_simulation.R`	1,000 repetitions per design; 20261022
`06_extra_figures.R`	Additional deterministic figures
`validate.R`	Data and output integrity checks
`report.Rmd`	Report source; tables read executed output files

Estimation uses the R *survey* package (Lumley, 2004), with regional strata, respondent weights and population counts declared in `svydesign`. The simulation checks closed-form estimates and standard errors against this implementation for both outcomes and both design formulas before repeating sampling; the checks consume no random numbers. Seeds are listed above. The output retains average estimated variances as well as the performance measures reported in the analysis.

The `data/` folder contains the full population and both selected samples. The `output/` folder contains planning tables, estimates, replicate results, performance summaries and software versions. The README provides run instructions and a field dictionary.

Appendix C: Declaration of generative AI use

Tools used: Claude.

Parts of the project for which AI was used: Simulation design and code. Error and bug checking. Grammar and spell check. Text generation and formatting.

How the work was reviewed and verified: All code was checked against known theory and academic literature. validate.R script performs integrity checks; closed-form estimators checked against the survey package; reported numbers independently recomputed from the data; all text reviewed and edited by the author.