

Practical Sheet in R

Distributions

Binomial Distribution

R has four in-built functions to generate binomial distribution. They are described below.

```
dbinom(x, size, prob)
pbinom(x, size, prob)
qbinom(p, size, prob)
rbinom(n, size, prob)
```

Following is the description of the parameters used –

- **x** is a vector of numbers.
- **p** is a vector of probabilities.
- **n** is number of observations.
- **size** is the number of trials.
- **prob** is the probability of success of each trial.

dbinom()

This function gives the probability density distribution at each point.

```
x <- seq(0,50,by = 1)
y <- dbinom(x,50,0.5) (# Creates a binomial distribution for 50 numbers with  $p = 0.5$ )
png(file = "dbinom.png")
plot(x,y)
dev.off()
```

pbinom()

This function gives the cumulative probability of an event. It is a single value representing the probability.

Probability of getting 26 or less heads from a 51 tosses of a coin.

```
print(x <- pbinom(26,51,0.5))
```

qbinom()

This function takes the probability value and gives a number whose cumulative value matches the probability value.

How many heads will come out with a probability of 0.25 when a coin is tossed 51 times

```
print(x <- qbinom(0.25,51,1/2))
```

rbinom()

This function generates required number of random values of given probability from a given sample.

Find 8 random values from a sample of 150 with probability of 0.4.

```
print(x <- rbinom(8,150,.4))
```

Poisson Distribution

R has four in-built functions to generate poisson distribution. They are described below.

```
dpois(x, lambda)
```

```
ppois(q, lambda)
```

```
qpois(p, lambda)
```

```
rpois(n, lambda)
```

- x is a vector of numbers.
- p is a vector of probabilities.
- q is the vector of quantiles
- n is number of observations.

dpois()

This function gives the probability density distribution at each point.

```
x <- seq(0,20,by = 1)
```

```
y<-dpois(x,1)
```

```
png(file = "dpois.png")
```

```
plot(x,y)
```

```
dev.off()
```

ppois()

If there are twelve cars crossing a bridge per minute on average, find the probability of having sixteen or less cars crossing the bridge in a particular minute.

```
ppois(16, lambda=12)
```

If there are twelve cars crossing a bridge per minute on average, find the probability of having seventeen or more cars crossing the bridge in a particular minute.

```
ppois(16, lambda=12, lower=FALSE)
```

qpois()

$qpois(p, \lambda)$ is the smallest integer x such that $P(X \leq x) \geq p$.

Find the smallest integer x such that $P(X \leq x) \geq 0.75$.

```
qpois(0.75, lambda=12)
```

rpois()

Find 16 values of a random value that is Poisson distributed with mean = 16

```
rpois(16, lambda=12)
```

Geometric Distribution

R has four in-built functions to generate geometric distribution. They are described below.

```
dgeom(x, prob, log = FALSE )  
pgeom(q, prob, lower.tail = TRUE, log.p = FALSE)  
qgeom(p, prob, lower.tail = TRUE, log.p = FALSE)  
rgeom(n, prob)
```

- x is a vector of numbers of failures in a sequence of Bernoulli trials before success occurs
- **prob** is the success of each trial.
- p is a vector of probabilities.
- q is the vector of quantiles
- n is number of observations.
- Lower.tail = TRUE represents $P[X \leq x]$ else $P[X > x]$

dgeom()

This function gives the probability density distribution at each point.

```
x <- seq(0,20,by = 1)  
y<-dgeom(x,0.6,log = FALSE)  
png(file = "dgeom.png")  
plot(x,y)  
dev.off()
```

pgeom()

The probability of having at most three failures before a success with probability of success 0.1

```
pgeom(3, 0.1, lower.tail = TRUE, log.p = FALSE)
```

qgeom()

What is the smallest value of x such that $P(X \leq x) \geq 0.3$ for a geometric distribution with probability of success 0.1?

```
qgeom(0.3, 0.1, lower.tail = TRUE, log.p = FALSE)
```

Gamma Distribution

R has four in-built functions to generate geometric distribution. They are described below.

```
dgamma(x, k, theta, log = FALSE )  
pgamma(q, k, theta, lower.tail = TRUE, log.p = FALSE)  
qgeom(p, k, theta, lower.tail = TRUE, log.p = FALSE)  
rgeom(n, k, theta)
```

- x is a vector of numbers of failures in a sequence of Bernoulli trials before success occurs
- k is the shape parameter
- θ is the scale parameter.
- p is a vector of probabilities.
- q is the vector of quantiles
- n is number of observations.
- Lower.tail = TRUE represents $P[X \leq x]$ else $P[X > x]$

dgamma()

This function gives the probability density distribution at each point.

```
x <- seq(0,20,by = 0.01)  
y<-dgamma(x,2,2,log = FALSE)  
png(file = "dgamma.png")  
plot(x,y)  
dev.off()
```

Exercise

Q1) Plot the curve for a Binomial distribution with $n = 10$, $p = 0.4$. Increase the value of n in steps of 5 till 30. What do you observe?

Q2) Plot the curve for a Poisson distribution with $\lambda = 0.1, 1, 10, 20$. What do you observe?

Q3) Plot the curve for a Binomial distribution with $n = 50$, $p = 0.1$. Plot the curve for Poisson distribution with $\lambda = 5$. Give your observation.

Q4) Try to do a similar exercise by changing the value of p in the geometric distribution. Can you explain the trend with mathematical explanation

Q5) Plot the curves for gamma distribution for $\theta = 2, 1, 0.5$ and $k = 1, 2, 3, 5, 9$