



Department of Mathematics and Natural Sciences  
MAT215 : Complex Variables and Laplace Transformations  
Assignment 1  
Date: June 28, 2025

Deadline : July 6,2024

Summer 2025

Total Marks: 90

Name: Priyanti Biswas

Section: 04

ID: 23201039

Use this page as the cover page of your assignment. No late submission will be graded.

(1) Express each of the following complex numbers in polar form. (any 1)

(a)  $2 + 2\sqrt{3}i$ , (b)  $-\sqrt{6} - \sqrt{2}i$ , (c)  $-3i$ . [5]

(2) Find each of the indicated roots and locate them graphically. (any 1)

(a)  $(-1 + i)^{1/3}$ , (b)  $(-2\sqrt{3} - 2i)^{1/4}$  [5]

(3) Describe graphically the region represented by each of the following: (any 2)

(a)  $1 < |z + i| \leq 2$ , (b)  $\operatorname{Re}\{z^2\} > 1$ , (c)  $|z + 3i| > 4$ , (d)  $|z + 2 - 3i| + |z - 2 + 3i| < 10$ , (e)  $\operatorname{Re}(1/z) \leq 1/2$

(f)  $\pi/2 < \arg z < 3\pi/2, |z| > 2$  [6]

(4) Show that,  $\ln(1 - i) = \frac{1}{2}\ln 2 + \left(2n + \frac{7}{4}\right)\pi i$  [5]

(5) Prove that (any 1)

(a)  $\cos^{-1} z = \frac{1}{i} \ln \left(z + \sqrt{z^2 - 1}\right)$ , (b)  $\cot^{-1} z = \frac{1}{2i} \ln \left(\frac{z+i}{z-i}\right)$  indicating any restrictions. [5]

(6) Prove that  $\lim_{z \rightarrow 1+i} \frac{z^2 - z + 1 - i}{z^2 - 2z + 2} = 1 - \frac{1}{2}i$ . [5]

(7) Let  $f(z) = \frac{2z-1}{3z+2}$ . Prove that  $\lim_{h \rightarrow 0} \frac{f(z_0+h) - f(z_0)}{h} = \frac{7}{(3z_0+2)^2}$  provided  $z_0 \neq -2/3$ . [5]

(8) If  $\left(\frac{3}{2} + \frac{i\sqrt{3}}{2}\right)^{50} = 3^{25}(x+iy)$ , where  $x$  and  $y$  are reals then find the values of  $x$  and  $y$ . [5]

(9) Evaluate using theorems on limits. (any 2)

(a)  $\lim_{z \rightarrow 2i} (iz^4 + 3z^2 - 10i)$ , (b)  $\lim_{z \rightarrow i/2} \frac{(2z-3)(4z+i)}{(iz-1)^2}$ ,

$$(c) \lim_{z \rightarrow e^{\pi/4}} \frac{z^2}{z^4 + z + 1}, (d) \lim_{z \rightarrow i} \frac{z^2 + 1}{z^6 + 1}, (e) \lim_{z \rightarrow 1+i} \left\{ \frac{z - 1 - i}{z^2 - 2z + 2} \right\}^2$$

(10) Use exponential form to compute (any 1)

$\checkmark (1 + \sqrt{3}i)^{2011}$ ; ii.  $(1 + \sqrt{3}i)^{-2011}$ .

[5]

(11) Sketch the region in xy plane represented by the following set of points: (any 2)

$\checkmark (a) \operatorname{Re}(\bar{z} - 1) = 2$  (b)  $\operatorname{Im}(z^2) = 4$  (c)  $\left| \frac{2z - 3}{2z + 3} \right| = 1$  (d)  $\operatorname{Re}(z) + \operatorname{Im}(z) = 0$

[4]

(12) Solve the following equations: (any 1)

$\checkmark (i) \cosh z = \frac{1}{2}$  (ii)  $\sinh z = i$ .

[5]

(13) Find all the roots of the following equations. (any 1)

$\checkmark (i) z^6 = 64$  (ii)  $z^4 + z^2 + 1 = 0$ .

[5]

(14) Let  $f(z) = \frac{z^2 + 4}{z - 2i}$  if  $z \neq 2i$ , while  $f(2i) = 3 + 4i$ .

[5]

(a) Prove that  $\lim_{z \rightarrow 2i} f(z)$  exists and determine its value (b) Is  $f(z)$  continuous at  $z = 2i$ ? Explain. (c) Is  $f(z)$  continuous at points  $z \neq 2i$ ? Explain.

(15) Find all values of  $z$  such that: (any 1)

$\checkmark (i) e^z = 1 + \sqrt{3}i$  (ii)  $e^{2z-1} = 1$

[5]

(16) Evaluate the following Limits using L'Hospital's rule (any 2)

$\checkmark (i) \lim_{z \rightarrow 2i} \frac{z^2 + 4}{2z^2 + (3 - 4i)z - 6i}$  (ii)  $\lim_{z \rightarrow 0} \frac{z - \sin z}{z^3}$  (iii)  $\lim_{z \rightarrow 0} \left( \frac{\sin z}{z} \right)^{\frac{1}{z}}$

[5]

(17) Find the modulus and argument of the following complex numbers: (any 1)

(i)  $\frac{\sqrt{3} + i}{\sqrt{3} - i}$  (ii)  $\left( \frac{1 + \sqrt{3}i}{1 - \sqrt{3}i} \right)^2$

[5]

(18) If  $f(z) = \begin{cases} \frac{z^2 - 4}{z^2 - 3z + 2}, & z \neq 2 \\ kz^2 + 6, & z = 2 \end{cases}$ , find  $k$  such that  $f(z)$  becomes continuous at  $z = 2$ .

[5]

$$1. a) 2 + 2\sqrt{3} i$$

$$\text{modulus, } r = \sqrt{(2)^2 + (2\sqrt{3})^2} = 4$$

$$\text{Argument, } \theta = \tan^{-1} \left| \frac{2\sqrt{3}}{2} \right| = \pi/3$$

$$\therefore \text{in polar form, } z = 4e^{i\pi/3}$$

$$2. a) (-1+i)^{1/3}$$

Now,  $z = (-1+i)^{1/3}$  consider  $(-1+i)$ :  $\begin{matrix} x = -1 \\ y = 1 \end{matrix}$

$$\text{mod}(z) = \sqrt{(-1)^2 + (1)^2} = \sqrt{2}$$

$$\text{Arg}(z) = \pi - \tan^{-1} \left| \frac{1}{-1} \right| + 2n\pi = \pi - \frac{\pi}{4} + 2n\pi = \frac{3\pi}{4} + 2n\pi$$

$$(-1+i)^{1/3} = (\sqrt{2})^{1/3} \left( \cos \frac{1}{3} \left( \frac{3\pi}{4} + 2n\pi \right) + i \sin \frac{1}{3} \left( \frac{3\pi}{4} + 2n\pi \right) \right)$$

$$\left[ \because (\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta) \right]$$

$$= (\sqrt{2})^{1/3} \cos \left( \frac{\pi}{4} + \frac{2n\pi}{3} \right) + i \sin \left( \frac{\pi}{4} + \frac{2n\pi}{3} \right)$$


$$\text{no. of roots} = 3$$

$$n = 0, 1, 2$$

$$n=0, z_0 = (\sqrt{2})^{1/3} \left( \cos \left( \frac{\pi}{4} \right) + i \sin \left( \frac{\pi}{4} \right) \right) = 0.707 + 0.707i$$

$$n=1, z_1 = (\sqrt{2})^{1/3} \left( \cos \left( \frac{\pi}{4} + \frac{2\pi}{3} \right) + i \sin \left( \frac{\pi}{4} + \frac{2\pi}{3} \right) \right) = -1.08 + 0.29i$$

$$n=2, z_2 = (\sqrt{2})^{1/3} \left( \cos \left( \frac{\pi}{4} + \frac{4\pi}{3} \right) + i \sin \left( \frac{\pi}{4} + \frac{4\pi}{3} \right) \right) = 0.29 - 1.08i$$



$$\frac{360^\circ}{3} = 120^\circ$$

~~Re~~ Re

Regular Triangle

$$3. e) \operatorname{Re}(\frac{1}{z}) \leq \frac{1}{2}$$

$$\Rightarrow \operatorname{Re}\left(\frac{1}{x+iy}\right) \leq \frac{1}{2}$$

$$\Rightarrow \operatorname{Re}\left(\frac{1(x-iy)}{x^2+y^2}\right) \leq \frac{1}{2}$$

$$\Rightarrow \frac{x}{x^2+y^2} \leq \frac{1}{2}$$

$$\Rightarrow 2x \leq x^2+y^2$$


$$\Rightarrow x^2+y^2 \geq 2x$$

$$\Rightarrow x^2 - 2x + y^2 \geq 0$$

$$\Rightarrow x^2 - 2x + 1 + y^2 \geq 1$$

$$\Rightarrow (x-1)^2 + (y-0)^2 \geq 1^2$$

$\Rightarrow$



$$f) \pi/2 < \underbrace{\arg z}_0 < 3\pi/2, |z| > 2$$

Given,  $|z| > 2$

$$\Rightarrow 90^\circ < \arg z < 270^\circ$$


$$\Rightarrow \sqrt{x^2+y^2} > 2$$

$$\Rightarrow 90^\circ < \theta < 270^\circ$$

$$\Rightarrow x^2+y^2 > 2^2$$

center (0, 0)

$$r = 2$$



$$4. \quad 1-i \Rightarrow x=1, y=-1$$

$$r = \sqrt{(1)^2 + (-1)^2} = \sqrt{2}$$

$$\theta = -\tan^{-1}\left|\frac{-1}{1}\right| = -\frac{\pi}{4}$$

Now,  $i(\theta + 2n\pi), n > 0$

$$1-i = r e^{i(\frac{7\pi}{4} + 2n\pi)}$$

$$= \sqrt{2} e^{i(\frac{7\pi}{4} + 2n\pi)}$$

$$\ln(1-i) = \ln(\sqrt{2} e^{i(2n\pi + \frac{7\pi}{4})})$$

$$= \ln \sqrt{2} + \ln e$$

$$= \ln 2^{\frac{1}{2}} + (2n\pi + \frac{7\pi}{4})i$$

$$= \frac{1}{2} \ln 2 + (2n\pi + \frac{7\pi}{4})i$$

$$= \frac{1}{2} \ln 2 + (2n + \frac{7}{4})\pi i$$

$$\therefore \ln(1-i) = \frac{1}{2} \ln 2 + (2n + \frac{7}{4})\pi i \quad [\text{Proved}]$$

$$5. b) \cot^{-1} z = \theta$$

$$\Rightarrow z = \cot \theta = \frac{\cos \theta}{\sin \theta} = \frac{e^{i\theta}}{e^{-i\theta}}$$

$$\Rightarrow z = \frac{e^{i\theta} + e^{-i\theta}}{2i}$$

$$\Rightarrow \frac{z+i}{z-i} = \frac{e^{i\theta} - e^{-i\theta}}{e^{i\theta} + e^{-i\theta}}$$

$$\Rightarrow \frac{z+i}{z-i} = \frac{e^{i\theta}}{e^{-i\theta}}$$

$$\Rightarrow e^{2i\theta} = \frac{z+i}{z-i}$$

$$\Rightarrow \ln(e^{2i\theta}) = \ln\left(\frac{z+i}{z-i}\right)$$

$$\Rightarrow 2i\theta = \ln\left(\frac{z+i}{z-i}\right)$$

$$\Rightarrow \theta = \frac{1}{2i} \ln \left( \frac{z+i}{z-i} \right)$$

$$\therefore \cot^{-1} z = \frac{1}{2i} \ln \left( \frac{z+i}{z-i} \right) \quad [\text{Proved}]$$

This expression is undefined at  $z = \pm i$  because of division by zero.

The logarithm is multivalued and undefined when the argument is zero or negative real.

So, we avoid:

$$z = i \Rightarrow \ln(0)$$

$$z = -i \Rightarrow \ln(\infty)$$

$$6. \lim_{z \rightarrow 1+i} \frac{z^2 - z + 1 - i}{z^2 - 2z + 2}$$

$$= \lim_{z \rightarrow 1+i} \frac{2z - 1}{2z - 2} \quad [\because \text{Apply L'Hospital's Rule}]$$

$$= \lim_{z \rightarrow 1+i} \frac{2(1+i) - 1}{2(1+i) - 2}$$

$$= \frac{2+2i-1}{2+2i-2}$$

$$= \frac{1+2i}{2i}$$

$$= \frac{(1+2i)}{2i} \cdot \frac{(-2i)}{(-2i)}$$

$$= \frac{-4i^2 - 2i}{-4i^2} = \frac{4-2i}{4} = 1 - \frac{1}{2}i$$

$$\therefore \lim_{z \rightarrow 1+i} \frac{z^2 - z + 1 - i}{z^2 - 2z + 2} = 1 - \frac{1}{2}i \quad [\text{Proved}]$$

$$7. \quad f(z) = \frac{2z-1}{3z+2}$$

$$f(z_0 + h) = \frac{2(z_0 + h) - 1}{3(z_0 + h) + 2} = \frac{2z_0 + 2h - 1}{3z_0 + 3h + 2}$$

$$f(z_0) = \frac{2z_0 - 1}{3z_0 + 2}$$

$$\begin{aligned} f(z_0 + h) - f(z_0) &= \frac{2z_0 + 2h - 1}{3z_0 + 3h + 2} - \frac{2z_0 - 1}{3z_0 + 2} \\ &= \frac{(2z_0 + 2h - 1)(3z_0 + 2) - (2z_0 - 1)(3z_0 + 3h + 2)}{(3z_0 + 3h + 2)(3z_0 + 2)} \\ &= \frac{6z_0^2 + 6z_0 h - 3z_0 + 4z_0 + 4h - 2 - 6z_0^2 + 3z_0 - 6z_0 h + 3h - 4z_0 + 2}{(3z_0 + 3h + 2)(3z_0 + 2)} \end{aligned}$$

$$= \frac{7h}{(3z_0 + 3h + 2)(3z_0 + 2)}$$

$$L.H.S = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\cancel{7h}}{(3z_0 + 3h + 2)(3z_0 + 2)} \times \frac{1}{\cancel{h}}$$

$$= \lim_{h \rightarrow 0} \frac{\cancel{7}}{(3z_0 + 3h + 2)(3z_0 + 2)}$$

$$= \frac{\cancel{7}}{(3z_0 + 2)(3z_0 + 2)}$$

$$= \frac{\cancel{7}}{(3z_0 + 2)^2} = R.H.S$$

[Proved]

$$8. z = \frac{3}{2} + \frac{i\sqrt{3}}{2}$$

$$r = \sqrt{\left(\frac{3}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} = \sqrt{3}$$

$$\theta = \tan^{-1}\left(\frac{\sqrt{3}/2}{3/2}\right) = \pi/6$$

$$z \text{ in polar form: } z = \sqrt{3} \left( \cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right)$$

Now,

$$\left(\frac{3}{2} + \frac{i\sqrt{3}}{2}\right)^{50} = 3^{25} (x+iy)$$

$$\Rightarrow (\sqrt{3})^{50} \left( \cos \frac{50\pi}{6} + i \sin \frac{50\pi}{6} \right) = 3^{25} (x+iy)$$

$$\Rightarrow 3^{25} \left( \cos \frac{50\pi}{6} + i \sin \frac{50\pi}{6} \right) = 3^{25} (x+iy)$$

$$\begin{cases} x = \cos\left(\frac{50\pi}{6}\right) = \frac{1}{2} \\ y = \sin\left(\frac{50\pi}{6}\right) = \frac{\sqrt{3}}{2} \end{cases}$$

$$9. a) \lim_{z \rightarrow 2i} (iz^4 + 3z^2 - 10i)$$

$$= i(2i)^4 + 3(2i)^2 - 10i$$

$$= i(16i^4) + 3(4i^2) - 10i$$

$$= 16i^5 + 12i^2 - 10i$$

$$= 16i - 12 - 10i$$

$$= 6i - 12$$

$$i^5 = i^2 \cdot i^2 \cdot i = i$$

$$b) \lim_{z \rightarrow i/2} \frac{(2z-3)(4z+i)}{(iz-1)^2}$$

$$= \frac{(i-3)(2i+i)}{i^2 \cdot \frac{i^2}{4} - 2i \cdot \frac{i}{2} + 1}$$

$$= \frac{(i-3) \cdot 3i}{\frac{1}{4} + 1 + 1}$$

$$= \frac{3i - 9i}{\frac{1}{4} + 2}$$

$$= \frac{-3 - 9i}{\frac{9}{4}}$$

$$= (-3 - 9i) \times \frac{4}{9}$$

$$= -\frac{12}{9} - \frac{36i}{9}$$

$$= -\frac{4}{3} - 4i$$

$$10. \text{ i) } (1 + \sqrt{3}i)^{2011}$$

$$z = (1 + \sqrt{3}i)$$

$$r = \sqrt{(1)^2 + (\sqrt{3})^2} = \sqrt{4} = 2$$

$$\theta = \tan^{-1} \left| \frac{\sqrt{3}}{1} \right| = \frac{\pi}{3} + 2n\pi \quad \frac{\pi}{3} + 2n\pi$$

$$z \text{ in exponential form} = 2^{2011} e^{2011 \cdot (\frac{\pi}{3} + 2n\pi)}$$

$$\text{Now, } (1 + \sqrt{3}i)^{2011} = 2^{2011} \cdot e^{2011 \cdot \frac{\pi}{3}}$$

$$= 2^{2011} \cdot \left\{ \cos \left( \frac{2011\pi}{3} + \frac{4022n\pi}{3} \right) + i \sin \left( \frac{2011\pi}{3} + \frac{4022n\pi}{3} \right) \right\}$$

$$= 2^{2011} \cdot \left\{ \cos\left(\frac{2011\pi}{3}\right) + i \sin\left(\frac{2011\pi}{3}\right) \right\} [n=0]$$

$$= 2^{2011} \cdot \left\{ \frac{1}{2} + i \sin \frac{\sqrt{3}}{2} \right\}$$

$$= \frac{2^{2011}}{2} + \frac{\sqrt{3} \cdot 2^{2011}}{2} i$$

$$= 2^{2010} + \sqrt{3} \cdot 2^{2010} i$$

11. a)  $\operatorname{Re}(\bar{z} - 1) = 2$

We know,

$$z = x + iy$$


$$\bar{z} = x - iy$$

$$\operatorname{Re}(x - iy - 1) = 2$$

$$\Rightarrow x - 1 = 2$$

$$\Rightarrow x = 2 + 1$$

$$\therefore x = 3$$



$$b) \operatorname{Im}(z^2) = 4$$

We know,

$$z = x + iy$$

Now,

$$\operatorname{Im}(z^2) = 4$$


$$\Rightarrow \operatorname{Im}\{(x+iy)^2\} = 4$$

$$\Rightarrow \operatorname{Im}(x^2 + 2xyi - y^2) = 4$$

$$\Rightarrow 2xy = 4$$

$$\Rightarrow y = 2/x$$

|     |    |   |    |   |      |     |      |     |
|-----|----|---|----|---|------|-----|------|-----|
| $x$ | -1 | 1 | -2 | 2 | -3   | 3   | -4   | 4   |
| $y$ | -2 | 2 | -1 | 1 | -2/3 | 2/3 | -1/2 | 1/2 |



$$12. 1) \cosh z = 1/2$$

$$\Rightarrow \frac{1}{2} (e^z + e^{-z}) = 1/2$$

$$\Rightarrow e^z + e^{-z} = 1$$

$$\Rightarrow (e^z)^2 + 1 = e^z \quad [\text{multiply by } e^z \text{ to avoid the -ve power}]$$

$$\Rightarrow (e^z)^2 - e^z + 1 = 0$$

$$\Rightarrow e^z = \frac{1 \pm \sqrt{1-4}}{2}$$

$$\Rightarrow e^z = \frac{1 \pm i\sqrt{3}}{2}$$

$$\Rightarrow e^z = \frac{1}{2} \pm i \frac{\sqrt{3}}{2}$$

Consider the complex number  $\frac{1}{2} \pm i \frac{\sqrt{3}}{2}$  for  $e^z = \frac{1}{2} - i \frac{\sqrt{3}}{2}$

$$\text{for } e^z = \frac{1}{2} + i \frac{\sqrt{3}}{2}$$



$$r = \sqrt{\left(\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} = 1$$

$$\theta = \tan^{-1}\left(\frac{\sqrt{3}/2}{1/2}\right) = \pi/3$$

$$r = \sqrt{\left(\frac{1}{2}\right)^2 + \left(-\frac{\sqrt{3}}{2}\right)^2} = 1$$

$$\theta = -\tan^{-1}\left(-\frac{\sqrt{3}/2}{1/2}\right) = -\pi/3$$

$$= 2\pi - |\pi/3| = \frac{5\pi}{3}$$



$$\text{For, } e^z = \frac{1}{2} + i \frac{\sqrt{3}}{2} = r e^{i(\theta + 2n\pi)}, n \geq 0$$

$$= 1 e^{i(\pi/3 + 2n\pi)}$$

$$e^z = e^{i(\pi/3 + 2n\pi)}$$

$$\Rightarrow z = i(\pi/3 + 2n\pi) = \pi i (1/3 + 2n); n \geq 0$$

$$\text{form, } e^z = \frac{1}{2} - \frac{i\sqrt{3}}{2} = re^{i(\theta + 2n\pi)}, n > 0$$

$$= 1e^{i\left(\frac{5\pi}{3} + 2n\pi\right)}$$

$$\Rightarrow e^z = e^{i\left(\frac{5\pi}{3} + 2n\pi\right)}$$

$$\Rightarrow z = i\left(\frac{5\pi}{3} + 2n\pi\right)$$

$$= ri\left(\frac{5}{3} + 2n\right); n > 0$$

$$13.1) z^6 = 64^{1/6} = (64+0i)^{1/6}$$

$$\Rightarrow z = (64)^{1/6} = \sqrt[6]{(64)^n} = 64$$

$$\tan^{-1}\left(\frac{0}{64}\right) = 0^\circ$$

$$z = r^{1/6} \left( \cos \frac{1}{6}(\theta + 2n\pi) + i \sin \frac{1}{6}(\theta + 2n\pi) \right)$$

$$= (64)^{1/6} \left( \cos \frac{1}{6}(0 + 2n\pi) + i \sin \frac{1}{6}(0 + 2n\pi) \right)$$

$$= 2 \left( \cos \frac{n\pi}{3} + i \sin \frac{n\pi}{3} \right)$$

6 roots means  $n = 0, 1, 2, 3, 4, 5$

$$n=0, z_0 = 2 \left( \cos 0^\circ + i \sin 0^\circ \right) = 2(1+0i) = 2$$

$$n=1, z_1 = 2 \left( \cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \right) = 2 \left( \frac{1}{2} + i \frac{\sqrt{3}}{2} \right) = 1 + i\sqrt{3}$$

$$n=2, z_2 = 2 \left( \cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3} \right) = 2 \left( -\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) = -1 + i\sqrt{3}$$

$$n=3, z_3 = 2 \left( \cos \frac{3\pi}{3} + i \sin \frac{3\pi}{3} \right) = 2(-1+i0) = -2$$

$$n=4, z_4 = 2 \left( \cos \frac{4\pi}{3} + i \sin \frac{4\pi}{3} \right) = 2 \left( -\frac{1}{2} - i \frac{\sqrt{3}}{2} \right) = -1 - i\sqrt{3}$$

$$n=5, z_5 = 2 \left( \cos \frac{5\pi}{3} + i \sin \frac{5\pi}{3} \right) = 2 \left( \frac{1}{2} - i \frac{\sqrt{3}}{2} \right) = 1 - i\sqrt{3}$$

$$14. f(z) = \begin{cases} \frac{z^2 + 4}{z - 2i}, & z \neq 2i \\ 3 + 4i, & z = 2i \end{cases}$$

a)  $\lim_{z \rightarrow i} f(z)$

$$= \lim_{z \rightarrow i} \frac{z^2 + 4}{z - 2i}$$

$$= \frac{i^2 + 4}{i - 2i} = \frac{-1 + 4}{-i} = -\frac{3}{i} \times \frac{i}{i} = 3i$$

$\therefore \lim_{z \rightarrow i} f(z)$  exists and value is  $3i$ .

b)  $\lim_{z \rightarrow 2i} f(z) = \lim_{z \rightarrow 2i} \frac{z^2 + 4}{z - 2i}$

$$= \frac{\cancel{4i^2 + 4}}{\cancel{2i}} \cdot \frac{2i}{1} = \frac{2 \cdot 2i}{1} = 4i$$

but function value  $f(2i) = 3 + 4i$

limit  $\neq$  function

$\therefore$  Not continuous at  $z = 2i$ .

c)  $f(z) = \begin{cases} \frac{z^2 + 4}{z - 2i}, & z \neq 2i \\ 3 + 4i, & z = 2i \end{cases}$

continuous at all points other than  
~~2i~~ because where the denominator is non zero.

We considering only  $z \neq 2i$ , the denominator  
 $z - 2i \neq 0$ . So, function is continuous.


$$15.1) e^z = 1 + \sqrt{3}i$$

consider  $e^z = 1 + \sqrt{3}i$  (Given)

$1 + \sqrt{3}i$  is a complex number

$$r = \sqrt{1^2 + (\sqrt{3})^2} = 2$$

$$\theta = \tan^{-1}\left(\frac{\sqrt{3}}{1}\right) = \frac{\pi}{3}$$



$$1 + \sqrt{3}i = re^{i(\theta + 2n\pi)} \quad n > 0$$

$$(r = 2, \theta = \frac{\pi}{3})$$

$$e^z = 2e^{(2n + \frac{1}{3})\pi i}$$

$$\Rightarrow e^z = 2e^{(2n + \frac{1}{3})\pi i}$$

$$\Rightarrow \ln e^z = \ln [2e^{(2n + \frac{1}{3})\pi i}] \quad [\because \text{take } \ln \text{ on both sides}]$$

$$\Rightarrow z = \ln 2 + \ln e^{(2n + \frac{1}{3})\pi i}$$

$$\therefore z = \ln 2 + (2n + \frac{1}{3})\pi i$$

$$16.1) \lim_{z \rightarrow 2i} \frac{z^2 + 4}{2z^2 + 3z - 4iz - 6i}$$

$$= \lim_{z \rightarrow 2i} \frac{2z}{4z + 3 - 4i}$$

$$= \frac{4i}{8i + 3 - 4i}$$

$$= \frac{4i}{3 + 4i}$$

$$= \frac{4i(3 - 4i)}{9 - 16i^2}$$

$$= \frac{12i - 16i^2}{9 + 16} = \frac{16 + 12i}{25}$$

$$= \frac{16}{25} + \frac{12}{25}i$$

$$\begin{aligned}
 \text{II) } & \lim_{z \rightarrow 0} \frac{z - \sin z}{z^3} \\
 &= \lim_{z \rightarrow 0} \frac{1 - \cos z}{3z^2} \\
 &= \lim_{z \rightarrow 0} \frac{\sin z}{6z} \\
 &\stackrel{H}{=} \lim_{z \rightarrow 0} \frac{\cos z}{6} = \frac{1}{6} \quad \lim_{z \rightarrow 0} \frac{\sin z}{z} \quad [\because \lim_{z \rightarrow 0} \frac{\sin z}{z} = 1]
 \end{aligned}$$

$$\begin{aligned}
 17. \text{ II) } & \left( \frac{1 + \sqrt{3}i}{1 - \sqrt{3}i} \right)^2 \\
 &= \frac{1 + 2\sqrt{3}i + 3i^2}{1 - 2\sqrt{3}i + 3i^2} \\
 &= \frac{-2 + 2\sqrt{3}i}{-2 - 2\sqrt{3}i} \\
 &= \frac{-(1 - \sqrt{3}i)}{-(1 + \sqrt{3}i)} \\
 &= \frac{(1 - \sqrt{3}i)(1 - \sqrt{3}i)}{1 - 3i^2} \\
 &= \frac{1 + 2\sqrt{3}i + 3i^2}{4} \\
 &= \frac{-2 - 2\sqrt{3}i}{4} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i
 \end{aligned}$$

$$z = -\frac{1}{2} - \frac{\sqrt{3}}{2} i$$

modulus:

$$|z| = \sqrt{(-\frac{1}{2})^2 + (\frac{-\sqrt{3}}{2})^2}$$

$$= \sqrt{\frac{1}{4} + \frac{3}{4}}$$

$$= 1$$

Argument:

$$\theta = \pi + \tan^{-1} \left| \frac{-\frac{\sqrt{3}}{2}}{-\frac{1}{2}} \right|$$


$$= \pi + \frac{\pi}{3}$$

$$= 2\pi - \frac{2\pi}{3}$$

$$= \frac{5\pi}{3}$$

$$= -\frac{3\pi + \pi}{3}$$

$$= -\frac{2\pi}{3}$$



$$18. f(2) = k \cdot 2^2 + b = 4k + b$$

$$\begin{aligned}\lim_{z \rightarrow 2} f(z) &= \lim_{z \rightarrow 2} \frac{z^2 - 4}{z^2 - 3z + 2} \\&= \lim_{z \rightarrow 2} \frac{2z}{2z - 3} \\&= \frac{2 \cdot 2}{2 \cdot 2 - 3} = 4\end{aligned}$$

For continuity,

Function value = limit

$$\Rightarrow 4k + b = 4$$

$$\Rightarrow 4k = -2$$

$$\Rightarrow k = -\frac{1}{2}$$