

# MAT215: Complex Variables & Laplace Transformations

## Chapter : Laplace Transformations

Laplace Transformation introduction

Laplace transform of elementary functions



Inspiring Excellence

BRAC University  
Department of Mathematics  
Summer 2025

Instructor: Kazi Hafizur Rahman

# Laplace transformation

## Functional transformation

input function  $f(x) = x^2$   $\xrightarrow{d/dx} 2x$  Functional transform of  $f(x)$

$f(x) = x^2 \xrightarrow{\int dx} \frac{x^3}{3} + c$  (Functional Transformation)

## Integral transformation - (A special functional transformation)

$f(x) \rightarrow$  input function defined on  $[a, b]$

$F(y) = \int_a^b K(x, y) f(x) dx$  Kernel of the integral transformation

$f(x) \xrightarrow{\text{input}} \int_a^b K(x, y) f(x) dx \xrightarrow{\text{integral transformation}} F(y) \rightarrow$  Integral transform of  $f(x)$

function of time domain

$$f(t); t \geq 0$$

# Integral transformation (for a function of time domain)

for a function  $f(t)$  defined on  $t \geq 0$ , an integral transformation has the general form

$$F(\lambda) = \int_0^{\infty} K(\lambda, t) f(t) dt$$

↳ input function  
Kernel  
→ Integral transform of  $f(t)$

Laplace transformation →

A special integral transformation where  
 $K(\lambda, t) = e^{-\lambda t}$

$f(t) \rightarrow$  function defined on  $t \geq 0$ . Then

$$F(\lambda) = \int_0^{\infty} e^{-\lambda t} f(t) dt$$

↳ input  
Kernel of the transformation  
→ Laplace transform of  $f(t)$  provided  
that the integral converges

Notation:

$$F(\lambda) = \int_0^{\infty} e^{-\lambda t} f(t) dt$$

$\mathcal{L}\{f(t)\} = F(\lambda) \rightarrow$  Laplace transform of  $f(t)$  is  $F(\lambda)$

(i)


$$\int uv dx = u \int v dx - \int (u' \int v dx) dx \text{ (integration by parts formula)}$$

choosing  $u, v$  : LIATE formula

## (ii) Shortcut for $u-v$ integration (Tabular method)


$$I = \int_a^b x^3 e^x dx$$

$$= [x^3 e^x - 3x^2 e^x + 6x e^x - 6 e^x]_a^b$$



(iii)  $\lim_{x \rightarrow \infty} e^{mx}$  is finite  
if  $m < 0$

In fact  $\lim_{x \rightarrow \infty} e^{mx} = 0$   
if  $m < 0$



## Laplace transformation of some elementary functions

### Problem

Find  $\mathcal{L}\{f(t)\}$  where  $f(t) = 1$ .

Solution:  $\mathcal{L}\{f(t)\} = \int_0^\infty e^{-st} f(t) dt$

$= \int_0^\infty e^{-st} dt \rightarrow$  (Improper integral of 1st kind)

$$= \lim_{P \rightarrow \infty} \int_0^P e^{-st} dt$$

$$= \lim_{P \rightarrow \infty} \left[ \frac{e^{-st}}{-s} \right]_0^P$$

For the improper integral to converge we must have

$$= -\frac{1}{s} \lim_{P \rightarrow \infty} (e^{-sP} - 1)$$


$$\begin{aligned} & \lim_{P \rightarrow \infty} e^{-sP} \text{ to be finite} \\ & \sim -s < 0 \text{ or } s > 0 \end{aligned}$$

$$= -\frac{1}{s} \left( \lim_{P \rightarrow \infty} e^{-sP} - 1 \right)$$

$$= -\frac{1}{s} (0 - 1) \quad \text{when } s > 0$$

$$= \frac{1}{s} \quad \text{when } s > 0$$

$$\mathcal{L}\{f(t)\} = \frac{1}{s} \quad \text{when } s > 0$$



### Problem

Find  $\mathcal{L}\{t\}$ .

**Solution:**

$$\mathcal{L}\{f(t)\} = \int_0^\infty e^{-st} f(t) dt$$

$$= \int_0^\infty e^{-st} t dt$$

$$= \lim_{P \rightarrow \infty} \int_0^P e^{-st} t dt$$

$$= \lim_{P \rightarrow \infty} \frac{t e^{-st}}{-s} - \frac{e^{-st}}{s^2} \Big|_0^P$$

$$= - \lim_{P \rightarrow \infty} \frac{t e^{-st}}{s} + \frac{e^{-st}}{s^2} \Big|_0^P$$

$$= - \lim_{P \rightarrow \infty} \left\{ \frac{P e^{-sP}}{s} + \frac{e^{-sP}}{s^2} - \frac{1}{s^2} \right\}$$

$$= - \left( 0 + 0 - \frac{1}{s^2} \right) \text{ when } s > 0$$

$$= \frac{1}{s^2} \text{ when } s > 0$$

$$\mathcal{L}\{t^2\} = \frac{1}{s^3} \text{ when } s > 0$$

### Problem

Find  $\mathcal{L}\{t^2\}$ .


$$\text{Ans: } \mathcal{L}\{t^2\} = \frac{2}{s^3}; s > 0$$

### Problem

Find  $\mathcal{L}\{t^3\}$ .

$$\text{Ans: } \mathcal{L}\{t^3\} = \frac{6}{s^4}; s > 0$$

$$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}; s > 0 \quad \text{where } n \text{ is a positive integer}$$



## Problem

Find  $\mathcal{L}\{t^n\}$  where n is a positive integer.

**Solution:**

$$\mathcal{L}\{t^n\} = \int_0^\infty e^{-st} t^n dt$$

$\int uv dx = u \int v dx - \int (u' \int v dx) dx$  (integration by parts formula)  
choosing  $u, v$ : LIATE formula

$$= \lim_{P \rightarrow \infty} \int_0^P e^{-st} t^n dt$$

$$u = t^n, v = e^{-st}$$

$$= \lim_{P \rightarrow \infty} \left\{ \left[ t^n \frac{e^{-st}}{-s} \right]_0^P - \int_0^P n t^{n-1} \frac{e^{-st}}{-s} dt \right\}$$



$$= \lim_{P \rightarrow \infty} \left\{ \frac{P^n}{-s e^{sP}} + \frac{n}{s} \int_0^P e^{-st} t^{n-1} dt \right\}$$

$$= \lim_{P \rightarrow \infty} \frac{P^n}{-s e^{sP}} + \frac{n}{s} \lim_{P \rightarrow \infty} \int_0^P e^{-st} t^{n-1} dt$$

$$= I + \frac{n}{s} \int_0^\infty e^{-st} t^{n-1} dt$$

$$= I + \frac{n}{s} \mathcal{L}\{t^{n-1}\} \quad \text{--- (i)}$$

**computation of I :**

$$I = \lim_{P \rightarrow \infty} \frac{P^n}{-s e^{sP}} \quad \frac{\infty}{\infty} \text{ form} \rightarrow \text{L'hospital Rule}$$

$$= \lim_{P \rightarrow \infty} \frac{n P^{n-1}}{-s^2 e^{sP}}$$

$$= \lim_{P \rightarrow \infty} \frac{n(n-1) P^{n-2}}{-s^3 e^{sP}}$$

⋮

$$= \lim_{P \rightarrow \infty} \frac{n(n-1) \dots 1 P^{n-n}}{-s^{n+1} e^{sP}}$$

$$= \lim_{P \rightarrow \infty} \frac{n!}{-s^{n+1} e^{sP}}$$

$$= \frac{n!}{-s^{n+1}} \lim_{P \rightarrow \infty} e^{-sP} = \frac{n!}{-s^{n+1}} \cdot 0 \text{ if } s > 0 = 0 \text{ if } s > 0$$

Substitute this value in (i) ⇒

$$\mathcal{L}\{t^n\} = \frac{n}{s} \mathcal{L}\{t^{n-1}\}$$

$$\therefore \mathcal{L}\{t^{n-1}\} = \frac{n-1}{s} \mathcal{L}\{t^{n-2}\}$$

$$\therefore \mathcal{L}\{t^n\} = \frac{n}{s} \mathcal{L}\{t^{n-1}\}$$

$$= \frac{n}{s} \cdot \frac{n-1}{s} \mathcal{L}\{t^{n-2}\}$$

$$= \frac{n}{s} \cdot \frac{n-1}{s} \cdot \frac{n-2}{s} \mathcal{L}\{t^{n-3}\}$$

⋮  
⋮  
⋮

$$= \frac{n(n-1) \cdots 1}{s^n} \mathcal{L}\{t^{n-n}\}$$

$$= \frac{n!}{s^n} \mathcal{L}\{1\}$$

$$= \frac{n!}{s^{n+1}} \text{ if } s > 0$$



$$\therefore \mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}} \text{ if } s > 0 \quad \text{where } n \in \mathbb{Z}^+$$

Gamma function:

$$\Gamma(n) = \int_0^\infty e^{-t} t^{n-1} dt$$

$$(i) \quad \Gamma(n) = (n-1)! \text{ if } n \in \mathbb{Z}^+$$

$$\therefore \mathcal{L}\{t^n\} = \frac{\Gamma(n+1)}{s^{n+1}} \text{ if } s > 0 \quad \text{where } n \in \mathbb{Z}^+$$

Question: What if  $n$  is not an integer

$$\therefore \mathcal{L}\{t^n\} = \frac{\Gamma(n+1)}{s^{n+1}} \text{ if } s > 0$$

where  $n$  is not an integer.

**Hint:** Utilize gamma function defn

## Problem

Find  $\mathcal{L}\{e^{at}\}$ .

**Solution:**

$$\mathcal{L}\{f(t)\} = \int_0^\infty e^{-st} f(t) dt$$

$$\mathcal{L}\{e^{at}\} = \int_0^\infty e^{-st} e^{at} dt$$

$$= \lim_{p \rightarrow \infty} \int_0^p e^{-(s-a)t} dt$$

$$= \lim_{p \rightarrow \infty} \frac{e^{-(s-a)t}}{-(s-a)} \Big|_0^p$$

$$= -\frac{1}{s-a} \lim_{p \rightarrow \infty} \left\{ e^{-(s-a)p} - 1 \right\}$$

$$= -\frac{1}{s-a} \left\{ \lim_{p \rightarrow \infty} e^{-(s-a)p} - 1 \right\}$$

$$= -\frac{1}{s-a} (0-1) \quad \text{when } s-a > 0 \text{ or } s > a$$

$$= \frac{1}{s-a} \quad \text{when } s > a$$

$$\mathcal{L}\{e^{at}\} = \frac{1}{s-a} \quad \text{when } s > a$$



H.W.

## Problem

Find (i)  $\mathcal{L}\{\sin at\}$  and (ii)  $\mathcal{L}\{\cos at\}$ .

**Solution:**

$$\mathcal{L}\{f(t)\} = \int_0^\infty e^{-st} f(t) dt$$

$$\mathcal{L}\{\sin at\} = \int_0^\infty e^{-st} \sin at dt$$

$$\mathcal{L}\{\cos at\} = \int_0^\infty e^{-st} \cos at dt$$



$$\begin{aligned}
d \{ e^{iat} \} &= \int_0^\infty e^{-st} e^{iat} dt \\
&= \lim_{P \rightarrow \infty} \int_0^P e^{-(s-ai)t} dt \\
&= \lim_{P \rightarrow \infty} \left[ \frac{e^{-(s-ai)t}}{-(s-ai)} \right]_0^P \\
&= -\frac{1}{s-ai} \lim_{P \rightarrow \infty} \{ e^{-(s-ai)P} - 1 \} \\
&= -\frac{1}{s-ai} \lim_{P \rightarrow \infty} [e^{-sP} \{ \cos(ap) + i \sin(ap) \} - 1] \\
&= -\frac{1}{s-ai} \left[ \lim_{P \rightarrow \infty} e^{-sP} \{ \cos(ap) + i \sin(ap) \} - 1 \right] \\
&= -\frac{1}{s-ai} (0-1) \text{ if } s>0 \\
&= \frac{1}{s-ai} \text{ if } s>0 \\
&= \frac{s+ai}{(s-ai)(s+ai)} = \frac{s+ai}{s^2+a^2} = \frac{s}{s^2+a^2} + i \frac{a}{s^2+a^2}
\end{aligned}$$

So we have,

$$\int_0^\infty e^{-st} e^{iat} dt = \frac{s}{s^2+a^2} + i \frac{a}{s^2+a^2}$$

$$\Rightarrow \int_0^\infty e^{-st} \{ \cos at + i \sin at \} dt = \frac{s}{s^2+a^2} + i \frac{a}{s^2+a^2}$$

$$\Rightarrow \int_0^\infty e^{-st} \cos at dt + i \int_0^\infty e^{-st} \sin at dt = \frac{s}{s^2+a^2} + i \frac{a}{s^2+a^2}$$

$$\int_0^\infty e^{-st} \cos at dt = \frac{s}{s^2+a^2}; s>0 \quad \& \quad \int_0^\infty e^{-st} \sin at dt = \frac{a}{s^2+a^2}; s>0$$

$$\Rightarrow d \{ \cos at \} = \frac{s}{s^2+a^2} ; s>0 \quad \&$$

$$d \{ \sin at \} = \frac{a}{s^2+a^2} ; s>0$$

# MAT215: Complex Variables & Laplace Transformations

## Chapter : Laplace Transformations

Linearity of Laplace transformation

1st Translation theorem

Multiplication by  $t^n$



Inspiring Excellence

BRAC University  
Department of Mathematics  
Summer 2025

Instructor: Kazi Hafizur Rahman

## Linearity of Laplace transformation

(i)  $\mathcal{L}\{f(t) + g(t)\} = \mathcal{L}\{f(t)\} + \mathcal{L}\{g(t)\}$

(ii)  $\mathcal{L}\{\alpha f(t)\} = \alpha \mathcal{L}\{f(t)\}$



**Reason:**

$$\begin{aligned} \text{(i)} \quad \mathcal{L}\{f(t) + g(t)\} &= \int_0^\infty e^{-st} \{f(t) + g(t)\} dt \\ &= \int_0^\infty e^{-st} f(t) dt + \int_0^\infty e^{-st} g(t) dt \\ &= \mathcal{L}\{f(t)\} + \mathcal{L}\{g(t)\} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \mathcal{L}\{\alpha f(t)\} &= \int_0^\infty e^{-st} (\alpha f(t)) dt \\ &= \alpha \int_0^\infty e^{-st} f(t) dt = \alpha \mathcal{L}\{f(t)\} \end{aligned}$$

## Linearity of Laplace transformation (Compact form)

$$\mathcal{L}\{\alpha f(t) + \beta g(t)\} = \alpha \mathcal{L}\{f(t)\} + \beta \mathcal{L}\{g(t)\}$$


### Problem

Find (i)  $\mathcal{L}\{\sinh at\}$  and (ii)  $\mathcal{L}\{\cosh at\}$ .

**Solution:**  $\sinh(at) = \frac{e^{at} - e^{-at}}{2}$


$$\begin{aligned} \mathcal{L}\{\sinh(at)\} &= \mathcal{L}\left\{\frac{e^{at} - e^{-at}}{2}\right\} \\ &= \frac{1}{2} (\mathcal{L}\{e^{at}\} - \mathcal{L}\{e^{-at}\}) \\ &\quad \text{with } s > a \quad \text{and } s > -a \\ &= \frac{1}{2} \left( \frac{1}{s-a} - \frac{1}{s+a} \right); \quad s > |a| \end{aligned}$$

$$a = 2$$



$$s > 2 \quad (s > a)$$

$$a = -2$$



$$s > 2 \quad (s > -a)$$

$$s > |a|$$

Therefore we have,

$$\mathcal{L}\{\sinh(at)\} = \frac{1}{2} \left( \frac{1}{s-a} - \frac{1}{s+a} \right); \quad s > |a|$$

$$= \frac{1}{2} \frac{s+a - s-a}{s^2 - a^2}; \quad s > |a|$$

$$\mathcal{L}\{\sinh at\} = \frac{a}{s^2 - a^2}; \quad s > |a| \quad (\text{Ans})$$

$$\mathcal{L}\{\cosh at\} = \frac{s}{s^2 - a^2}; \quad s > |a| \quad (\text{Ans})$$

### Laplace Transforms of Elementary Functions (Real $s$ )

Function $f(t)$	$\mathcal{L}\{f(t)\} = F(s)$	Function $f(t)$	$\mathcal{L}\{f(t)\} = F(s)$
1	$\frac{1}{s}, \quad s > 0$	$\sin(at)$	$\frac{a}{s^2 + a^2}, \quad s > 0$
$t^n, n \in \mathbb{Z}^+$	$\frac{n!}{s^{n+1}}, \quad s > 0$	$\cos(at)$	$\frac{s}{s^2 + a^2}, \quad s > 0$
$t^n, n \notin \mathbb{Z}$	$\frac{\Gamma(n+1)}{s^{n+1}}, \quad s > 0$	$\sinh(at)$	$\frac{a}{s^2 - a^2}, \quad s >  a $
$e^{at}$	$\frac{1}{s-a}, \quad s > a$	$\cosh(at)$	$\frac{s}{s^2 - a^2}, \quad s >  a $



## Problem

Find:

(i)  $\mathcal{L}\{3e^{-2t}\}$     (ii)  $\mathcal{L}\{4t^3 - e^{-t}\}$     (iii)  $\mathcal{L}\{(t^2 + 1)^2\}$

(iv)  $\mathcal{L}\{7 \sin 2t - 3 \cos 4t\}$      ~~$\mathcal{L}\{(4e^{2t} - 2)^3\}$~~

**Solution:** (iii)  $\mathcal{L}\{(t^2 + 1)^2\} = \mathcal{L}\{t^4 + 2t^2 + 1\}$

$$= \mathcal{L}\{t^4\} + 2\mathcal{L}\{t^2\} + \mathcal{L}\{1\}$$

$$= \frac{4!}{s^5} + 2 \cdot \frac{2!}{s^3} + \frac{1}{s}; s > 0$$

(iv)  $\mathcal{L}\{7 \sin 2t - 3 \cos 4t\} = 7\mathcal{L}\{\sin 2t\} - 3\mathcal{L}\{\cos 4t\}$

$$= 7 \cdot \frac{2}{s^2 + 4} - 3 \cdot \frac{s}{s^2 + 16}; s > 0$$

(Ans.)

(ii)  $\mathcal{L}\{4t^3 - e^{-t}\} = 4\mathcal{L}\{t^3\} - \mathcal{L}\{e^{-t}\}$

$$\downarrow s > 0 \quad \downarrow s > -1$$

$$= 4 \cdot \frac{3!}{s^4} - \frac{1}{s+1}; s > 0 \quad (\text{Ans.})$$

## 1st Translation theorem

$$\mathcal{L}\{f(t)\} = F(s) \Rightarrow \mathcal{L}\{e^{at}f(t)\} = F(s-a) \text{ for any } a \in \mathbb{R}$$

$$\mathcal{L}\{e^{at}f(t)\} = F(s)|_{s \rightarrow s-a} = \mathcal{L}\{f(t)\}|_{s \rightarrow s-a}$$

## 1st Translation theorem

$$\mathcal{L}\{e^{at}f(t)\} = \mathcal{L}\{f(t)\}|_{s \rightarrow s-a} \text{ for any real number } a.$$



## Problem

Find: (i)  $\mathcal{L}\{e^{-3t}t^3\}$  (ii)  $\mathcal{L}\{5e^{3t}\sin 4t\}$

## Solution:

$$\begin{aligned}
 \text{(i)} \quad \mathcal{L}\{e^{-3t}t^3\} &= \alpha \{t^3\}|_{s \rightarrow s+3} \\
 &= \frac{3!}{s^4} |_{s \rightarrow s+3}; \quad s > 0 |_{s \rightarrow s+3} \\
 &= \frac{3!}{(s+3)^4}; \quad s+3 > 0 \quad (\text{Ans:})
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii)} \quad \mathcal{L}\{5e^{3t}\sin 4t\} &= 5 \alpha \{e^{3t}\sin 4t\} \\
 &= 5 \alpha \{\sin 4t\}|_{s \rightarrow s-3} \\
 &= 5 \frac{4}{s^2 + 16} |_{s \rightarrow s-3}; \quad s > 0 |_{s \rightarrow s-3} \\
 &= \frac{20}{(s-4)^2 + 16}; \quad s > 4 \quad (\text{Ans:})
 \end{aligned}$$

## Problem

Find:

(i)  $\mathcal{L}\{(t+2)^2 e^t\}$  (ii)  $\mathcal{L}\{e^{-t}(3 \sinh 2t - 5 \cosh 3t)\}$

(iii)  $\mathcal{L}\{e^{-4t} \cosh 2t\}$  (iv)  $\mathcal{L}\{e^{2t}(3 \sin 4t - 4 \cos 3t)\}$

## Solution:

$$\begin{aligned}
 \text{(iv)} \quad \mathcal{L}\{e^{2t}(3 \sin 4t - 4 \cos 3t)\} &= \alpha \{3 \sin 4t - 4 \cos 3t\}|_{s \rightarrow s-2} \\
 &= 3 \alpha \{\sin 4t\} - 4 \alpha \{\cos 3t\}|_{s \rightarrow s-2} \\
 &\quad s > 0 \downarrow \quad \downarrow s > 0
 \end{aligned}$$

$$= 3 \frac{4}{\lambda^2 + 16} - 4 \frac{\lambda}{\lambda^2 + 9} \Big|_{\lambda \rightarrow \lambda-2} ; \quad \lambda > 0 \Big|_{\lambda \rightarrow \lambda-2}$$

$$= \frac{12}{(\lambda-2)^2 + 16} - \frac{4(\lambda-2)}{(\lambda-2)^2 + 9} ; \quad \lambda-2 > 0 \quad (\text{A.M.})$$

Laplace transformation of the form  $t^n f(t)$

If  $\mathcal{L}\{f(t)\} = F(s)$ , then  $\mathcal{L}\{t^n f(t)\} = ?$

**Derivation:**

$$\mathcal{L}\{f(t)\} = F(s)$$



$$F(s) = \int_0^\infty e^{-st} f(t) dt$$

$$\Rightarrow \frac{d}{ds}(F(s)) = \frac{d}{ds} \left( \int_0^\infty e^{-st} f(t) dt \right)$$

Assume that order of differentiation  
& integration can be interchanged

$$= \int_0^\infty \frac{d}{ds} (e^{-st} f(t)) dt$$

$$\frac{d}{ds}(F(s)) = \int_0^\infty f(t) e^{-st} (-t) dt$$

$$\Rightarrow - \frac{d}{ds}(F(s)) = \int_0^\infty e^{-st} (t f(t)) dt$$

$$\mathcal{L}\{t f(t)\} = - \frac{d}{ds} (\mathcal{L}\{f(t)\})$$



$$\mathcal{L}\{t^n f(t)\} = \mathcal{L}\{t \underbrace{\frac{(t f(t))}{g(t)}}\}$$

$$= - \frac{d}{ds} (\alpha \{ f(t) \})$$

$$= - \frac{d}{ds} (- \frac{d}{ds} (\alpha \{ f(t) \}))$$

$$\alpha \{ t^v f(t) \} = (-1)^v \frac{d^v}{ds^v} (\alpha \{ f(t) \})$$

$$\alpha \{ t^n f(t) \} = (-1)^n \frac{d^n}{ds^n} (\alpha \{ f(t) \})$$

Laplace transformation of the form  $t^n f(t)$

$$\mathcal{L}\{t^n f(t)\} = (-1)^n \frac{d^n}{ds^n} [\mathcal{L}\{f(t)\}]$$



### Problem

Find: (i)  $\mathcal{L}\{t \sin t\}$     (ii)  $\mathcal{L}\{t^2 e^{2t}\}$     (iii)  $\mathcal{L}\{te^{-3t} \sin t\}$

### Solution:

$$\begin{aligned}
 \text{(i)} \quad \mathcal{L}\{t \sin t\} &= - \frac{d}{ds} (\alpha \{\sin t\}) \\
 &= - \frac{d}{ds} \left( \frac{1}{s^2 + 1} \right) ; s > 0 \\
 &= - \frac{(s^2 + 1) \cdot 0 - 1 \cdot 2s}{(s^2 + 1)^2} \\
 &= \frac{2s}{(s^2 + 1)^2} ; s > 0 \quad (\text{Ans.})
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii)} \quad \mathcal{L}\{t^2 e^{2t}\} &= (-1)^2 \frac{d^2}{ds^2} (\alpha \{ e^{2t} \}) \\
 &= \frac{d^2}{ds^2} \left( \frac{1}{s-2} \right) ; s > 2
 \end{aligned}$$

$$= (-1) (-2) (\lambda - 2)^{-3}; \quad \lambda > 2$$

$$= \frac{2}{(\lambda - 2)^3}; \quad \lambda > 2$$

$$\begin{aligned} \text{(iii)} \mathcal{L}\{te^{-3t} \sin t\} &= \mathcal{L}\{e^{-3t} t \sin t\} \\ &= \mathcal{L}\{t \sin t\} \Big|_{\lambda \rightarrow \lambda + 3} \\ &= \frac{2\lambda}{(\lambda^2 + 1)^2} \Big|_{\lambda \rightarrow \lambda + 3} : \quad \lambda > 0 \Big|_{\lambda \rightarrow \lambda + 3} \\ &= \frac{2(\lambda + 3)}{\{(\lambda + 3)^2 + 1\}^2}; \quad \lambda > -3 \quad (\text{Ans}) \end{aligned}$$

~~H.W.~~

### Brain Teaser

Find  $\mathcal{L}\{te^{-3t} \cos t \sin t\}$ .

# MAT215: Complex Variables & Laplace Transformations

## Chapter : Laplace Transformations

Unit Step function

Piecewise function representation in terms of Unit Step function

2nd Translation Theorem



Inspiring Excellence

BRAC University  
Department of Mathematics  
Summer 2025

Instructor: Kazi Hafizur Rahman

## Unit Step Function

$$u(t) = \begin{cases} 1 & \text{if } t \geq 0 \\ 0 & \text{if } t < 0 \end{cases}$$

$a$  is any positive real number

$$u(t-a) = \begin{cases} 1 & \text{if } t-a \geq 0 \\ 0 & \text{if } t-a < 0 \end{cases}$$




$$u(t-a) = \begin{cases} 1 & \text{if } t \geq a \\ 0 & \text{if } t < a \end{cases}$$

↓

restrict this function in non-negative Domain

$$u(t-a) = \begin{cases} 1 & \text{if } t \geq a \\ 0 & \text{if } 0 \leq t < a \end{cases}$$

↓  
(Unit step function at  $a$ )




### Problem




Plot the functions: (i)  $u(t - \frac{\pi}{2})$     (ii)  $1 - u(t - \frac{\pi}{2})$

### Solution:

(i)  $u(t - \frac{\pi}{2})$



$$(ii) 1 - u(t - \frac{\pi}{2})$$



## 2 Geometric Properties of Unit Step Function

(i) Multiplication by  $u(t-a)$ :


$$f(t) = \sin t$$

$$g(t) = u(t - \pi/2)$$

$$h(t) = \sin t \cdot u(t - \pi/2)$$

$$\sin t \cdot u(t - \pi/2) = \begin{cases} \sin t; & t \geq \pi/2 \\ 0 & ; 0 \leq t < \pi/2 \end{cases}$$

$$f(t) \cdot u(t-a) = \begin{cases} f(t); & t \geq a \\ 0 & ; 0 \leq t < a \end{cases}$$



(i) right portion to  $a$  (including  $a$ )  $\rightarrow$  as it is

(ii) left portion to  $a$  (excluding  $a$ )  $\rightarrow$  nullifies

## (ii) Multiplication by $1 - u(t-a)$ :

$$f(t) = \sin t$$

$$g(t) = 1 - u(t - \pi/2)$$

$$h(t) = \sin t \{1 - u(t - \pi/2)\}$$

$$\sin t \{1 - u(t - \pi/2)\}$$

$$= \begin{cases} \sin t ; & 0 \leq t < \pi/2 \\ 0 ; & t \geq \pi/2 \end{cases}$$


$$f(t) \{1 - u(t-a)\}$$

$$= \begin{cases} f(t) ; & 0 \leq t < a \\ 0 ; & t \geq a \end{cases}$$



(i) left portion to  $a$  (excluding  $a$ )  $\rightarrow$  as it is


(ii) right portion to  $a$  (including  $a$ )  $\rightarrow$  nullifies





# Piecewise Function in Terms of Unit Step

## Function

$$h(t) = \begin{cases} f(t); & 0 \leq t < a \\ g(t); & t \geq a \end{cases}$$



$$h(t) = f(t) (1 - u(t-a)) + g(t) u(t-a)$$



## Piecewise function representation through unit step function

Suppose that,

$$h(t) = \begin{cases} f(t); & 0 \leq t < a \\ g(t); & t \geq a \end{cases}$$



Then,

$$h(t) = f(t)(1 - u(t-a)) + g(t)u(t-a)$$

## 2nd Translation Theorem

$$\mathcal{L}\{f(t)\} = F(s) \Rightarrow \mathcal{L}\{f(t-a)u(t-a)\} = F(s)e^{-as}$$

for any positive value of a

$$\mathcal{L}\{f(t-a)u(t-a)\} = \mathcal{L}\{f(t)\}e^{-as}$$

## 2nd Translation Theorem

$$\mathcal{L}\{f(t-a)u(t-a)\} = \mathcal{L}\{f(t)\}e^{-as} \text{ for any positive value of a.}$$

### Problem

Find:

(i)  $\mathcal{L}\{(t^2 + 1)u(t-1)\}$     (ii)  $\mathcal{L}\{(t-1)u(t-1)\}$

(iii)  $\mathcal{L}\{e^{-2t}u(t-1)\}$     (iv)  $\mathcal{L}\{\cos 2tu(t-\pi)\}$

(v)  $\mathcal{L}\{u(t-1)(t^2 + 1)\}$

### Solution:

$$(i) \mathcal{L}\{(t^2 + 1)u(t-1)\} = \mathcal{L}\{(t-1)^2 + 2(t-1)u(t-1)\}$$

$$= \mathcal{L}\{(t-1)^2 + 2(t-1) + 2u(t-1)\}$$

$$f(t) = t^2 + 2t + 2$$

$$= \mathcal{L}\{f(t-1)u(t-1)\}$$

$$= \mathcal{L}\{f(t)\}e^{-s}$$

$$= \mathcal{L}\{t^2 + 2t + 2\}e^{-s}$$

$$= (\mathcal{L}\{t^2\} + 2\mathcal{L}\{t\} + 2\mathcal{L}\{1\})e^{-s}$$

$$= \left(\frac{2}{s^3} + \frac{2}{s^2} + \frac{2}{s}\right)e^{-s}; s > 0 \text{ (A.M.)}$$

$$(iv) \mathcal{L}\{\cos 2tu(t-\pi)\} = \mathcal{L}\{\cos(2(t-\pi) + 2\pi) u(t-\pi)\}$$

$$f(t) = \cos(2t + 2\pi) = \cos(2t)$$

$$= \mathcal{L}\{f(t-\pi) u(t-\pi)\}$$

$$= \mathcal{L}\{f(t)\} e^{-\pi s}$$

$$= \mathcal{L}\{\cos(2t)\} e^{-\pi s}$$

$$= -\frac{s}{s^2 + 4} e^{-\pi s}; \quad s > 0 \quad (\text{Ans.})$$

$$(ii) \mathcal{L}\{(t-1)u(t-1)\} = \mathcal{L}\{f(t-1)u(t-1)\}$$

$$= \mathcal{L}\{f(t)\} e^{-s} \quad f(t) = t$$

$$= \mathcal{L}\{t\} e^{-s}$$

$$= \frac{1}{s^2} e^{-s}; \quad s > 0 \quad (\text{Ans.})$$

$$(iii) \mathcal{L}\{e^{-2t}u(t-1)\} = \mathcal{L}\{e^{-2(t-1)-2} u(t-1)\}$$

$$f(t) = e^{-2t-2}$$

$$= \mathcal{L}\{f(t-1)u(t-1)\}$$

$$= \mathcal{L}\{f(t)\} e^{-s}$$

$$= \mathcal{L}\{e^{-2t} e^{-2}\} e^{-s}$$

$$= e^{-2} \cancel{\{e^{-2t}\}} e^{-s}$$

$$= e^{-2-s} \frac{1}{s+2}; \quad s>-2 \quad (\text{Ans.})$$

## Problem

Find  $\mathcal{L}\{f(t)\}$  where

$$(i) f(t) = \begin{cases} 2t+1; & 0 \leq t < 2 \\ 3t; & t \geq 2 \end{cases} \quad (ii) f(t) = \begin{cases} 0; & 0 \leq t < \pi \\ \sin t; & t \geq \pi \end{cases}$$

$$(iii) f(t) = \begin{cases} \sin t; & 0 \leq t < 2\pi \\ 0; & t \geq 2\pi \end{cases} \quad (iv) f(t) = \begin{cases} 0; & 0 \leq t < \pi \\ 3 \cos t; & t \geq \pi \end{cases}$$

## Solution:

$$(i) f(t) = \begin{cases} 2t+1; & 0 \leq t < 2 \\ 3t; & t \geq 2 \end{cases}$$

$$f(t) = (2t+1)(1 - u(t-2)) + 3t u(t-2)$$

$$= 2t+1 - (2t+1)u(t-2) + 3t u(t-2)$$

$$= 2t+1 + u(t-2)(3t-2t-1)$$

$$f(t) = 2t+1 + (t-1)u(t-2)$$

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{2t+1 + (t-1)u(t-2)\}$$

$$= 2\mathcal{L}\{t\} + \mathcal{L}\{1\} + \mathcal{L}\{(t-1)u(t-2)\}$$

$$= 2 \frac{1}{\lambda \gamma} + \frac{1}{\lambda} + \alpha \{ (t-2+1) u(t-2) \}$$

$$f(t) = t+1$$

$$= \frac{2}{\lambda \gamma} + \frac{1}{\lambda} + \alpha \{ f(t-2) u(t-2) \}$$

$$= \frac{2}{\lambda \gamma} + \frac{1}{\lambda} + \alpha \{ t+1 \} e^{-2\lambda}$$

$$= \frac{2}{\lambda \gamma} + \frac{1}{\lambda} + \left( \frac{1}{\lambda \gamma} + \frac{1}{\lambda} \right) e^{-2\lambda}; \lambda > 0 \text{ (Ans.)}$$

$$(iii) f(t) = \begin{cases} \sin t; & 0 \leq t < 2\pi \\ 0; & t \geq 2\pi \end{cases}$$

$$f(t) = \sin t (1 - u(t-2\pi))$$

$$\alpha \{ f(t) \} = \alpha \{ \sin t (1 - u(t-2\pi)) \}$$

$$= \alpha \{ \sin t - \sin t u(t-2\pi) \}$$

$$= \alpha \{ \sin t \} - \alpha \{ \sin t u(t-2\pi) \}$$

$$= \alpha \{ \sin t \} - \alpha \{ \sin(t-2\pi+2\pi) u(t-2\pi) \}$$

$$f(t) = \sin(t+2\pi) = \sin(t)$$

$$= \alpha \{ \sin t \} - \alpha \{ f(t-2\pi) u(t-2\pi) \}$$

$$= \alpha \{ \sin t \} - \alpha \{ f(t) \} e^{-2\pi\lambda}$$

$$= \alpha \{ \sin t \} - \alpha \{ \sin t \} e^{-2\pi\lambda}$$

$$= \alpha \{ \sin t \} (1 - e^{-2\pi\lambda})$$

$$= \frac{1}{\lambda \gamma + 1} (1 - e^{-2\pi\lambda}); \lambda > 0 \text{ (Ans.)}$$

$$(iv) f(t) = \begin{cases} 0; & 0 \leq t < \pi \\ 3 \cos t; & t \geq \pi \end{cases}$$

$$f(t) = 3 \cos t u(t - \pi)$$

$$g(t) = \cos(t + \pi)$$

$$\alpha \{f(t)\} = \alpha \{3 \cos t u(t - \pi)\}$$

$$= -\text{const}$$

$$= 3 \alpha \{\cos(t - \pi + \pi) u(t - \pi)\}$$

$$= 3 \alpha \{g(t - \pi) u(t - \pi)\}$$

$$= 3 \alpha \{g(t)\} e^{-\pi \lambda}$$

$$= 3 \alpha \{-\text{const}\} e^{-\pi \lambda} = -\frac{3 \lambda}{\lambda^2 + 1} e^{-\pi \lambda}, \lambda > 0 \quad (\text{Ans.})$$

### Brain Teaser

Write the following function in terms of unit step functions.

$$h(t) = \begin{cases} f_1(t), & 0 \leq t < t_1 \\ f_2(t), & t_1 \leq t < t_2 \\ f_3(t), & t \geq t_2 \end{cases}$$

### Brain Teaser

Find the Laplace transform of

$$h(t) = \begin{cases} 0, & 0 \leq t < \pi \\ \sin t, & \pi \leq t < 2\pi \\ 0, & t \geq 2\pi \end{cases}$$

# MAT215: Complex Variables & Laplace Transformations

## Chapter : Laplace Transformations

Partial Fraction Decomposition



Inspiring Excellence

BRAC University  
Department of Mathematics  
Summer 2025

Instructor: Kazi Hafizur Rahman

## Partial Fraction Decomposition

$\frac{P(s)}{Q(s)}$  → polynomial &  $\text{Deg } (Q(s)) > \text{Deg } (P(s))$   
 $\frac{P(s)}{Q(s)}$  → polynomial

$Q(s)$  has distinct linear factors

$$\frac{P(s)}{(a_1s+b_1)(a_2s+b_2)\dots(a_ns+b_n)} = \frac{-1}{\overbrace{\phantom{\frac{A_1}{a_1s+b_1} + \frac{A_2}{a_2s+b_2} + \dots + \frac{A_n}{a_ns+b_n}}}^{\text{---}}}$$

$$= \frac{A_1}{a_1s+b_1} + \frac{A_2}{a_2s+b_2} + \dots + \frac{A_n}{a_ns+b_n}$$

### Problem

Partially decompose  $\frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)}$

### Solution:

$$\frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)} = \frac{A}{s+4} + \frac{B}{s-1} + \frac{C}{s-2}$$

$$\Rightarrow s^2 + 6s + 9 = A(s-1)(s-2) + B(s+4)(s-2) + C(s+4)(s-1)$$

$\downarrow$   
valid for any value of  $s$

$$(i) s = -4 \Rightarrow 16 - 24 + 9 = A(-5)(-6) \Rightarrow A = 1/30$$

$$(ii) s = 1 \Rightarrow 0 = -16/5$$

$$(iii) s = 2 \Rightarrow C = 25/6$$

$$\frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)} = \frac{1/30}{s+4} + \frac{-16/5}{s-1} + \frac{25/6}{s-2} \quad (\text{Ans!})$$

$Q(s)$  has repeated linear factor

$$\frac{P(s)}{(as+b)^n} = \frac{A_1}{as+b} + \frac{A_2}{(as+b)^2} + \dots + \frac{A_n}{(as+b)^n}$$

### Problem

Partially decompose  $\frac{2s+5}{(s-3)^2}$

Solution:

$$\frac{2s+5}{(s-3)^2} = \frac{A}{s-3} + \frac{B}{(s-3)^2}$$

$$\Rightarrow 2s+5 = A(s-3) + B$$

$$(i) s=3 \Rightarrow 11 = B$$

$$(ii) \text{ co-efficient of } s : 2 = A$$

$$\frac{2s+5}{(s-3)^2} = \frac{2}{s-3} + \frac{11}{(s-3)^2} \quad (\text{Ans})$$

$Q(s)$  has irreducible quadratic factor(not repeated)

$$\frac{P(s)}{as^2+bs+c} = \frac{-1 \rightarrow A s + B}{as^2+bs+c}$$

### Problem

Partially decompose  $\frac{6s^2+50}{(s+3)(s^2+4)}$

**Solution:**

$$\frac{6s^2 + 50}{(s+3)(s^2+4)} = \frac{A}{s+3} + \frac{Bs+C}{s^2+4}$$

$$\Rightarrow 6s^2 + 50 = A(s^2 + 4) + (Bs + C)(s + 3)$$

$$(i) s = -3 \Rightarrow 104 = 13A \Rightarrow A = 8$$

$$(ii) s^2 : 6 = A + B \Rightarrow B = -2$$

$$(iii) \text{ constant : } 50 = 4A + 3C \Rightarrow C = 6$$

$$\frac{6s^2 + 50}{(s+3)(s^2+4)} = \frac{8}{s+3} + \frac{-2s+6}{s^2+4} \quad (\text{Ans!})$$

**Problem**

Partially decompose  $\frac{2s+1}{s(s+1)(s^2+4s+6)}$

**Solution:**

$$\frac{2s+1}{s(s+1)(s^2+4s+6)} = \frac{A}{s} + \frac{B}{s+1} + \frac{Cs+D}{s^2+4s+6}$$

$$\Rightarrow 2s+1 = A(s+1)(s^2+4s+6) + Bs(s^2+4s+6) + (Cs+D)s$$

$$(i) s=0 \Rightarrow 1 = A \cdot 6 \Rightarrow A = 1/6$$

$$(ii) s=-1 \Rightarrow -1 = -B(1-4+6) \Rightarrow B = 1/3$$

$$(iii) s^2 : 0 = A + B + C \Rightarrow C = -(A+B) = -1/2$$

$$(iv) \lambda : 2 = 6A + 4A + 6B + D \Rightarrow D = -5/3$$

$$\frac{2s+1}{s(s+1)(s^2+4s+6)} = \frac{1/6}{\lambda} + \frac{1/3}{\lambda+1} + \frac{-1/2\lambda - 5/3}{\lambda^2+4\lambda+6} \quad (\text{Ans})$$

### Problem

Partially decompose  $\frac{3s}{(s+1)(s^2+1)}$

**Solution:**

$$\frac{3s}{(s+1)(s^2+1)} = \frac{A}{s+1} + \frac{Bs+C}{s^2+1}$$

$$\Rightarrow 3s = A(s^2+1) + (Bs+C)(s+1)$$

$$(i) s=-1 \Rightarrow -3 = 2A \Rightarrow A = -3/2$$

$$(ii) s^2: 0 = A + B \Rightarrow B = 3/2$$

$$(iii) \text{constant: } 0 = A + C \Rightarrow C = 3/2$$

$$\frac{3s}{(s+1)(s^2+1)} = \frac{-3/2}{s+1} + \frac{3/2s + 3/2}{s^2+1} \quad (\text{Ans})$$

### Problem

Partially decompose  $\frac{s}{(s^2+4)(s^2+9)}$

**Solution:**

$$\frac{s}{(s^2+4)(s^2+9)} = \frac{As+B}{s^2+4} + \frac{Cs+D}{s^2+9}$$

$$\Rightarrow s = (A\lambda + B)(s^2 + 9) + (C\lambda + D)(s^2 + 4)$$

$$\Rightarrow s = A\lambda^3 + 9A\lambda + B\lambda^2 + 9B + C\lambda^3 + 4C\lambda + D\lambda^2 + 4D$$

$$\Rightarrow s = (A+C)\lambda^3 + (B+D)\lambda^2 + (9A+4C)\lambda + 9B+4D$$

$$\lambda^3: 0 = A + C \quad \text{(i)} \Rightarrow A = -C \Rightarrow A = -1/5$$

$$\lambda^2: 0 = B + D \quad \text{(ii)} \Rightarrow B = -D \Rightarrow B = 0$$

$$\lambda: 1 = 9A + 4C \quad \text{(iii)} \Rightarrow -9C + 4C = 1 \Rightarrow C = -1/5$$

$$\text{constant: } 0 = 9B + 4D \quad \text{(iv)} \Rightarrow -9D + 4D = 0 \Rightarrow D = 0$$

$$\frac{s}{(s^2 + 4)(s^2 + 9)} = \frac{-1/5s}{s^2 + 4} + \frac{1/5s}{s^2 + 9} \quad (\text{Ans:})$$

$Q(s)$  has repeated irreducible quadratic factor

$$\frac{P(s)}{(as^2 + bs + c)^n} = \frac{A_1s + B_1}{as^2 + bs + c} + \frac{A_2s + B_2}{(as^2 + bs + c)^2} + \dots + \frac{A_ns + B_n}{(as^2 + bs + c)^n}$$

### Problem

H.W.

Partially decompose  $\frac{s+3}{(s-1)(s^2+1)^2}$

**Solution:**

$$\frac{s+3}{(s-1)(s^2+1)^2} = \frac{A}{s-1} + \frac{Bs+C}{s^2+1} + \frac{Ds+E}{(s^2+1)^2}$$

# MAT215: Complex Variables & Laplace Transformations

## Chapter : Laplace Transformations

Inverse of Laplace transform meaning  
Elementary inverse Laplace transforms & Linearity of  
Inverse Laplace transform  
1st Translation theorem in inverse form  
Inverse using Partial fraction decomposition



BRAC University  
Department of Mathematics  
Summer 2025

Instructor: Kazi Hafizur Rahman

## Inverse Laplace Transformation

$f(t) \rightarrow$  input function

$$F(s) = \int_0^{\infty} e^{-st} f(t) dt \Leftrightarrow \mathcal{L}\{f(t)\} = F(s) \Leftrightarrow \mathcal{L}^{-1}\{F(s)\} = f(t)$$

$F(s) \rightarrow$  laplace transform of  $f(t)$

$f(t) \rightarrow$  Inverse laplace transform of  $F(s)$

- $\mathcal{L}\{1\} = \frac{1}{s} \Leftrightarrow \mathcal{L}^{-1}\left\{\frac{1}{s}\right\} = 1$

### Inverse Laplace Transforms of some known expressions

$F(s)$	$\mathcal{L}^{-1}\{F(s)\} = f(t)$	$F(s)$	$\mathcal{L}^{-1}\{F(s)\} = f(t)$
$\frac{1}{s}$	1	$\frac{a}{s^2 + a^2}$	$\sin(at)$
$\frac{n!}{s^{n+1}}, n \in \mathbb{Z}^+$	$t^n$	$\frac{s}{s^2 + a^2}$	$\cos(at)$
$\frac{\Gamma(n+1)}{s^{n+1}}, n \notin \mathbb{Z}$	$t^n$	$\frac{a}{s^2 - a^2}$	$\sinh(at)$
$\frac{1}{s-a}$	$e^{at}$	$\frac{s}{s^2 - a^2}$	$\cosh(at)$

### Linearity of Inverse Laplace transformation

$$\mathcal{L}^{-1}\{\alpha F(s) + \beta G(s)\} = \alpha \mathcal{L}^{-1}\{F(s)\} + \beta \mathcal{L}^{-1}\{G(s)\}$$

## Problem

Find:

- (i)  $\mathcal{L}^{-1}\left\{\frac{1}{s^4}\right\}$
- (ii)  $\mathcal{L}^{-1}\left\{\frac{12}{4-3s}\right\}$
- (iii)  $\mathcal{L}^{-1}\left\{\frac{23s-15}{s^2+8}\right\}$
- (iv)  $\mathcal{L}^{-1}\left\{\frac{2s-5}{s^2-9}\right\}$
- (v)  $\mathcal{L}^{-1}\left\{\frac{1}{s^{3/2}}\right\}$

**Solution:** (i)  $\mathcal{L}^{-1}\left\{\frac{1}{s^4}\right\}$

$$\mathcal{L}^{-1}\left\{\frac{n!}{s^{n+1}}\right\} = t^n$$

$$\begin{aligned}\mathcal{L}^{-1}\left\{\frac{1}{s^4}\right\} &= \mathcal{L}^{-1}\left\{\frac{3!}{s^4} \cdot \frac{1}{3!}\right\} \\ &= \frac{1}{6} \mathcal{L}^{-1}\left\{\frac{9!}{s^4}\right\} = \frac{1}{6} t^3 \quad (\text{Ans:})\end{aligned}$$

Try to generate results of known Laplace transform  
in the argument of  $\mathcal{L}^{-1}$

$$\begin{aligned}\text{(ii)} \quad \mathcal{L}^{-1}\left\{\frac{12}{4-3s}\right\} &= \mathcal{L}^{-1}\left\{\frac{12}{-3(s-4/3)}\right\} \\ &= -4 \mathcal{L}^{-1}\left\{\frac{1}{s-4/3}\right\} \\ &= -4 e^{4/3 t} \quad (\text{Ans:})\end{aligned}$$

$$\begin{aligned}\text{(iii)} \quad \mathcal{L}^{-1}\left\{\frac{23s-15}{s^2+8}\right\} &= 23 \mathcal{L}^{-1}\left\{\frac{s}{s^2+8}\right\} - 15 \mathcal{L}^{-1}\left\{\frac{1}{s^2+8}\right\} \\ &= 23 \mathcal{L}^{-1}\left\{\frac{s}{s^2+(\sqrt{8})^2}\right\} - 15 \mathcal{L}^{-1}\left\{\frac{1}{s+(\sqrt{8})^2}\right\}\end{aligned}$$

$$= 23 \cos(\sqrt{8}t) - 15 \alpha^{-1} \left\{ \frac{\sqrt{8}}{\lambda^2 + (\sqrt{8})^2} - \frac{1}{\sqrt{8}} \right\}$$

$$= 23 \cos(\sqrt{8}t) - 15/\sqrt{8} \sin(\sqrt{8}t) \quad (\text{Ans:})$$

$$(iv) \mathcal{L}^{-1}\left\{\frac{2s-5}{s^2-9}\right\} = 2\alpha^{-1} \left\{ \frac{s}{s^2-9} \right\} - 5\alpha^{-1} \left\{ \frac{1}{s^2-9} \right\}$$

$$= 2 \cosh(3t) - 5/3 \sinh(3t) \quad (\text{Ans:})$$

$$(v) \mathcal{L}^{-1}\left\{\frac{1}{s^{3/2}}\right\} = \alpha^{-1} \left\{ \frac{\Gamma(1/2+1)}{\lambda^{1/2+1}} \cdot \frac{1}{\Gamma(1/2+1)} \right\}$$

$$= \frac{1}{\Gamma(1/2+1)} \alpha^{-1} \left\{ \frac{\Gamma(1/2+1)}{\lambda^{1/2+1}} \right\}$$

$$= \frac{1}{\Gamma(1/2+1)} t^{1/2} \quad (\text{Ans:})$$

### First Translation Theorem in inverse form

$$\alpha \{f(t)\} = f(s) \Rightarrow \alpha \{e^{at} f(t)\} = f(s-a)$$

$$\downarrow \qquad \qquad \qquad \downarrow$$

$$\alpha^{-1} \{F(s)\} = f(t) \Rightarrow \alpha^{-1} \{F(s-a)\} = e^{at} f(t)$$

$$\alpha^{-1} \{F(s)\}|_{s \rightarrow s-a} = e^{at} \alpha^{-1} \{F(s)\}$$

### First Translation Theorem in inverse form

$$\mathcal{L}^{-1}\{F(s)|_{s \rightarrow s-a}\} = e^{at} \mathcal{L}^{-1}\{F(s)\}$$

## Problem

Find:

$$(i) \mathcal{L}^{-1}\left\{\frac{1}{(s-4)^2}\right\} \quad (ii) \mathcal{L}^{-1}\left\{\frac{2}{(s-3)^5}\right\}$$

$$(iii) \mathcal{L}^{-1}\left\{\frac{s-3}{(s-3)^2+1}\right\} \quad (iv) \mathcal{L}^{-1}\left\{\frac{6s-4}{(s^2-4s+20)}\right\}$$

**Solution:**

$$\begin{aligned} (i) \mathcal{L}^{-1}\left\{\frac{1}{(s-4)^2}\right\} &= \mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} \Big|_{s \rightarrow s-4} \\ &= \mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} e^{4t} \\ &= t e^{4t} \quad (\text{Ans:}) \end{aligned}$$

$$\begin{aligned} (ii) \mathcal{L}^{-1}\left\{\frac{2}{(s-3)^5}\right\} &= \mathcal{L}^{-1}\left\{\frac{2}{s^5}\right\} \Big|_{s \rightarrow s-3} \\ &= \mathcal{L}^{-1}\left\{\frac{2}{s^5}\right\} e^{3t} \\ &= \mathcal{L}^{-1}\left\{\frac{2 \cdot 4!}{s^5} \cdot \frac{1}{4!}\right\} e^{3t} \\ &= \frac{1}{12} t^4 e^{3t} \quad (\text{Ans:}) \end{aligned}$$

$$\begin{aligned} (iii) \mathcal{L}^{-1}\left\{\frac{s-3}{(s-3)^2+1}\right\} &= \mathcal{L}^{-1}\left\{\frac{s}{s^2+1}\right\} \Big|_{s \rightarrow s-3} \\ &= \mathcal{L}^{-1}\left\{\frac{s}{s^2+1}\right\} e^{3t} \\ &= \text{const } e^{3t} \quad (\text{Ans:}) \end{aligned}$$

$$(iv) \mathcal{L}^{-1}\left\{\frac{6s-4}{(s^2-4s+20)}\right\}$$

**Hint: completing the square**

$$\begin{aligned}
 \mathcal{L}^{-1}\left\{\frac{6s-4}{(s^2-4s+20)}\right\} &= \alpha^{-1} \left\{ \frac{6s-4}{(s-2)^2+16} \right\} \\
 &= 6 \alpha^{-1} \left\{ \frac{s}{(s-2)^2+16} \right\} - 4 \alpha^{-1} \left\{ \frac{1}{(s-2)^2+16} \right\} \\
 &= 6 \alpha^{-1} \left\{ \frac{s-2+2}{(s-2)^2+16} \right\} - 4 \alpha^{-1} \left\{ \frac{1}{(s-2)^2+16} \right\} \\
 &= 6 \alpha^{-1} \left\{ \frac{s-2}{(s-2)^2+16} \right\} + 12 \alpha^{-1} \left\{ \frac{1}{(s-2)^2+16} \right\} \\
 &\quad - 4 \alpha^{-1} \left\{ \frac{1}{(s-2)^2+16} \right\} \\
 &= 6 \alpha^{-1} \left\{ \frac{s}{s^2+16} \right\} \Big|_{s \rightarrow s-2} + 8 \alpha^{-1} \left\{ \frac{1}{s^2+16} \right\} \\
 &= 6 \alpha^{-1} \left\{ \frac{s}{s^2+16} \right\} e^{2t} + 8 \alpha^{-1} \left\{ \frac{1}{s^2+16} \right\} \Big|_{s \rightarrow s-2} \\
 &= 6 \cos(4t) e^{2t} + 8 \alpha^{-1} \left\{ \frac{1}{s^2+16} \right\} e^{2t} \\
 &= 6 \cos(4t) e^{2t} + 2 \sin(4t) e^{2t} \quad (\text{Ans!})
 \end{aligned}$$

## Problem

Find:

$$(i) \mathcal{L}^{-1}\left\{\frac{s^2+6s+9}{(s+4)(s-1)(s-2)}\right\} \quad (ii) \mathcal{L}^{-1}\left\{\frac{2s+5}{(s-3)^2}\right\}$$

$$(iii) \mathcal{L}^{-1}\left\{\frac{6s^2+50}{(s^2+4)(s+3)}\right\} \quad (iv) \mathcal{L}^{-1}\left\{\frac{3s}{(s+1)(s^2+1)}\right\}$$

$$(v) \mathcal{L}^{-1}\left\{\frac{2s+1}{s(s+1)(s^2+4s+6)}\right\} \quad (vi) \mathcal{L}^{-1}\left\{\frac{s}{(s^2+4)(s^2+9)}\right\}$$

**Solution:** (i)  $\mathcal{L}^{-1}\left\{\frac{s^2+6s+9}{(s+4)(s-1)(s-2)}\right\}$

$$\frac{s^2+6s+9}{(s+4)(s-1)(s-2)} = \frac{A}{s+4} + \frac{B}{s-1} + \frac{C}{s-2}$$

$$\Rightarrow s^2 + 6s + 9 = A(s-1)(s-2) + B(s+1)(s-2) + C(s+1)(s-1)$$

$\downarrow$   
valid for any value of  $s$

$$(i) s = -4 \Rightarrow 16 - 24 + 9 = A(-5)(-6) \Rightarrow A = 1/30$$

$$(ii) s = 1 \Rightarrow 0 = -16/5$$

$$(iii) s = 2 \Rightarrow 25/6$$

$$\frac{s^2+6s+9}{(s+4)(s-1)(s-2)} = \frac{1/30}{s+4} + \frac{-16/5}{s-1} + \frac{25/6}{s-2}$$

$$\begin{aligned} \mathcal{L}^{-1}\left\{\frac{s^2+6s+9}{(s+4)(s-1)(s-2)}\right\} &= \mathcal{L}^{-1}\left\{\frac{1/30}{s+4} + \frac{-16/5}{s-1} + \frac{25/6}{s-2}\right\} \\ &= 1/30 \mathcal{L}^{-1}\left\{\frac{1}{s+4}\right\} - 16/5 \mathcal{L}^{-1}\left\{\frac{1}{s-1}\right\} + 25/6 \mathcal{L}^{-1}\left\{\frac{1}{s-2}\right\} \\ &= 1/30 e^{-4t} - 16/5 e^t + 25/6 e^{2t} \text{ (Ans!)} \end{aligned}$$

$$(ii) \mathcal{L}^{-1}\left\{\frac{2s+5}{(s-3)^2}\right\}$$

$$\frac{2s+5}{(s-3)^2} = \frac{A}{s-3} + \frac{B}{(s-3)^2}$$

$$\Rightarrow 2s+5 = A(s-3) + B$$

$$(i) s=3 \Rightarrow 11 = B$$

$$(ii) \text{ co-efficient of } s: 2 = A$$

$$\frac{2s+5}{(s-3)^2} = \frac{2}{s-3} + \frac{11}{(s-3)^2}$$

$$\begin{aligned} d^{-1} \left\{ \frac{2s+5}{(s-3)^2} \right\} &= d^{-1} \left\{ \frac{2}{s-3} + \frac{11}{(s-3)^2} \right\} \\ &= 2 d^{-1} \left\{ \frac{1}{s-3} \right\} + 11 d^{-1} \left\{ \frac{1}{s^2} \right\} \Big|_{s=3} \\ &= 2e^{3t} + 11 t e^{3t} \\ &= 2e^{3t} + 11 t e^{3t} \quad (\text{Ans!}) \end{aligned}$$

$$(iii) \mathcal{L}^{-1}\left\{\frac{6s^2+50}{(s^2+4)(s+3)}\right\}$$

$$\frac{6s^2+50}{(s+3)(s^2+4)} = \frac{A}{s+3} + \frac{Bs+C}{s^2+4}$$

$$\Rightarrow 6s^2 + 50 = A(s^2 + 4) + (Bs + C)(s + 3)$$

$$(i) s = -3 \Rightarrow 104 = 13A \Rightarrow A = 8$$

$$(ii) s^2: 6 = A + B \Rightarrow B = -2$$

$$(iii) \text{ constant: } 50 = 4A + 3C \Rightarrow C = 6$$

$$\frac{6s^2 + 50}{(s+3)(s^2+4)} = \frac{8}{s+3} + \frac{-2s+6}{s^2+4}$$

$$\begin{aligned} \Rightarrow \alpha^{-1} \left\{ \frac{6s^2 + 50}{(s+3)(s^2+4)} \right\} &= \alpha^{-1} \left\{ \frac{8}{s+3} + \frac{-2s+6}{s^2+4} \right\} \\ &= 8 \alpha^{-1} \left\{ \frac{1}{s+3} \right\} - 2 \alpha^{-1} \left\{ \frac{s}{s^2+1} \right\} + 6 \alpha^{-1} \left\{ \frac{1}{s^2+1} \right\} \\ &= 8e^{-3t} - 2 \cos(2t) + 3 \sin(2t) \quad (\text{Ans:}) \end{aligned}$$

$$(iv) \mathcal{L}^{-1} \left\{ \frac{3s}{(s+1)(s^2+1)} \right\}$$

$$\frac{3s}{(s+1)(s^2+1)} = \frac{A}{s+1} + \frac{Bs+C}{s^2+1}$$

$$\Rightarrow 3s = A(s^2 + 1) + (Bs + C)(s + 1)$$

$$(i) s = -1 \Rightarrow -3 = 2A \Rightarrow A = -\frac{3}{2}$$

$$(ii) s^2: 0 = A + B \Rightarrow B = \frac{3}{2}$$

$$(iii) \text{ constant: } 0 = A + C \Rightarrow C = \frac{3}{2}$$

$$\frac{3s}{(s+1)(s^2+1)} = \frac{-\frac{3}{2}}{s+1} + \frac{\frac{3}{2}s + \frac{3}{2}}{s^2+1}$$

$$\alpha^{-1} \left\{ \frac{3s}{(s+1)(s^2+1)} \right\} = \alpha^{-1} \left\{ \frac{-3/2}{s+1} + \frac{3/2s + 3/2}{s^2+1} \right\}$$

$$= \frac{3}{2} \left( -\alpha^{-1} \left\{ \frac{1}{s+1} \right\} + \alpha^{-1} \left\{ \frac{s}{s^2+1} \right\} + \alpha^{-1} \left\{ \frac{1}{s^2+1} \right\} \right)$$

$$= \frac{3}{2} (-e^{-t} + \text{const} + \sin t) \quad (\text{Ans!})$$

(v)  $\mathcal{L}^{-1} \left\{ \frac{2s+1}{s(s+1)(s^2+4s+6)} \right\}$

$$\frac{2s+1}{s(s+1)(s^2+4s+6)} = \frac{A}{s} + \frac{B}{s+1} + \frac{Cs+D}{s^2+4s+6}$$

$$\Rightarrow s(s+1) = A(s+1)(s^2+4s+6) + Bs(s^2+4s+6) + (Cs+D)s$$

$$(i) s=0 \Rightarrow 1 = A \cdot 6 \Rightarrow A = 1/6$$

$$(ii) s=-1 \Rightarrow -1 = -B(1-4+6) \Rightarrow B = 1/3$$

$$(iii) s=0: 0 = A+B+C \Rightarrow C = -(A+B) = -1/2$$

$$(iv) s=2: 2 = (A+4A+6B+D) \Rightarrow D = -5/3$$

$$\frac{2s+1}{s(s+1)(s^2+4s+6)} = \frac{1/6}{s} + \frac{1/3}{s+1} + \frac{-1/2s - 5/3}{s^2+4s+6}$$

$$\alpha^{-1} \left\{ \frac{2s+1}{s(s+1)(s^2+4s+6)} \right\} = \alpha^{-1} \left\{ \frac{1/6}{s} + \frac{1/3}{s+1} + \frac{-1/2s - 5/3}{s^2+4s+6} \right\}$$

$$= \frac{1}{6} \alpha^{-1} \left\{ 1/s \right\} + \frac{1}{3} \alpha^{-1} \left\{ \frac{1}{s+1} \right\} - \frac{1}{2} \alpha^{-1} \left\{ \frac{s}{(s+2)^2+2} \right\}$$

$$- \frac{5}{3} \alpha^{-1} \left\{ \frac{1}{(s+2)^2+2} \right\}$$

$$= \frac{1}{6} + \frac{1}{3} e^{-t} - \frac{1}{2} \alpha^{-1} \left\{ \frac{s+2-2}{(s+2)^{\gamma}+2} \right\} \\ - \frac{5}{3} \alpha^{-1} \left\{ \frac{1}{(s+2)^{\gamma}+2} \right\}$$

$$= \frac{1}{6} + \frac{1}{3} e^{-t} - \frac{1}{2} \alpha^{-1} \left\{ \frac{s+2}{(s+2)^{\gamma}+(\sqrt{2})^{\gamma}} \right\} + \alpha^{-1} \left\{ \frac{1}{(s+2)^{\gamma}+2} \right\}$$

$$- \frac{5}{3} \alpha^{-1} \left\{ \frac{1}{(s+2)^{\gamma}+2} \right\}$$

$$= \frac{1}{6} + \frac{1}{3} e^{-t} - \frac{1}{2} \alpha^{-1} \left\{ \frac{s}{s^{\gamma}+(\sqrt{2})^{\gamma}} \Big|_{s \rightarrow s+2} \right\}$$

$$- \frac{2}{3} \alpha^{-1} \left\{ \frac{1}{s^{\gamma}+(\sqrt{2})^{\gamma}} \Big|_{s \rightarrow s+2} \right\}$$

$$= \frac{1}{6} + \frac{1}{3} e^{-t} - \frac{1}{2} \cos(\sqrt{2}t) e^{-2t} - \frac{2}{3\sqrt{2}} \sin(\sqrt{2}t) e^{-2t}$$

(Ans!)

(vi)  $\mathcal{L}^{-1} \left\{ \frac{s}{(s^2+4)(s^2+9)} \right\}$

$$\frac{s}{(s^2+4)(s^2+9)} = \frac{As+B}{s^2+4} + \frac{Cs+D}{s^2+9}$$

$$\Rightarrow s = (As+B)(s^2+9) + (Cs+D)(s^2+4)$$

$$\Rightarrow s = As^3 + 9As + Bs^2 + 9B + Cs^3 + 4Cs + Ds^2 + 4D$$

$$\Rightarrow s = (A+c)s^3 + (B+D)s^2 + (9A+4c)s + 9B+4D$$

$$\text{Case 1: } 0 = A + C \xrightarrow{\text{(i)}} A = -C \Rightarrow A = 1/5$$

$$\text{Case 2: } 0 = B + D \xrightarrow{\text{(ii)}} B = -D \Rightarrow B = 0$$

$$\text{Case 3: } 1 = 9A + 4C \xrightarrow{\text{(iii)}} -9C + 4C = 1 \Rightarrow C = -1/5$$

$$\text{constant: } 0 = 9B + 4D \xrightarrow{\text{(iv)}} -9D + 4D = 0 \Rightarrow D = 0$$

$$\frac{s}{(s^2+4)(s^2+9)} = \frac{1/5}{s^2+4} + \frac{-1/5}{s^2+9}$$

$$\begin{aligned} d^{-1} \left\{ \frac{s}{(s^2+4)(s^2+9)} \right\} &= d^{-1} \left\{ \frac{1/5}{s^2+4} + \frac{-1/5}{s^2+9} \right\} \\ &= 1/5 \{ \cos(2t) - \cos(3t) \} \end{aligned}$$

(Ans!)

# MAT215: Complex Variables & Laplace Transformations

## Chapter : Laplace Transformations

2nd Translation theorem in inverse form  
Laplace transform of derivatives  
Solving Initial Value Problem (IVP) using Laplace transform



BRAC University  
Department of Mathematics and Natural Sciences  
Summer 2025

Instructor: Kazi Hafizur Rahman

## Second Translation Theorem in inverse form

$$\alpha \{f(t)\} = F(s) \Rightarrow \alpha \{f(t-a) u(t-a)\} = e^{-as} F(s)$$

$$\downarrow \qquad \qquad \downarrow$$

$$\alpha^{-1}\{F(s)\} = f(t) \Rightarrow \alpha^{-1}\{e^{-as} F(s)\} = f(t-a) u(t-a)$$

$$\alpha^{-1}\{e^{-as} F(s)\} = \alpha^{-1}\{F(s)\}|_{t \rightarrow t-a} u(t-a)$$

## Second Translation Theorem in inverse form

$$\mathcal{L}^{-1}\{e^{-as}F(s)\} = \mathcal{L}^{-1}\{F(s)\}|_{t \rightarrow t-a} u(t-a)$$

### Problem

Find:

$$(i) \mathcal{L}^{-1}\left\{\frac{1}{(s-4)}e^{-2s}\right\} \quad (ii) \mathcal{L}^{-1}\left\{\frac{3s}{(s+1)(s^2+1)}e^{-\pi s}\right\}$$

### Solution:

$$(i) \mathcal{L}^{-1}\left\{\frac{1}{(s-4)}e^{-2s}\right\} = \alpha^{-1}\left\{\frac{1}{s-4}\right\}|_{t \rightarrow t-2} u(t-2)$$

$$= e^{4t}|_{t \rightarrow t-2} u(t-2)$$

$$= e^{4(t-2)} u(t-2) \quad (\text{Ans!})$$

$$(ii) \mathcal{L}^{-1}\left\{\frac{3s}{(s+1)(s^2+1)}e^{-\pi s}\right\} = \alpha^{-1}\left\{\frac{3s}{(s+1)(s^2+1)}\right\}|_{t \rightarrow t-\pi} u(t-\pi)$$

Finding  $\alpha^{-1}\left\{\frac{3s}{(s+1)(s^2+1)}\right\}$ :

$$\frac{3s}{(s+1)(s^2+1)} = \frac{A}{s+1} + \frac{Bs+C}{s^2+1}$$

$$\Rightarrow 3s = A(s^2+1) + (Bs+C)(s+1)$$

$$(i) s=-1 \Rightarrow -3 = 2A \Rightarrow A = -\frac{3}{2}$$

$$(ii) s^2 : 0 = A + B \Rightarrow B = \frac{3}{2}$$

$$(iii) \text{ constant} : 0 = A + C \Rightarrow C = \frac{3}{2}$$

$$\frac{3s}{(s+1)(s^2+1)} = \frac{-\frac{3}{2}}{s+1} + \frac{\frac{3}{2}s + \frac{3}{2}}{s^2+1}$$

$$\mathcal{L}^{-1} \left\{ \frac{3s}{(s+1)(s^2+1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{-\frac{3}{2}}{s+1} + \frac{\frac{3}{2}s + \frac{3}{2}}{s^2+1} \right\}$$

$$= \frac{3}{2} \left( -\mathcal{L}^{-1} \left\{ \frac{1}{s+1} \right\} + \mathcal{L}^{-1} \left\{ \frac{s}{s^2+1} \right\} + \mathcal{L}^{-1} \left\{ \frac{1}{s^2+1} \right\} \right)$$

$$= \frac{3}{2} (-e^{-t} + \text{const} + \sin t)$$

$$\mathcal{L}^{-1} \left\{ \frac{3s}{(s+1)(s^2+1)} e^{-\pi s} \right\} = \mathcal{L}^{-1} \left\{ \frac{3s}{(s+1)(s^2+1)} \right\} |_{t-\pi} u(t-\pi)$$

$$= \frac{3}{2} \left\{ -e^{-(t-\pi)} + \sin(t-\pi) + \cos(t-\pi) \right\} u(t-\pi) \quad (\text{Ans.})$$

## Laplace transformation of derivatives

If  $\mathcal{L}\{f(t)\} = F(s)$ , then

- (i)  $\mathcal{L}\{f'(t)\} = ?$
- (ii)  $\mathcal{L}\{f''(t)\} = ?$
- (iii)  $\mathcal{L}\{f'''(t)\} = ?$  if  $s > 0$
- (iv) What is the generalization?

**Solution:**  $\mathcal{L}\{f(t)\} = F(s)$

$$\mathcal{L}\{f'(t)\} = \int_0^\infty e^{-st} f'(t) dt$$

$$= \lim_{p \rightarrow \infty} \int_0^p e^{-st} f'(t) dt$$

$\int u v dx = u \int v dx - \int (u' \int v dx) dx$  (integration by parts formula)  
choosing  $u, v$ : LIATE formula

$$u = e^{-st}$$

$$v = f'(t)$$

$$= \lim_{p \rightarrow \infty} \left\{ e^{-sp} f(p) \Big|_0^p - \int_0^p e^{-st} (-s) f(t) dt \right\}$$

$$= \lim_{p \rightarrow \infty} \left\{ e^{-sp} f(p) - f(0) + s \int_0^p e^{-st} f(t) dt \right\}$$

$$= \lim_{p \rightarrow \infty} e^{-sp} f(p) - f(0) + s \lim_{p \rightarrow \infty} \int_0^p e^{-st} f(t) dt$$

$$= 0 - f(0) + s \int_0^\infty e^{-st} f(t) dt$$

$$= s \alpha \{f(t)\} - f(0)$$

$$\therefore \alpha \{f'(t)\} = s \alpha \{f(t)\} - f(0)$$

$$\alpha \{f''(t)\} = \alpha \{(f'(t))'\}$$

$$= s \alpha \{f'(t)\} - f'(0)$$

$$= s \int_0^\infty f(t) dt - f(0) = f'(0)$$

$$\alpha \int_0^\infty f''(t) dt = s^2 \int_0^\infty f(t) dt - s f(0) - f'(0)$$

$$\alpha \int_0^\infty f'''(t) dt = s \alpha \int_0^\infty f''(t) dt - f''(0)$$

$$= s \int_0^\infty s^2 \alpha \int_0^\infty f(t) dt - s f(0) - f'(0) - f''(0)$$

$$\alpha \int_0^\infty f'''(t) dt = s^3 \alpha \int_0^\infty f(t) dt - s^2 f(0) - s f'(0) - f''(0)$$

## Laplace transformation of derivatives

If  $\mathcal{L}\{f(t)\} = F(s)$ , then

$$\mathcal{L}\{f'(t)\} = s \cdot F(s) - f(0)$$

$$\mathcal{L}\{f''(t)\} = s^2 \cdot F(s) - s \cdot f(0) - f'(0)$$

$$\mathcal{L}\{f'''(t)\} = s^3 \cdot F(s) - s^2 \cdot f(0) - s \cdot f'(0) - f''(0)$$

Provided that  $s > 0$ .

## Initial Value Problem

$x' + 2x + 1 = 0 \rightarrow \text{unknown } x \rightarrow \text{number}$

(Algebraic equation)

$$\frac{dy}{dt} + \frac{dy}{dt} + y + \sin t = 0 \quad (i)$$

Initial Value Problem (IVP)

$$\begin{cases} y(0) = 2 \\ y'(0) = 4 \end{cases} \quad (ii)$$

unknown  $y(t)$

(initial conditions)

Differential equation for  
the unknown  $y(t)$

# Solving IVP Using Laplace Transformation



**Step 1: Apply Laplace Transform (  $y(t) \rightarrow Y(s)$  )**

**Step 2: Find  $Y(s)$**

**Step 3: Apply Inverse Laplace Transform (  $Y(s) \rightarrow y(t)$  )**

## Problem

Solve the following initial value problems:

$$(i) \frac{dy}{dt} + 3y = 13 \sin 2t, \quad y(0) = 6.$$

$$(ii) y'' - 3y' + 2y = e^{-4t}, \quad y(0) = 1, \quad y'(0) = 5.$$

$$(iii) y'' - 6y' + 9y = t^2 e^{3t}, \quad y(0) = 2, \quad y'(0) = 17$$

$$(iv) y'' + 4y' + 6y = 1 + e^{-t}, \quad y(0) = 0, \quad y'(0) = 0.$$

$$(v) y' + y = f(t), \quad y(0) = 5, \quad \text{where } f(t) = \begin{cases} 0, & 0 \leq t < \pi \\ 3 \cos t, & t \geq \pi \end{cases}$$

$$(i) \frac{dy}{dt} + 3y = 13 \sin 2t, \quad y(0) = 6.$$

**Solution:**

$$\alpha \{y'(t) + 3y\} = \alpha \{13 \sin 2t\}$$

$$\Rightarrow \alpha \{y'(t)\} + 3\alpha \{y(t)\} = 13 \frac{2}{s^2+4}$$

let,

$$\alpha \{y(t)\} = Y(s)$$

$$\alpha \{y'(t)\} = sY(s) - y(0)$$

$$sY(s) - y(0) + 3Y(s) = \frac{26}{s^2+4}$$

$$\Rightarrow (s+3)Y(s) = 6 + \frac{26}{s^2+4} = \frac{6s^2+50}{s^2+4}$$

$$\Rightarrow Y(s) = \frac{6s^2+50}{(s+3)(s^2+4)}$$

$$\Rightarrow \alpha^{-1}\{Y(s)\} = \alpha^{-1}\left\{\frac{6s^2+50}{(s+3)(s^2+4)}\right\}$$

$$\Rightarrow y(t) = \alpha^{-1}\left\{\frac{6s^2+50}{(s+3)(s^2+4)}\right\}$$

**Finding**  $\mathcal{L}^{-1}\left\{\frac{6s^2+50}{(s^2+4)(s+3)}\right\}$

$$\frac{6s^2+50}{(s+3)(s^2+4)} = \frac{A}{s+3} + \frac{Bs+C}{s^2+4}$$

$$\Rightarrow 6s^2 + 50 = A(s^2 + 4) + (Bs + C)(s + 3)$$

$$(i) s = -3 \Rightarrow 104 = 13A \Rightarrow A = 8$$

$$(ii) s^2 : 6 = A + B \Rightarrow B = -2$$

$$(iii) \text{ constant} : 50 = 4A + 3C \Rightarrow C = 6$$

$$\frac{6s^2 + 50}{(s+3)(s^2+4)} = \frac{8}{s+3} + \frac{-2s+6}{s^2+4}$$

$$\Rightarrow \mathcal{d}^{-1} \left\{ \frac{6s^2 + 50}{(s+3)(s^2+4)} \right\} = \mathcal{d}^{-1} \left\{ \frac{8}{s+3} + \frac{-2s+6}{s^2+4} \right\}$$

$$= 8 \mathcal{d}^{-1} \left\{ \frac{1}{s+3} \right\} - 2 \mathcal{d}^{-1} \left\{ \frac{s}{s^2+4} \right\} + 6 \mathcal{d}^{-1} \left\{ \frac{1}{s^2+4} \right\}$$

$$= 8e^{-3t} - 2 \cos(2t) + 3 \sin(2t)$$

Therefore,

$$y(t) = 8e^{-3t} - 2 \cos(2t) + 3 \sin(2t) \quad (\text{Ans})$$

$$(ii) y'' - 3y' + 2y = e^{-4t}, \quad y(0) = 1, \quad y'(0) = 5.$$

Solution:

$$\mathcal{d} \left\{ y'' - 3y' + 2y \right\} = \mathcal{d} \left\{ e^{-4t} \right\}$$

$$\Rightarrow \mathcal{d} \left\{ y'' \right\} - 3 \mathcal{d} \left\{ y' \right\} + 2 \mathcal{d} \left\{ y \right\} = \frac{1}{s+4}$$

Let,

$$\mathcal{d} \left\{ y(t) \right\} = Y(s)$$

$$\mathcal{d} \left\{ y'(t) \right\} = sY(s) - y(0) = sY(s) - 1$$

$$\mathcal{d} \left\{ y''(t) \right\} = s^2 Y(s) - sy(0) - y'(0) = s^2 Y(s) - s - 5$$

$$s^2 Y(s) - s - 5 - 9 \quad \{sY(s) - 1\} + 2Y(s) = \frac{1}{s+4}$$

$$\Rightarrow (s^2 - 9s + 2) Y(s) = s + 5 - 9 + \frac{1}{s+4}$$

$$\Rightarrow (s-1)(s-2) Y(s) = \frac{(s+2)(s+4) + 1}{s+4}$$

$$\Rightarrow Y(s) = \frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)}$$

$$\Rightarrow \mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)}\right\}$$

Finding  $\mathcal{L}^{-1}\left\{\frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)}\right\}$

$$\frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)} = \frac{A}{s+4} + \frac{B}{s-1} + \frac{C}{s-2}$$

$$\Rightarrow s^2 + 6s + 9 = A(s-1)(s-2) + B(s+4)(s-2) + C(s+4)(s-1)$$

↓  
valid for any value of s

$$(i) \ s = -4 \Rightarrow 16 - 24 + 9 = A(-5)(-6) \Rightarrow A = 1/30$$

$$(ii) \ s = 1 \Rightarrow 0 = -16/5$$

$$(iii) \ s = 2 \Rightarrow 0 = 25/6$$

$$\frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)} = \frac{1/30}{s+4} + \frac{-16/5}{s-1} + \frac{25/6}{s-2}$$

$$\begin{aligned}
 & \mathcal{d}^{-1} \left\{ \frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)} \right\} = \mathcal{d}^{-1} \left\{ \frac{1/30}{s+4} + \frac{-16/5}{s-1} + \frac{25/6}{s-2} \right\} \\
 & = 1/30 \mathcal{d}^{-1} \left\{ \frac{1}{s+4} \right\} - 16/5 \mathcal{d}^{-1} \left\{ \frac{1}{s-1} \right\} \\
 & \quad + 25/6 \mathcal{d}^{-1} \left\{ \frac{1}{s-2} \right\} \\
 & = 1/30 e^{-4t} - 16/5 e^t + 25/6 e^{2t}
 \end{aligned}$$

Therefore we have,

$$\begin{aligned}
 \Rightarrow y(t) &= \mathcal{d}^{-1} \left\{ \frac{s^2 + 6s + 9}{(s+4)(s-1)(s-2)} \right\} \\
 \therefore y(t) &= 1/30 e^{-4t} - 16/5 e^t + 25/6 e^{2t} \quad (\text{Ans})
 \end{aligned}$$

$$(iii) y'' - 6y' + 9y = t^2 e^{3t}, \quad y(0) = 2, \quad y'(0) = 17$$

Solution:

$$\begin{aligned}
 \mathcal{d} \left\{ y'' - 6y' + 9y \right\} &= \mathcal{d} \left\{ t^2 e^{3t} \right\} \\
 \Rightarrow \mathcal{d} \left\{ y'' \right\} - 6 \mathcal{d} \left\{ y' \right\} + 9 \mathcal{d} \left\{ y \right\} &= \mathcal{d} \left\{ t^2 \right\} \Big|_{s \rightarrow s-3}
 \end{aligned}$$

Let,

$$\mathcal{d} \left\{ y(t) \right\} = Y(s)$$

$$\mathcal{d} \left\{ y'(t) \right\} = sY(s) - y(0) = sY(s) - 2$$

$$\mathcal{d} \left\{ y''(t) \right\} = s^2 Y(s) - sy(0) - y'(0) = s^2 Y(s) - 2s - 1$$

$$s^2 Y(s) - 2s - 1 - 6 \left\{ sY(s) - 2 \right\} + 9Y(s) = \frac{2}{(s-3)^3}$$

$$\Rightarrow (s^2 - 6s + 9)Y(s) - 2s - 1 + 12 = \frac{2}{(s-3)^3}$$

$$\Leftrightarrow (\lambda - 3)^2 Y(\lambda) = 2\lambda + 5 + \frac{2}{(\lambda - 3)^3}$$

$$\Rightarrow Y(\lambda) = \frac{2\lambda + 5}{(\lambda - 3)^2} + \frac{2}{(\lambda - 3)^5}$$

$$\Leftrightarrow \alpha^{-1} \{Y(\lambda)\} = \alpha^{-1} \left\{ \frac{2\lambda + 5}{(\lambda - 3)^2} + \frac{2}{(\lambda - 3)^5} \right\}$$

$$\Rightarrow y(t) = \alpha^{-1} \left\{ \frac{2\lambda + 5}{(\lambda - 3)^2} \right\} + \alpha^{-1} \left\{ \frac{2}{(\lambda - 3)^5} \right\}$$

**Finding**  $\mathcal{L}^{-1}\left\{\frac{2}{(s-3)^5}\right\}$  &  $\mathcal{L}^{-1}\left\{\frac{2s+5}{(s-3)^2}\right\}$

$$(ii) \mathcal{L}^{-1}\left\{\frac{2}{(s-3)^5}\right\} = \alpha^{-1} \left\{ \frac{2}{s^5} \Big|_{s \rightarrow s-3} \right\}$$

$$= \alpha^{-1} \left\{ \frac{2}{s^5} \right\} e^{3t}$$

$$= \alpha^{-1} \left\{ \frac{2 \cdot 4!}{s^5} \cdot \frac{1}{4!} \right\} e^{3t}$$

$$= \frac{1}{12} t^4 e^{3t}$$

$$(ii) \mathcal{L}^{-1}\left\{\frac{2s+5}{(s-3)^2}\right\}$$

$$\frac{2s+5}{(s-3)^2} = \frac{A}{s-3} + \frac{B}{(s-3)^2}$$

$$\Rightarrow 2s+5 = A(s-3) + B$$

$$(i) s=3 \Rightarrow 11 = B$$

$$(ii) \text{ Coefficient of } s: 2 = A$$

$$\frac{2s+5}{(s-3)^2} = \frac{2}{s-3} + \frac{11}{(s-3)^2}$$

$$\begin{aligned} \mathcal{L}^{-1}\left\{\frac{2s+5}{(s-3)^2}\right\} &= \mathcal{L}^{-1}\left\{\frac{2}{s-3} + \frac{11}{(s-3)^2}\right\} \\ &= 2 \mathcal{L}^{-1}\left\{\frac{1}{s-3}\right\} + 11 \mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} \Big|_{s=3} \\ &= 2e^{3t} + 11 \mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} e^{3t} \\ &= 2e^{3t} + 11t e^{3t} \end{aligned}$$

Therefore we have,

$$y(t) = 2e^{3t} + 11t e^{3t} + \frac{1}{12} t^4 e^{3t} \quad (\text{Ans.})$$

$$(v) y' + y = f(t), \quad y(0) = 5, \quad \text{where } f(t) = \begin{cases} 0, & 0 \leq t < \pi \\ 3 \cos t, & t \geq \pi \end{cases}$$

$$\alpha \{y' + y\} = \alpha \{f(t)\}$$

$$\Rightarrow \alpha \{y'\} + \alpha \{y\} = \alpha \{f(t)\}$$

let,  $\alpha \{y(t)\} = Y(s)$

$$\alpha \{y'(t)\} = sY(s) - y(0) = sY(s) - 5$$

$$f(t) = 3 \cos t u(t-\pi)$$

$$g(t) = \cos(t+\pi)$$

$$\alpha \{f(t)\} = \alpha \{3 \cos t u(t-\pi)\} = -\cos t$$

$$= 3 \alpha \{\cos(t-\pi+\pi) u(t-\pi)\}$$

$$= 3 \alpha \{g(t-\pi) u(t-\pi)\}$$

$$= 3 \alpha \{g(t)\} e^{-\pi s}$$

$$= 3 \alpha \{-\cos t\} e^{-\pi s} = -\frac{3s}{s^2+1} e^{-\pi s}$$

$$sY(s) - 5 + Y(s) = -\frac{3s}{s^2+1} e^{-\pi s}$$

$$\Rightarrow (s+1)Y(s) = 5 - \frac{3s}{s^2+1} e^{-\pi s}$$

$$\Rightarrow Y(s) = \frac{5}{s+1} - \frac{3s}{(s+1)(s^2+1)} e^{-\pi s}$$

$$\Rightarrow y(t) = \alpha^{-1} \left\{ \frac{5}{s+1} \right\} - \alpha^{-1} \left\{ \frac{3s}{(s+1)(s^2+1)} e^{-\pi s} \right\}$$

$$\Rightarrow y(t) = 5e^{-t} - \alpha^{-1} \left\{ \frac{3s}{(s+1)(s^2+1)} e^{-\pi s} \right\}$$

**Finding**  $\mathcal{L}^{-1}\left\{\frac{3s}{(s+1)(s^2+1)}e^{-\pi s}\right\}$

$$\frac{3s}{(s+1)(s^2+1)} = \frac{A}{s+1} + \frac{Bs+C}{s^2+1}$$

$$\Rightarrow 3s = A(s^2+1) + (Bs+C)(s+1)$$

$$(i) s=-1 \Rightarrow -3 = 2A \Rightarrow A = -\frac{3}{2}$$

$$(ii) s^2 : 0 = A + B \Rightarrow B = \frac{3}{2}$$

$$(iii) \text{ constant} : 0 = A + C \Rightarrow C = \frac{3}{2}$$

$$\frac{3s}{(s+1)(s^2+1)} = \frac{-\frac{3}{2}}{s+1} + \frac{\frac{3}{2}s + \frac{3}{2}}{s^2+1}$$

$$\alpha^{-1} \left\{ \frac{3s}{(s+1)(s^2+1)} \right\} = \alpha^{-1} \left\{ \frac{-\frac{3}{2}}{s+1} + \frac{\frac{3}{2}s + \frac{3}{2}}{s^2+1} \right\}$$

$$= \frac{3}{2} \left( -\alpha^{-1} \left\{ \frac{1}{s+1} \right\} + \alpha^{-1} \left\{ \frac{s}{s^2+1} \right\} + \alpha^{-1} \left\{ \frac{1}{s^2+1} \right\} \right)$$

$$= \frac{3}{2} (-e^{-t} + \cos t + \sin t)$$

$$\mathcal{L}^{-1}\left\{\frac{3s}{(s+1)(s^2+1)}e^{-\pi s}\right\} = \alpha^{-1} \left\{ \frac{3s}{(s+1)(s^2+1)} \right\} |_{t-\pi} u(t-\pi)$$

$$= \frac{3}{2} \left\{ -e^{-(t-\pi)} + \sin(t-\pi) + \cos(t-\pi) \right\} u(t-\pi)$$

Therefore we have,

$$= 5e^{-t} - \frac{3}{2} \left\{ -e^{-(t-\pi)} + \sin(t-\pi) + \cos(t-\pi) \right\} u(t-\pi)$$

(Ans.)

(iv)  $y'' + 4y' + 6y = 1 + e^{-t}$ ,  $y(0) = 0$ ,  $y'(0) = 0$ .

Solution:

$$\begin{aligned} d \{y'' + 4y' + 6y\} &= d\{1 + e^{-t}\} \\ \Rightarrow d\{y''\} + 4d\{y'\} + 6d\{y\} &= \frac{1}{s} + \frac{1}{s+1} \end{aligned}$$

Let,

$$d\{y(t)\} = Y(s)$$

$$d\{y'(t)\} = sY(s) - y(0) = sY(s)$$

$$d\{y''(t)\} = s^2 Y(s) - sy(0) - y'(0) = s^2 Y(s)$$

$$\therefore s^2 Y(s) + 4s Y(s) + 6Y(s) = \frac{1}{s} + \frac{1}{s+1}$$

$$\Rightarrow (s^2 + 4s + 6) Y(s) = \frac{2s+1}{s(s+1)}$$

$$\Rightarrow Y(s) = \frac{2s+1}{s(s+1)(s^2+4s+6)}$$

$$\Rightarrow d^{-1}\{Y(s)\} = d^{-1}\left\{\frac{2s+1}{s(s+1)(s^2+4s+6)}\right\}$$

$$\Rightarrow y(t) = d^{-1}\left\{\frac{2s+1}{s(s+1)(s^2+4s+6)}\right\}$$

Finding  $\mathcal{L}^{-1}\left\{\frac{2s+1}{s(s+1)(s^2+4s+6)}\right\}$

$$\frac{2s+1}{s(s+1)(s^2+4s+6)} = \frac{A}{s} + \frac{B}{s+1} + \frac{Cs+D}{s^2+4s+6}$$

$$\Rightarrow 2s+1 = A(s+1)(s^2+4s+6) + Bs(s^2+4s+6) + (Cs+D)s$$

$$(i) s=0 \Rightarrow 1 = A \cdot 6 \Rightarrow A = 1/6$$

$$(ii) s=-1 \Rightarrow -1 = -B(1-4+6) \Rightarrow B = 1/3$$

$$(iii) s^2: 0 = A+B+C \Rightarrow C = -(A+B) = -1/2$$

$$(iv) s : 2 = CA + 4A + 6B + D \Rightarrow D = -5/3$$

$$\frac{2s+1}{s(s+1)(s^2+4s+6)} = \frac{1/6}{s} + \frac{1/3}{s+1} + \frac{-1/2s - 5/3}{s^2+4s+6}$$

$$\mathcal{L}^{-1}\left\{\frac{2s+1}{s(s+1)(s^2+4s+6)}\right\} = \mathcal{L}^{-1}\left\{\frac{1/6}{s} + \frac{1/3}{s+1} + \frac{-1/2s - 5/3}{s^2+4s+6}\right\}$$

$$= 1/6 \mathcal{L}^{-1}\left\{\frac{1}{s}\right\} + 1/3 \mathcal{L}^{-1}\left\{\frac{1}{s+1}\right\} - 1/2 \mathcal{L}^{-1}\left\{\frac{s}{(s+2)^2+2}\right\}$$

$$- 5/3 \mathcal{L}^{-1}\left\{\frac{1}{(s+2)^2+2}\right\}$$

$$= 1/6 + 1/3 e^{-t} - 1/2 \mathcal{L}^{-1}\left\{\frac{s+2-2}{(s+2)^2+2}\right\}$$

$$- 5/3 \mathcal{L}^{-1}\left\{\frac{1}{(s+2)^2+2}\right\}$$

$$\begin{aligned}
&= \frac{1}{6} + \frac{1}{3} e^{-t} - \frac{1}{2} d^{-1} \left\{ \frac{\frac{s+2}{s}}{(s+2)^s + (\sqrt{2})^s} \right\} + d^{-1} \left\{ \frac{\frac{1}{s}}{(s+2)^s + 2} \right\} \\
&\quad - \frac{5}{3} d^{-1} \left\{ \frac{1}{(s+2)^s + 2} \right\} \\
&= \frac{1}{6} + \frac{1}{3} e^{-t} - \frac{1}{2} d^{-1} \left\{ \frac{\frac{s}{s^s + (\sqrt{2})^s}}{|s \rightarrow s+2} \right\} \\
&\quad - \frac{2}{3} d^{-1} \left\{ \frac{1}{s^s + (\sqrt{2})^s} |_{s \rightarrow s+2} \right\} \\
&= \frac{1}{6} + \frac{1}{3} e^{-t} - \frac{1}{2} \cos(\sqrt{2}t) e^{-2t} - \frac{2}{3\sqrt{2}} \sin(\sqrt{2}t) e^{-2t}
\end{aligned}$$

Therefore we have,

$$y(t) = \frac{1}{6} + \frac{1}{3} e^{-t} - \frac{1}{2} \cos(\sqrt{2}t) e^{-2t} - \frac{2}{3\sqrt{2}} \sin(\sqrt{2}t) e^{-2t}$$

(Ans.)

# MAT215: Complex Variables & Laplace Transformations

## Chapter : Laplace Transformations

Solving Boundary Value Problem (BVP) using Laplace transform

Solving System of equations using Laplace transform



BRAC University  
Department of Mathematics and Natural Sciences  
Summer 2025

Instructor: Kazi Hafizur Rahman

# Solving BVP Using Laplace Transformation

Step 1: Apply Laplace Transform ( $y(t) \rightarrow Y(s)$ )

Step 2: Find  $Y(s)$

Step 3: Apply Inverse Laplace Transform ( $Y(s) \rightarrow y(t)$ )

Step 4: Find the unknown constant using boundary condition

## Problem

Solve the BVP  $y'' + 9y = \cos 2t$ ;  $y(0) = 1$ ,  $y(\pi/2) = -1$ .

## Solution:

$$\mathcal{L}\{y'' + 9y\} = \mathcal{L}\{\cos 2t\}$$

$$\Rightarrow \mathcal{L}\{y''\} + 9\mathcal{L}\{y\} = \frac{s^2}{s^2 + 4}$$

Let,

$$\mathcal{L}\{y(t)\} = Y(s)$$

$$\mathcal{L}\{y''(t)\} = s^2 Y(s) - sy(0) - y'(0) = s^2 Y(s) - s - y'(0)$$

$$\therefore s^2 Y(s) - s - y'(0) + 9Y(s) = \frac{s^2}{s^2 + 4}$$

$$\Rightarrow (s^2 + 9)Y(s) = s + y'(0) + \frac{s^2}{s^2 + 4}$$

$$\Rightarrow Y(s) = \frac{s}{s^2 + 9} + \frac{y'(0)}{s^2 + 9} + \frac{s^2}{(s^2 + 9)(s^2 + 4)}$$

$$\Rightarrow \mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{s}{s^2 + 9} + \frac{y'(0)}{s^2 + 9} + \frac{s^2}{(s^2 + 9)(s^2 + 4)}\right\}$$

$$\Rightarrow y(t) = \mathcal{L}^{-1}\left\{\frac{h}{s^2+9}\right\} + y'(0) \mathcal{L}^{-1}\left\{\frac{1}{s^2+9}\right\} \\ + \mathcal{L}^{-1}\left\{\frac{h}{(s^2+4)(s^2+9)}\right\}$$

$$\therefore y(t) = \cos(3t) + \frac{y'(0)}{3} \sin(3t) + \mathcal{L}^{-1}\left\{\frac{h}{(s^2+4)(s^2+9)}\right\}$$

$$\therefore y(t) = \cos(3t) + A \sin(3t) + \mathcal{L}^{-1}\left\{\frac{h}{(s^2+4)(s^2+9)}\right\}$$

Finding  $\mathcal{L}^{-1}\left\{\frac{s}{(s^2+4)(s^2+9)}\right\} \quad \vdots$

$$\frac{s}{(s^2+4)(s^2+9)} = \frac{As+B}{s^2+4} + \frac{Cs+D}{s^2+9}$$

$$\Rightarrow h = (As+B)(s^2+9) + (Cs+D)(s^2+4)$$

$$\Rightarrow h = As^3 + 9As + Bs^2 + 9B + Cs^3 + 4Cs + Ds^2 + 4D$$

$$\Rightarrow h = (A+C)s^3 + (B+D)s^2 + (9A+4C)s + 9B+4D$$

$$s^3: 0 = A+C \quad \text{(i)} \Rightarrow A = -C \Rightarrow A = 1/5$$

$$s^2: 0 = B+D \quad \text{(ii)} \Rightarrow B = -D \Rightarrow B = 0$$

$$s: 1 = 9A + 4C \quad \text{(iii)} \Rightarrow -9C + 4C = 1 \Rightarrow C = -1/5$$

$$\text{constant: } 0 = 9B + 4D \quad \text{(iv)} \Rightarrow -9D + 4D = 0 \Rightarrow D = 0$$

$$\frac{s}{(s^2+4)(s^2+9)} = \frac{1/5h}{s^2+4} + \frac{-1/5h}{s^2+9}$$

$$\mathcal{L}^{-1}\left\{\frac{s}{(s^2+4)(s^2+9)}\right\} = \mathcal{L}^{-1}\left\{\frac{1/5s}{s^2+4} + \frac{-1/5s}{s^2+9}\right\}$$

$$= \frac{1}{5} \left\{ \cos(2t) - \cos(3t) \right\}$$

$$y(t) = \cos(3t) + A \sin(3t) + \frac{1}{5} \left\{ \cos(2t) - \cos(3t) \right\}$$

$$y(\pi/2) = -1$$

$$\cos(3\pi/2) + A \sin(3\pi/2) + \frac{1}{5} \left\{ \cos(2\pi/2) - \cos(3\pi/2) \right\} = -1$$

$$0 - A + \frac{1}{5}(-1 - 0) = -1$$

$$\Rightarrow -A - \frac{1}{5} = -1 \Rightarrow A = \frac{4}{5}$$

Therefore we have,

$$y(t) = \cos(3t) + \frac{4}{5} \sin(3t) + \frac{1}{5} \left\{ \cos(2t) - \cos(3t) \right\} \quad (\text{Ans})$$

### Problem

$$(i) \mathcal{L}^{-1}\left\{\frac{2s}{(s+1)(s-1)}\right\} \quad (ii) \mathcal{L}^{-1}\left\{\frac{2}{(s+1)(s-1)(s^2+1)}\right\}$$

**Ans:**

$$(i) e^{-t} + e^t$$

$$(ii) \frac{1}{2} (e^t - e^{-t}) - \sin t$$

### Shortcut: Inverse of a $2 \times 2$ Matrix

$$\text{If } A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow A^{-1} = \frac{1}{\text{Det}(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \text{ provided } \text{Det}(A) \neq 0.$$

# Solving System of equations Using Laplace Transformation

**Step 1: Apply Laplace Transform (  $y(t) \rightarrow Y(s)$ ,  $x(t) \rightarrow X(s)$  )**


**Step 2: Find  $Y(s)$ ,  $X(s)$**

**Step 3: Apply Inverse Laplace Transform (  $Y(s) \rightarrow y(t)$ ,  $X(s) \rightarrow x(t)$  )**

## Problem

Solve the following system:

$$\begin{cases} \frac{dx}{dt} + y = \sin t \\ \frac{dy}{dt} + x = \cos t \end{cases}; \quad x(0) = 2, y(0) = 0.$$



## Solution:

$$\mathcal{L}\{x'(t) + y\} = \mathcal{L}\{\sin t\}$$

$$\mathcal{L}\{y'(t) + x\} = \mathcal{L}\{\cos t\}$$

let,

$$\mathcal{L}\{y(t)\} = Y(s), \quad \mathcal{L}\{x(t)\} = X(s)$$

$$\mathcal{L}\{y'(t)\} = sY(s) - y(0) = sY(s)$$

$$\mathcal{L}\{x'(t)\} = sX(s) - x(0) = sX(s) - 2$$

$$sX(s) - 2 + Y(s) = \frac{1}{s^2 + 1}$$

$$\therefore sY(s) + X(s) = \frac{s}{s^2 + 1}$$

$$5x(s) + 3y(s) = 2 + \frac{1}{s^2+1}$$

$\Rightarrow$

$$x(s) + 3y(s) = \frac{s}{s^2+1}$$

$$\Rightarrow \begin{pmatrix} s & 1 \\ 1 & s \end{pmatrix} \begin{pmatrix} x(s) \\ y(s) \end{pmatrix} = \begin{pmatrix} \frac{2s^2+3}{s^2+1} \\ \frac{s}{s^2+1} \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} x(s) \\ y(s) \end{pmatrix} = \begin{pmatrix} s & 1 \\ 1 & s \end{pmatrix}^{-1} \begin{pmatrix} \frac{2s^2+3}{s^2+1} \\ \frac{s}{s^2+1} \end{pmatrix}$$

$$= \frac{1}{s^2-1} \begin{pmatrix} s & -1 \\ -1 & s \end{pmatrix} \begin{pmatrix} \frac{2s^2+3}{s^2+1} \\ \frac{s}{s^2+1} \end{pmatrix}$$

$$\begin{pmatrix} x(s) \\ y(s) \end{pmatrix} = \begin{pmatrix} \frac{2s(s^2+1)}{(s^2+1)(s^2-1)} \\ \frac{-s(s^2+3)}{(s^2+1)(s^2-1)} \end{pmatrix}$$

$$\Rightarrow x(s) = \frac{2s(s^2+1)}{(s^2+1)(s^2-1)} \Rightarrow x(t) = e^{-t} \left\{ \frac{2s(s^2+1)}{(s^2+1)(s^2-1)} \right\}$$

$$y(s) = \frac{-s(s^2+3)}{(s^2+1)(s^2-1)} \Rightarrow y(t) = e^{-t} \left\{ \frac{-s(s^2+3)}{(s^2+1)(s^2-1)} \right\}$$

$$\Rightarrow x(t) = e^{-t} + e^t$$

$$\& y(t) = \frac{1}{2} (e^t - e^{-t}) - 3 \sin t \quad (\text{Ans:})$$





**Department of Mathematics and Natural Sciences**  
**MAT215 : Complex Variables and Laplace Transformations**  
**Laplace Transformation Formula Sheet**

### Laplace Transform definition

$$\mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

### Linearity of Laplace transformation

$$\mathcal{L}\{\alpha f(t) + \beta g(t)\} = \alpha \mathcal{L}\{f(t)\} + \beta \mathcal{L}\{g(t)\}$$

### Laplace Transforms of Elementary Functions (Real $s$ )

Function $f(t)$	$\mathcal{L}\{f(t)\} = F(s)$	Function $f(t)$	$\mathcal{L}\{f(t)\} = F(s)$
1	$\frac{1}{s}, \quad s > 0$	$\sin(at)$	$\frac{a}{s^2 + a^2}, \quad s > 0$
$t^n, n \in \mathbb{Z}^+$	$\frac{n!}{s^{n+1}}, \quad s > 0$	$\cos(at)$	$\frac{s}{s^2 + a^2}, \quad s > 0$
$t^n, n \notin \mathbb{Z}$	$\frac{\Gamma(n+1)}{s^{n+1}}, \quad s > 0$	$\sinh(at)$	$\frac{a}{s^2 - a^2}, \quad s >  a $
$e^{at}$	$\frac{1}{s-a}, \quad s > a$	$\cosh(at)$	$\frac{s}{s^2 - a^2}, \quad s >  a $

### 1st Translation theorem

$$\mathcal{L}\{e^{at} f(t)\} = \mathcal{L}\{f(t)\}|_{s \rightarrow s-a} \text{ for any real number } a.$$

### Laplace transformation of the form $t^n f(t)$

$$\mathcal{L}\{t^n f(t)\} = (-1)^n \frac{d^n}{ds^n} [\mathcal{L}\{f(t)\}]$$

## Piecewise function representation through unit step function

Suppose that,

$$h(t) = \begin{cases} f(t); & 0 \leq t < a \\ g(t); & t \geq a \end{cases}$$

Then,

$$h(t) = f(t)(1 - u(t - a)) + g(t)u(t - a)$$

## 2nd Translation Theorem

$$\mathcal{L}\{f(t-a)u(t-a)\} = \mathcal{L}\{f(t)\}e^{-as} \text{ for any positive value of } a.$$

## Inverse Laplace Transforms of some known expressions

$F(s)$	$\mathcal{L}^{-1}\{F(s)\} = f(t)$	$F(s)$	$\mathcal{L}^{-1}\{F(s)\} = f(t)$
$\frac{1}{s}$	1	$\frac{a}{s^2 + a^2}$	$\sin(at)$
$\frac{n!}{s^{n+1}}, n \in \mathbb{Z}^+$	$t^n$	$\frac{s}{s^2 + a^2}$	$\cos(at)$
$\frac{\Gamma(n+1)}{s^{n+1}}, n \notin \mathbb{Z}$	$t^n$	$\frac{a}{s^2 - a^2}$	$\sinh(at)$
$\frac{1}{s-a}$	$e^{at}$	$\frac{s}{s^2 - a^2}$	$\cosh(at)$

## Linearity of Inverse Laplace transformation

$$\mathcal{L}^{-1}\{\alpha F(s) + \beta G(s)\} = \alpha \mathcal{L}^{-1}\{F(s)\} + \beta \mathcal{L}^{-1}\{G(s)\}$$

## First Translation Theorem in inverse form

$$\mathcal{L}^{-1}\{F(s)|_{s \rightarrow s-a}\} = e^{at} \mathcal{L}^{-1}\{F(s)\}$$

## Second Translation Theorem in inverse form

$$\mathcal{L}^{-1}\{e^{-as}F(s)\} = \mathcal{L}^{-1}\{F(s)\}|_{t \rightarrow t-a} u(t-a)$$

## Laplace transformation of derivatives

If  $\mathcal{L}\{f(t)\} = F(s)$ , then

$$\mathcal{L}\{f'(t)\} = s \cdot F(s) - f(0)$$

$$\mathcal{L}\{f''(t)\} = s^2 \cdot F(s) - s \cdot f(0) - f'(0)$$

$$\mathcal{L}\{f'''(t)\} = s^3 \cdot F(s) - s^2 \cdot f(0) - s \cdot f'(0) - f''(0)$$