

Large N_c

Su HOUNG Lee



Theme:

1. Large N_c limit
2. Tetraquarks
3. Witten Veneziano formula

Ref:


- E. Witten, Nucl. Phys. B 160 (1980) 57
- E. Witten, Nucl. Phys. B 156 (1979) 213
- G. Veneziano, Nucl. Phys B 159 (1979) 213
- S. Weinberg, PRL110 (2013) 261601

II Feynman Diagram for Large N_c -1

1. QCD is a $SU(N)$ gauge theory with quarks in the fundamental representation

$$L = -\frac{1}{4} \text{Tr} \left[\left(\partial_\mu A_\nu - \partial_\nu A_\mu - ig [A_\mu, A_\nu] \right)^2 \right] + \bar{\psi} (i \not{D} - m) \psi$$



$$(A_\mu)_j^i = A_\mu^a \frac{\lambda_{ij}^a}{2}, \quad \psi_j, \bar{\psi}^i \quad i = 1, \dots, N_c$$



II Feynman Diagram for Large N_c -2

1. Counting of coupling g : quantum correction should have a smooth large N_c limit

$$(A_\mu)_i^j = A_\mu^a \frac{\lambda_{ij}^a}{2}, \quad \psi_j, \bar{\psi}^i \quad i = 1, \dots, N_c$$



Total counting: $g^2 N_c \approx O(1)$

$$g \rightarrow 1/N_c^{1/2}$$

Rule 1 : count total coupling $n \quad g^n \approx 1/N_c^{n/2}$


Rule 2 : additional factors can be obtained by drawing everything in terms of quark lines

II Feynman Diagram for Large N_c-3


(1)




(2)




(5)




(3)




(4)



(6)




(7)




Planar diagrams survive in the large N_c limit $8 \rightarrow 10$


(8)





(9)




(10)




(11)




(12)





(13)



(14)




(15)




quark loops are suppress compared to gluon loop (11)


(8)





(9)




(10)




(11)




(12)




(13)



(14)




(15)




quark loops are suppress compared to gluon loop (11)


(8)





(9)




(10)




(11)




(12)





(13)



(14)



(15)



quark lines are only at the boundary

(11)



(12)



(13)




(14)




(15)



(16)



(17)




III Properties of mesons in large N_c limit

Assumption: confinement persists in the large N_c limit

- Then can show mesons are free, stable, non interacting. Meson mass has smooth limit and the number of mesons are infinity. \rightarrow to prove this, look at two point function

$$P(k) = \int dx e^{ikx} \langle J(x), J(0) \rangle, \quad J = \bar{\psi} \psi$$



Only one meson state can contribute: $\bar{\psi}^i A_i^j A_j^k \psi_k$

$P(k)$: should be of order N_c for any k



$\langle J | N \rangle$ should be of order $N_c^{1/2}$

m_n^2 should be of order 1

There should be infinite number of mesons

$$P(k) = \sum_n \frac{|\langle J | n \rangle|^2}{k^2 - m_n^2} \xrightarrow{k \rightarrow \text{large}} \ln k^2$$

III Properties of mesons in large N_c limit: Decay



(22)

$$\langle JJJ \rangle = \sum \text{[triangle diagram with external lines at vertices]} + \sum \text{[triangle diagram with external lines at midpoints]}$$


- Using color decomposition as before, can show that only single meson states can interpolate

First term give : $|\langle J | n \rangle|^3 \Gamma_{nnn} \rightarrow \left(N_c^{1/2}\right)^3 \frac{1}{N_c^{1/2}}$


Therefore meson decay $\Gamma_{A \rightarrow BC} \propto 1/N_c$

III Properties of mesons in large N_c limit: elastic scattering

(28)




(29)



III Properties of mesons in large N_c limit: Glueball



Assumption: composed of two gluons $G_{\mu\nu}G^{\mu\nu}$

$$\langle G^2, G^2 \rangle = \sum \frac{|\langle G^2 | n \rangle|^2}{k^2 - m_G^2}$$



- Then can show glueballs are free, stable and non interacting and infinite in number. \rightarrow

Amplitude for glue to mix with meson: $1/N_c^{1/2}$



$$\langle J | n \rangle C_{nG} \langle G | J_G \rangle \propto N_c^{1/2} C_{nG} N_c = (1/N_c^{1/2})^4 N_c^3 \rightarrow C_{nG} = 1/N_c^{1/2}$$


Amplitude for k mesons and 1 glue state: $N_c^{-l - \frac{k}{2} + 1}$

IV Attractiveness of large N_c limit

- *Mesons are $q\bar{q}$ states $qqqq$ such as quark sea are suppressed*
- *Zweig rule suppression : decay from $q\bar{q}$ creation is suppressed*
- *Octet singlet degeneracy: contribution from disconnected diagrams are suppressed*

V Baryon in the large N_c limit


- Baryon mass is of order N_c*



$$N_c^2 \times \frac{1}{N_c}$$


VI Baryon in the large N_c limit : Scattering

- Baryon Baryon interaction



VI Baryon-Meson Scattering

- *Baryon-Meson*



$$N_c \times \frac{1}{N_c} \times 1$$

Tetraquarks in Large N_c


S. Weinberg, PRL110 (2013) 261601

Original argument why there is no Tetraquark

- Tetraquark in quark-antiquark basis

$$Q = c_1 (\bar{q} \Gamma q) (\bar{q} \Gamma q) + c_2 (\bar{q} \Gamma \lambda^a q) (\bar{q} \Gamma \lambda^a q)$$

$$\langle Q(x), Q(0) \rangle = \underbrace{\langle B(x), B(0) \rangle \langle B(x), B(0) \rangle}_{N_c^2} + \underbrace{\langle B(x) B(x), B(0) B(0) \rangle}_{N_c}_{connected}$$




Tetraquark lives here

So what? Tetraquark – current coupling

- Tetraquark current coupling

$$\langle 0 | Q(x) | T \rangle \propto N_c^{1/2}$$

$$Q(x) = B(x)B(x)$$







Tetraquark – meson+meson decay width

- Tetraquark current coupling

$$Q(x) = B(x)B(x)$$

$$\langle 0 | Q(x)B(y)B(z) | 0 \rangle = \langle 0 | B(x)B(y) | 0 \rangle \langle 0 | B(x)B(z) | 0 \rangle + \langle 0 | Q(x)B(y)B(z) | 0 \rangle_{\text{connected}}$$



Hence Tetraquark-meson-meson coupling $N_c^{-1/2}$

→ Decay width is N_c^{-1}

η' mass


E. Witten, Nucl. Phys. B 156 (1979) 213

G. Veneziano, Nucl. Phys B 159 (1979) 213


η' mass? Witten-Veneziano formula - I

- Correlation function $P(k) = -i \int dx e^{ikx} \langle G\tilde{G}(x), G\tilde{G}(0) \rangle$
- Contributions from glue only $P_0(k=0) \neq 0$ from low energy theorem
- When massless quarks are added $P(k) = -i \int dx e^{ikx} \langle \partial^\mu j_\mu^5(x), \partial^\mu j_\mu^5(0) \rangle \propto k^\mu k^\nu P_{\mu\nu} \xrightarrow{k=0} 0$

- Large N_c argument
$$P(k) = \sum_{\text{glueballs}} \frac{\langle 0 | G\tilde{G} | \text{glueball} \rangle^2}{k^2 - m_n^2} + \sum_{\text{mesons}} \frac{\langle 0 | G\tilde{G} | \text{meson} \rangle^2}{k^2 - m_n^2}$$



N_c^2



N_c

- Need η' meson
$$+ \frac{\langle 0 | G\tilde{G} | \eta' \rangle^2}{k^2 - m_{\eta'}^2} \quad \text{with } m_{\eta'}^2 \approx O\left(\frac{1}{N_c}\right)$$

$$\rightarrow P(k=0) = \boxed{P_0(0)} - \frac{\langle 0 | G\tilde{G} | \eta' \rangle^2}{m_{\eta'}^2} = 0$$

Witten-Veneziano formula – II

- η' meson
$$\frac{\langle 0 | G\tilde{G} | \eta' \rangle^2}{m_{\eta'}^2} = P_0(0)$$

$$\frac{\left(\frac{4\pi}{\alpha}\right)^2 \left(\frac{1}{N_F} \sqrt{N_F} m_{\eta'}^2 f_{\eta'}\right)^2}{m_{\eta'}^2} = \left(\frac{4\pi}{3\alpha}\right)^2 \frac{8}{11N/3} \left\langle \frac{\alpha}{\pi} G^2 \right\rangle$$

Lee, Zahed (01)

$$m_{\eta'}^2 f_{\eta'}^2 = \frac{8}{11N} \left\langle \frac{\alpha}{\pi} G^2 \right\rangle \rightarrow 250 \text{ MeV} < m_{\eta'} < 432 \text{ MeV} \quad \text{at } m \rightarrow 0 \text{ limit}$$

Should be related to $m_{\eta'}(958) - m_{\eta}(547) = 411 \text{ MeV}$

Few Formula in Large N_c

- *Meson*

$$m(1), \quad \Gamma(1/N_c), \quad g_{mmm}(1/N_c^{1/2}), \quad \langle 0 | \bar{q} \Gamma q | m \rangle (N_c^{1/2}), \quad \langle 0 | GG | m \rangle (N_c^{1/2})$$

- *Glueball*

$$m(1), \quad \Gamma(1/N_c^2), \quad g_{ggg}(1/N_c), \quad \langle 0 | \bar{q} \Gamma q | m \rangle (N_c), \quad \langle 0 | GG | g \rangle (N_c)$$

- *Baryon*

$$m(N_c), \quad g_{mBB}(N_c^{1/2}), \quad \langle B | \bar{q} \Gamma q | B \rangle (N_c), \quad \langle B | GG | B \rangle (N_c)$$

Witten-Veneziano formula – III N_c counting and glueball

- η' meson

$$\begin{array}{l}
 O(N_c^1) \rightarrow \frac{\langle 0 | G \tilde{G} | \eta' \rangle^2}{m_{\eta'}^2} = P_0(0) = \left(\frac{4\pi}{3\alpha} \right)^2 \frac{8}{11N/3} \left\langle \frac{\alpha}{\pi} G^2 \right\rangle \leftarrow O(N_c^2) \\
 O(1/N_c^1) \rightarrow
 \end{array}$$

η' mass is a large $1/N_c$ correction

- glueball

$$\begin{array}{l}
 O(N_c^2) \rightarrow \frac{\langle 0 | GG | g \rangle^2}{m_g^2} = S_0(0) = \left(\frac{4\pi}{3\alpha} \right)^2 \frac{18}{11N/3} \left\langle \frac{\alpha}{\pi} G^2 \right\rangle \leftarrow O(N_c^2) \\
 O(1) \rightarrow
 \end{array}$$

Witten-Veneziano formula – IV

- *Low energy theorem is a Non-perturbative effect*

$$S(q) = i \int dx e^{iqx} \left\langle \frac{3\alpha}{4\pi} G^2(x) \frac{3\alpha}{4\pi} G^2(0) \right\rangle = \frac{18}{11} \left\langle \frac{\alpha}{\pi} G^2 \right\rangle$$

$$P(q) = i \int dx e^{iqx} \left\langle \frac{3\alpha}{4\pi} G\tilde{G}(x) \frac{3\alpha}{4\pi} G\tilde{G}(0) \right\rangle = -\frac{8}{11} \left\langle \frac{\alpha}{\pi} G^2 \right\rangle$$

η' mass is a large $1/N_c$ correction

Witten-Veneziano formula at finite T (Kwon, Morita, Wolf, Lee: PRD 12)

- Large N_c counting $P(k) = i \int dx e^{ikx} \langle G\tilde{G}(x), G\tilde{G}(0) \rangle_m$



$$N_c^2$$

$$N_c^2$$


$$N_c$$

- At finite temperature, only gluonic effect is important


$$P(k) = \underbrace{\sum_{\text{glueballs}} \frac{\langle 0 | G\tilde{G} | \text{glueball} \rangle^2}{k^2 - m_n^2}}_{\text{Glue } N_c^2} + \underbrace{\sum_{\text{mesons}} \frac{\langle 0 | G\tilde{G} | \text{meson} \rangle^2}{k^2 - m_n^2}}_{\text{Quark } N_c} + \underbrace{\text{Scattering Term}}_{\text{Quark } N_c^2 \text{ ?}}$$

$$\rightarrow P(k=0) = P_0(0) - \frac{\langle 0 | G\tilde{G} | \eta' \rangle_c^2}{m_{\eta'}^2} + \text{scattering ?}$$

- *Large N_c argument for Nucleon Scattering Term*




Witten




That is, scattering terms are of order N_c and can be safely neglected

$$\rightarrow \frac{\langle n | G\tilde{G} | n \rangle^2}{m_N} \rho_{\text{density}} \qquad N_c^2 \left(\frac{1}{N_c} \right) = N_c$$

- *Large N_c argument for Meson Scattering Term*



Witten



$$N_c^2 \left(\frac{1}{N_c^{1/2}} \right)^2 \left(\frac{1}{N_c^{1/2}} \right)^2 = 1$$

That is, scattering terms are of order 1 and can be safely neglected

$$\rightarrow P_0(0) = \frac{\langle 0 | G\tilde{G} | \eta' \rangle^2}{m_{\eta'}^2}$$

WV relation remains the same


- *LET (Novikov, Shifman, Vainshtein, Zhakarov) at finite temperature : Ellis, Kapusta, Tang (98)*

$$\frac{d}{d(-1/4g_0^2)} \langle Op \rangle = -i \int dx e^{ikx} \langle Op(x), g_0^2 GG(0) \rangle$$

$$\langle Op \rangle_T = \text{const} \left[M_0 \exp \left(-\frac{8\pi^2}{bg_0^2} \right) \right]^d + c' T^d = \langle Op \rangle_{T_0} + c' T^d$$

$$\frac{d}{d(-1/4g_0^2)} \langle Op \rangle_T = \frac{32\pi^2}{b} \left(d - T \frac{\partial}{\partial T} \right) \langle Op \rangle_T = \frac{32\pi^2}{b} \left(d - T \frac{\partial}{\partial T} \right) \langle Op \rangle_{T_0}$$

- *Lee, Zahed (2001) $P_0(0) = -\frac{2}{b} \left(d - T \frac{\partial}{\partial T} \right) \left\langle \frac{\alpha}{\pi} G^2 \right\rangle$*



→ Morita et al. (2012)

Weak T dependence even near T_c

$$\rightarrow P_0(0) = \frac{\langle 0 | G \tilde{G} | \eta' \rangle_c^2}{m_{\eta'}^2}$$

- $\langle 0 | G\tilde{G} | \eta' \rangle$ at finite temperature

$$P(k) = \int d^4x e^{ikx} \left[\langle G\tilde{G}(x), G\tilde{G}(0) \rangle \right] = \frac{\langle 0 | G\tilde{G} | \eta' \rangle^2}{k^2 - m_\eta^2} + \dots$$

$$= k^\mu k^\nu \int d^4x e^{ikx} \left(\frac{4\pi}{\alpha N_F} \right)^2 \left[\langle \bar{q}(x) i\gamma_\mu \gamma^5 q(x), \bar{q}(0) i\gamma_\nu \gamma^5 q(0) \rangle - \langle \bar{q}(x) i\gamma_\mu q(x), \bar{q}(0) i\gamma_\nu q(0) \rangle \right]$$

chiral sym restored phase \rightarrow

$$= k^\mu k^\nu \int d^4x e^{ikx} \left(\frac{4\pi}{\alpha N_F} \right)^2 \left[\langle \bar{q}(x) i\gamma_\mu \gamma^5 \tau^a q(x), \bar{q}(0) i\gamma_\nu \gamma^5 \tau^a q(0) \rangle - \langle \bar{q}(x) i\gamma_\mu \tau^a q(x), \bar{q}(0) i\gamma_\nu \tau^a q(0) \rangle \right]$$

$= 0$ for any k^μ , when Chiral symmetry is restored

Therefore, $\langle 0 | G\tilde{G} | \eta' \rangle \rightarrow 0$ when chiral symmetry gets restored

- *W-V formula at finite temperature:*

$$\begin{array}{c}
 \langle \bar{q}q \rangle^2 \\
 \uparrow \\
 \frac{\langle 0 | G\tilde{G} | \eta' \rangle^2}{m_{\eta'}^2} = P_0(0) \longrightarrow \left(\frac{4\pi}{3\alpha} \right)^2 \frac{2}{11} \left(d - T \frac{\partial}{\partial T} \right) \left\langle \frac{\alpha}{\pi} G^2 \right\rangle
 \end{array}$$


Smooth temperature dependence even near T_c

Therefore , $m_{\eta'} - m_{\eta} \propto \langle \bar{q}q \rangle$


→ eta' mass should decrease at finite temperature

Experimental evidence of property change of η' in matter ?

CBELSA/TAPS coll



$$V = -(37 \pm 10 \pm 10) \text{ MeV} + i(10 \pm 2.5) \text{ MeV}$$



10 % reduction of mass from around 400 MeV from chiral symmetry breaking

Summary

1. η' correlation functions should exhibit symmetry breaking from N-point function in $SU(N)$ flavor even when chiral symmetry is restored.
→ For $SU(2)$, $UA(1)$ effect will be broken in the two point function

2. In W-V formula η' mass is related to quark condensate and thus should reduce at finite temperature independent of flavor **due to chiral symmetry restoration**
→
 - a) Could serve as signature of chiral symmetry restoration
 - b) Dilepton in Heavy Ion collision
 - c) Measurements from nuclear targets seems to support it ?

Summary

1. Chiral symmetry breaking in Correlator

$$\langle \bar{q}(0)q(0) \rangle \text{ or } \langle VV - AA \rangle \xrightarrow{m \rightarrow 0} \langle \psi_{\lambda=0}^+ \psi_{\lambda=0} \times \text{form factor} \rangle$$

2. $U_A(1)$ breaking effects in Correlators

$$\langle \eta' \eta' - \sigma\sigma \rangle \xrightarrow{m \rightarrow 0} \langle \psi_{\lambda=0}^+ \psi_{\lambda=0} \times \text{form factor} \rangle + m^{N_f-2} (\text{zero mode})$$

→ Restored in $SU(3)$ and real world

3. WV formula suggest mass of η' reduces in medium and at finite temperature: due to **chiral symmetry restoration**
4. Renewed interest in Theory and Experiments both for nuclear matter and at may be at finite T