

# Assignment 1

Simon Etter, 2020

Deadline: TODO

Total marks: 20

## 1 Floating-point numbers [4 marks]

Answer the following questions using pen-and-paper-style analysis [2 marks each].

1. What is the smallest positive integer which cannot be represented exactly in `Float64`?
2. What is the largest positive and finite integer which can be represented exactly in `Float64`?

*Hint.* You can check your answers using Julia.

## 2 Big O notation [4 marks]

Consider the statement

$$(1 + O(\varepsilon))^{-1} = 1 + O(\varepsilon) \quad \text{for } \varepsilon \rightarrow 0.$$

1. [2 marks] Verify this statement using the definition

$$f(x) = O(g(x)) \text{ for } x \rightarrow x_0 \iff \lim_{x \rightarrow x_0} \frac{|f(x)|}{|g(x)|} < \infty.$$

2. [2 marks] Verify this statement using the definition

$$f(x) = O(g(x)) \text{ for } x \rightarrow x_0 \iff \exists \delta, C > 0 \forall x \in x_0 + [-\delta, \delta] : |f(x)| \leq C |g(x)|.$$

## 3 The `expm1` function [8 marks]

Consider the function  $f(x) = \exp(x) - 1$  and its numerical implementation  $\tilde{f}(x) = \mathbf{T}(\exp(x)) \ominus 1$  (we ignore the rounding of the input of  $\tilde{f}(x)$  for simplicity).

1. [2 marks] Show that  $\kappa(f, x)$  is bounded for all  $x \in [-1, 1]$ .
2. [2 marks] Complete `expm1_errors()` such that it produces a plot demonstrating that

$$\frac{|\tilde{f}(x) - f(x)|}{|f(x)|} = O(x^{-1}) \quad \text{for } x \rightarrow 0.$$

3. [2 marks] Determine  $x \in [-1, 1]$  such that  $\frac{|\tilde{f}(x) - f(x)|}{|f(x)|} = 1$ . Demonstrate analytically that your choice of  $x$  indeed leads to a relative error of 1.
4. [1 mark] Determine functions  $g(y)$ ,  $h(x)$  such that  $f(x) = g(h(x))$  and at least one of the two functions is ill conditioned for  $x \approx 0$ . Compute the condition number of the ill-conditioned function.

5. [1 mark] Explain in your own words why Julia provides a special function `expm1(x)` for evaluating  $f(x) = \exp(x) - 1$ .
6. [unmarked] Extend `expm1_errors()` such that it also shows the relative errors for  $\tilde{f}(x) = \text{expm1}(x)$ .

## 4 Rounding errors when solving linear systems [4 marks]

Assume  $\tilde{x} \in \mathbb{R}^n$  is a numerically computed approximation to the solution of the linear system  $Ax = b$  where  $A \in \mathbb{R}^{n \times n}$  and  $x, b \in \mathbb{R}^n$ .

1. [3 marks] Determine a constant  $C \in \mathbb{R}$  depending only on the singular values  $\sigma_1 \geq \dots \geq \sigma_n > 0$  of  $A$  and  $\|x\|_2, \|b\|_2$  such that

$$\frac{\|\tilde{x} - x\|_2}{\|x\|_2} = C O(\text{eps}()).$$

You may assume that the numerical solution  $\tilde{x}$  is computed using a backward stable algorithm, i.e.  $\tilde{x}$  is the solution to  $\tilde{A}\tilde{x} = b$  where  $\tilde{A} \in \mathbb{R}^{n \times n}$  satisfies

$$\frac{\|\tilde{A} - A\|_2}{\|A\|_2} = O(\text{eps}()).$$

2. [1 mark] Complete the function `linear_system_error()` to check your answer to Task 1.  
*Hint.* You can compute the singular values of **A** using `svdvals(A)`, and you can compute the 2-norm of a vector **x** using `norm(x)`. The estimated relative error may be one or two order of magnitudes larger than the exact relative error (order of magnitude = power of 10).