

Example 8.2

Confidence Interval Estimates

OpenStax
Introductory Statistics 2e

Example


Suppose scores on exams in statistics are normally distributed with an unknown population mean and a population standard deviation of three points. A random sample of 36 scores is taken and gives a sample mean of 68. Find a confidence interval estimate for the population mean exam score.

$$\bar{X} - Z_{\alpha} \left(\frac{\sigma}{\sqrt{n}} \right) \leq \mu \leq \bar{X} + Z_{\alpha} \left(\frac{\sigma}{\sqrt{n}} \right)$$

$$\bar{X} = 68, \sigma = 3, n = 36, Z_{0.05} \text{ from calculator}$$


Example

Suppose scores on exams in statistics are normally distributed with an unknown population mean and a population standard deviation of three points. A random sample of 36 scores is taken and gives a sample mean of 68. Find a confidence interval estimate for the population mean exam score.



Example

Suppose scores on exams in statistics are normally distributed with an unknown population mean and a population standard deviation of three points. A random sample of 36 scores is taken and gives a sample mean of 68. Find a confidence interval estimate for the population mean exam score.



Example

Suppose scores on exams in statistics are normally distributed with an unknown population mean and a population standard deviation of three points. A random sample of 36 scores is taken and gives a sample mean of 68. Find a confidence interval estimate for the population mean exam score.

$$\bar{X} - Z_{\alpha} \left(\frac{\sigma}{\sqrt{n}} \right) \leq \mu \leq \bar{X} + Z_{\alpha} \left(\frac{\sigma}{\sqrt{n}} \right)$$

$$\bar{X} = 68, \sigma = 3, n = 36, Z_{0.05} \text{ from calculator}$$

$$68 - 1.645 \left(\frac{3}{\sqrt{36}} \right) \leq \mu \leq 68 + 1.645 \left(\frac{3}{\sqrt{36}} \right)$$

Example

Suppose scores on exams in statistics are normally distributed with an unknown population mean and a population standard deviation of three points. A random sample of 36 scores is taken and gives a sample mean of 68. Find a confidence interval estimate for the population mean exam score.

$$\bar{X} - Z_{\alpha} \left(\frac{\sigma}{\sqrt{n}} \right) \leq \mu \leq \bar{X} + Z_{\alpha} \left(\frac{\sigma}{\sqrt{n}} \right)$$

$$\bar{X} = 68, \sigma = 3, n = 36, Z_{0.05} \text{ from calculator}$$

$$68 - 1.645 \left(\frac{3}{\sqrt{36}} \right) \leq \mu \leq 68 + 1.645 \left(\frac{3}{\sqrt{36}} \right)$$

$$68 - 0.822 \leq \mu \leq 68 + 0.822$$

Example

Suppose scores on exams in statistics are normally distributed with an unknown population mean and a population standard deviation of three points. A random sample of 36 scores is taken and gives a sample mean of 68. Find a confidence interval estimate for the population mean exam score.

$$\bar{X} - Z_{\alpha} \left(\frac{\sigma}{\sqrt{n}} \right) \leq \mu \leq \bar{X} + Z_{\alpha} \left(\frac{\sigma}{\sqrt{n}} \right)$$

$$\bar{X} = 68, \sigma = 3, n = 36, Z_{0.05} \text{ from calculator}$$

$$68 - 1.645 \left(\frac{3}{\sqrt{36}} \right) \leq \mu \leq 68 + 1.645 \left(\frac{3}{\sqrt{36}} \right)$$

$$68 - 0.822 \leq \mu \leq 68 + 0.822$$

$$67.178 \leq \mu \leq 68.822$$

Example

Suppose scores on exams in statistics are normally distributed with an unknown population mean and a population standard deviation of three points. A random sample of 36 scores is taken and gives a sample mean of 68. Find a confidence interval estimate for the population mean exam score.

```
NORMAL FLOAT AUTO REAL RADIAN MP
invNorm(0.95,0,1)
.....1.644853626
Ans*3/√36
......822426813
68-Ans
.....67.17757319
68+.822426813
.....68.82242681
```