

Étale groupoid algebras

Benjamin Steinberg, City College of New York



August 1, 2014

Groups, Rings and Group Rings

Outline

Étale Groupoids

Étale Groupoid Algebras

Representation Theory

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).
- They simultaneously generalize:

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).
- They simultaneously generalize:
 - commutative C^* -algebras

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).
- They simultaneously generalize:
 - commutative C^* -algebras
 - group C^* -algebras

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).
- They simultaneously generalize:
 - commutative C^* -algebras
 - group C^* -algebras
 - group action C^* -algebras

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).
- They simultaneously generalize:
 - commutative C^* -algebras
 - group C^* -algebras
 - group action C^* -algebras
 - Cuntz-Krieger/graph C^* -algebras

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).
- They simultaneously generalize:
 - commutative C^* -algebras
 - group C^* -algebras
 - group action C^* -algebras
 - Cuntz-Krieger/graph C^* -algebras
 - inverse semigroup C^* -algebras.

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).
- They simultaneously generalize:
 - commutative C^* -algebras
 - group C^* -algebras
 - group action C^* -algebras
 - Cuntz-Krieger/graph C^* -algebras
 - inverse semigroup C^* -algebras.
- Algebraic properties can often be seen from the groupoid.

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).
- They simultaneously generalize:
 - commutative C^* -algebras
 - group C^* -algebras
 - group action C^* -algebras
 - Cuntz-Krieger/graph C^* -algebras
 - inverse semigroup C^* -algebras.
- Algebraic properties can often be seen from the groupoid.
- Morita equivalence of groupoid algebras is often explained by a Morita equivalence of the groupoids.

Background

- Groupoid C^* -algebras have been a fruitful and unifying notion in the theory of operator algebras since J. Renault's influential monograph (1980).
- They simultaneously generalize:
 - commutative C^* -algebras
 - group C^* -algebras
 - group action C^* -algebras
 - Cuntz-Krieger/graph C^* -algebras
 - inverse semigroup C^* -algebras.
- Algebraic properties can often be seen from the groupoid.
- Morita equivalence of groupoid algebras is often explained by a Morita equivalence of the groupoids.
- Groupoids are espoused by Connes as noncommutative models of spaces.

Algebraic analogues

- Many groupoid C^* -algebras have analogues in the context of associative algebras.

Algebraic analogues

- Many groupoid C^* -algebras have analogues in the context of associative algebras.
- Group algebras and inverse semigroup algebras are obvious examples.

Algebraic analogues

- Many groupoid C^* -algebras have analogues in the context of associative algebras.
- Group algebras and inverse semigroup algebras are obvious examples.
- **Leavitt path algebras** are algebraic analogues of Cuntz-Krieger C^* -algebras.

Algebraic analogues

- Many groupoid C^* -algebras have analogues in the context of associative algebras.
- Group algebras and inverse semigroup algebras are obvious examples.
- **Leavitt path algebras** are algebraic analogues of Cuntz-Krieger C^* -algebras.
- The ring of \mathbb{k} -valued continuous functions with compact support on a totally disconnected space is an algebraic analogue of a commutative C^* -algebra.

Algebraic analogues

- Many groupoid C^* -algebras have analogues in the context of associative algebras.
- Group algebras and inverse semigroup algebras are obvious examples.
- **Leavitt path algebras** are algebraic analogues of Cuntz-Krieger C^* -algebras.
- The ring of \mathbb{k} -valued continuous functions with compact support on a totally disconnected space is an algebraic analogue of a commutative C^* -algebra.
- Over a field, these are precisely the idempotent-generated commutative algebras.

Algebraic analogues

- Many groupoid C^* -algebras have analogues in the context of associative algebras.
- Group algebras and inverse semigroup algebras are obvious examples.
- **Leavitt path algebras** are algebraic analogues of Cuntz-Krieger C^* -algebras.
- The ring of \mathbb{k} -valued continuous functions with compact support on a totally disconnected space is an algebraic analogue of a commutative C^* -algebra.
- Over a field, these are precisely the idempotent-generated commutative algebras.
- Surprising similarities between operator algebras and their algebraic analogues have been known for some time.

History

- In 2009 [Adv. Math. '10], I introduced an algebraic analogue of groupoid C^* -algebras for a class of étale groupoids.

History

- In 2009 [Adv. Math. '10], I introduced an algebraic analogue of groupoid C^* -algebras for a class of étale groupoids.
- My hope was that it would explain many of the similarities between the algebraic and analytic setting, especially for Leavitt path algebras.

History

- In 2009 [Adv. Math. '10], I introduced an algebraic analogue of groupoid C^* -algebras for a class of étale groupoids.
- My hope was that it would explain many of the similarities between the algebraic and analytic setting, especially for Leavitt path algebras.
- Initially, I focused on applications to inverse semigroup algebras.

History

- In 2009 [Adv. Math. '10], I introduced an algebraic analogue of groupoid C^* -algebras for a class of étale groupoids.
- My hope was that it would explain many of the similarities between the algebraic and analytic setting, especially for Leavitt path algebras.
- Initially, I focused on applications to inverse semigroup algebras.
- Groupoid algebras over \mathbb{C} were rediscovered later by L. O. Clark, C. Farthing, A. Sims and M. Tomforde, who have kindly dubbed them “Steinberg algebras.”

History

- In 2009 [Adv. Math. '10], I introduced an algebraic analogue of groupoid C^* -algebras for a class of étale groupoids.
- My hope was that it would explain many of the similarities between the algebraic and analytic setting, especially for Leavitt path algebras.
- Initially, I focused on applications to inverse semigroup algebras.
- Groupoid algebras over \mathbb{C} were rediscovered later by L. O. Clark, C. Farthing, A. Sims and M. Tomforde, who have kindly dubbed them “Steinberg algebras.”
- Many of my hopes have since been borne out by J. Brown, L. O. Clark, C. Farthing, A. Sims and M. Tomforde, who seem to produce new results faster than I can keep up with.

Groupoids

- A **groupoid** \mathcal{G} is a small category in which all morphisms are isomorphisms.

Groupoids

- A **groupoid** \mathcal{G} is a small category in which all morphisms are isomorphisms.
- $\mathcal{G}^{(0)}$ is the **object** set.

Groupoids

- A **groupoid** \mathcal{G} is a small category in which all morphisms are isomorphisms.
- $\mathcal{G}^{(0)}$ is the **object** set.
- $\mathcal{G}^{(1)}$ is the **arrow** set.

Groupoids

- A **groupoid** \mathcal{G} is a small category in which all morphisms are isomorphisms.
- $\mathcal{G}^{(0)}$ is the **object** set.
- $\mathcal{G}^{(1)}$ is the **arrow** set.
- $d, r: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ are the **domain** and **range** maps.

Groupoids

- A **groupoid** \mathcal{G} is a small category in which all morphisms are isomorphisms.
- $\mathcal{G}^{(0)}$ is the **object** set.
- $\mathcal{G}^{(1)}$ is the **arrow** set.
- $d, r: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ are the **domain** and **range** maps.
- $m: \mathcal{G}^{(2)} \rightarrow \mathcal{G}^{(1)}$ is the **multiplication** map.

Groupoids

- A **groupoid** \mathcal{G} is a small category in which all morphisms are isomorphisms.
- $\mathcal{G}^{(0)}$ is the **object** set.
- $\mathcal{G}^{(1)}$ is the **arrow** set.
- $d, r: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ are the **domain** and **range** maps.
- $m: \mathcal{G}^{(2)} \rightarrow \mathcal{G}^{(1)}$ is the **multiplication** map.
- $\iota: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(1)}$ is the **inversion** map.

Groupoids

- A **groupoid** \mathcal{G} is a small category in which all morphisms are isomorphisms.
- $\mathcal{G}^{(0)}$ is the **object** set.
- $\mathcal{G}^{(1)}$ is the **arrow** set.
- $d, r: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ are the **domain** and **range** maps.
- $m: \mathcal{G}^{(2)} \rightarrow \mathcal{G}^{(1)}$ is the **multiplication** map.
- $\iota: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(1)}$ is the **inversion** map.
- We view $\mathcal{G}^{(0)} \subseteq \mathcal{G}^{(1)}$ by identifying objects and identity arrows.

Groupoids

- A **groupoid** \mathcal{G} is a small category in which all morphisms are isomorphisms.
- $\mathcal{G}^{(0)}$ is the **object** set.
- $\mathcal{G}^{(1)}$ is the **arrow** set.
- $d, r: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ are the **domain** and **range** maps.
- $m: \mathcal{G}^{(2)} \rightarrow \mathcal{G}^{(1)}$ is the **multiplication** map.
- $\iota: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(1)}$ is the **inversion** map.
- We view $\mathcal{G}^{(0)} \subseteq \mathcal{G}^{(1)}$ by identifying objects and identity arrows.
- Often $\mathcal{G}^{(0)}$ is called the **unit space** of \mathcal{G} .

Étale groupoids

- A **topological groupoid** is a groupoid \mathcal{G} with a topology on $\mathcal{G}^{(1)}$ such that if $\mathcal{G}^{(0)}$ is given the induced topology, then all the structure maps are continuous.

Étale groupoids

- A **topological groupoid** is a groupoid \mathcal{G} with a topology on $\mathcal{G}^{(1)}$ such that if $\mathcal{G}^{(0)}$ is given the induced topology, then all the structure maps are continuous.
- **Étale groupoids** form one of the most important classes of topological groupoids.

Étale groupoids

- A **topological groupoid** is a groupoid \mathcal{G} with a topology on $\mathcal{G}^{(1)}$ such that if $\mathcal{G}^{(0)}$ is given the induced topology, then all the structure maps are continuous.
- **Étale groupoids** form one of the most important classes of topological groupoids.
- \mathcal{G} is **étale** if $d: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ is a local homeomorphism.

Étale groupoids

- A **topological groupoid** is a groupoid \mathcal{G} with a topology on $\mathcal{G}^{(1)}$ such that if $\mathcal{G}^{(0)}$ is given the induced topology, then all the structure maps are continuous.
- **Étale groupoids** form one of the most important classes of topological groupoids.
- \mathcal{G} is **étale** if $d: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ is a local homeomorphism.
- This forces all the structure maps to be local homeomorphisms and $\mathcal{G}^{(0)}$ to be open.

Étale groupoids

- A **topological groupoid** is a groupoid \mathcal{G} with a topology on $\mathcal{G}^{(1)}$ such that if $\mathcal{G}^{(0)}$ is given the induced topology, then all the structure maps are continuous.
- **Étale groupoids** form one of the most important classes of topological groupoids.
- \mathcal{G} is **étale** if $d: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ is a local homeomorphism.
- This forces all the structure maps to be local homeomorphisms and $\mathcal{G}^{(0)}$ to be open.
- We assume $\mathcal{G}^{(0)}$ is locally compact Hausdorff, but $\mathcal{G}^{(1)}$ need not be Hausdorff.

Étale groupoids

- A **topological groupoid** is a groupoid \mathcal{G} with a topology on $\mathcal{G}^{(1)}$ such that if $\mathcal{G}^{(0)}$ is given the induced topology, then all the structure maps are continuous.
- **Étale groupoids** form one of the most important classes of topological groupoids.
- \mathcal{G} is **étale** if $d: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ is a local homeomorphism.
- This forces all the structure maps to be local homeomorphisms and $\mathcal{G}^{(0)}$ to be open.
- We assume $\mathcal{G}^{(0)}$ is locally compact Hausdorff, but $\mathcal{G}^{(1)}$ need not be Hausdorff.
- Renault calls an étale groupoid \mathcal{G} **ample** if $\mathcal{G}^{(0)}$ has a basis of compact open sets.

Étale groupoids

- A **topological groupoid** is a groupoid \mathcal{G} with a topology on $\mathcal{G}^{(1)}$ such that if $\mathcal{G}^{(0)}$ is given the induced topology, then all the structure maps are continuous.
- **Étale groupoids** form one of the most important classes of topological groupoids.
- \mathcal{G} is **étale** if $d: \mathcal{G}^{(1)} \rightarrow \mathcal{G}^{(0)}$ is a local homeomorphism.
- This forces all the structure maps to be local homeomorphisms and $\mathcal{G}^{(0)}$ to be open.
- We assume $\mathcal{G}^{(0)}$ is locally compact Hausdorff, but $\mathcal{G}^{(1)}$ need not be Hausdorff.
- Renault calls an étale groupoid \mathcal{G} **ample** if $\mathcal{G}^{(0)}$ has a basis of compact open sets.
- Many important C^* -algebras come from ample groupoids: group algebras, Cuntz-Krieger algebras and inverse semigroup algebras.

Discrete groups

- A discrete group is the same thing as a one-object étale groupoid.

Discrete groups

- A discrete group is the same thing as a one-object étale groupoid.
- Indeed, if $\mathcal{G}^{(0)}$ is discrete, then because d is a local homeomorphism also $\mathcal{G}^{(1)}$ is discrete.

Discrete groups

- A discrete group is the same thing as a one-object étale groupoid.
- Indeed, if $\mathcal{G}^{(0)}$ is discrete, then because d is a local homeomorphism also $\mathcal{G}^{(1)}$ is discrete.
- More generally, any discrete groupoid is an étale groupoid.

Discrete groups

- A discrete group is the same thing as a one-object étale groupoid.
- Indeed, if $\mathcal{G}^{(0)}$ is discrete, then because d is a local homeomorphism also $\mathcal{G}^{(1)}$ is discrete.
- More generally, any discrete groupoid is an étale groupoid.
- In fact, discrete groups and groupoids are ample.

Spaces

- Let X be a locally compact Hausdorff space.

Spaces

- Let X be a locally compact Hausdorff space.
- Then X is an étale groupoid in which all arrows are identities.

Spaces

- Let X be a locally compact Hausdorff space.
- Then X is an étale groupoid in which all arrows are identities.
- That is, $\mathcal{G}^{(0)} = X = \mathcal{G}^{(1)}$.

Spaces

- Let X be a locally compact Hausdorff space.
- Then X is an étale groupoid in which all arrows are identities.
- That is, $\mathcal{G}^{(0)} = X = \mathcal{G}^{(1)}$.
- X is an ample groupoid if and only if it has a basis of compact open sets.

Spaces

- Let X be a locally compact Hausdorff space.
- Then X is an étale groupoid in which all arrows are identities.
- That is, $\mathcal{G}^{(0)} = X = \mathcal{G}^{(1)}$.
- X is an ample groupoid if and only if it has a basis of compact open sets.
- This guarantees X has enough continuous maps into any non-trivial discrete ring to separate points.

Group actions

- Let G be a discrete group acting on a locally compact Hausdorff space X .

Group actions

- Let G be a discrete group acting on a locally compact Hausdorff space X .
- The **semidirect product** $G \ltimes X$ has object space X and arrow space $G \times X$.

Group actions

- Let G be a discrete group acting on a locally compact Hausdorff space X .
- The **semidirect product** $G \ltimes X$ has object space X and arrow space $G \times X$.
- Think

$$(g, x) = gx \xleftarrow{g} x.$$

Group actions

- Let G be a discrete group acting on a locally compact Hausdorff space X .
- The **semidirect product** $G \ltimes X$ has object space X and arrow space $G \times X$.
- Think

$$(g, x) = gx \xleftarrow{g} x.$$

- Composition is

$$ghx \xleftarrow{g} hx \xleftarrow{h} x$$

Group actions

- Let G be a discrete group acting on a locally compact Hausdorff space X .
- The **semidirect product** $G \ltimes X$ has object space X and arrow space $G \times X$.
- Think

$$(g, x) = gx \xleftarrow{g} x.$$

- Composition is

$$ghx \xleftarrow{g} hx \xleftarrow{h} x$$

- Inversion is

$$gx \xleftarrow{g} x$$

Deaconu-Renault groupoid

- Let $Q = (Q^{(0)}, Q^{(1)})$ be a finite quiver.

Deaconu-Renault groupoid

- Let $Q = (Q^{(0)}, Q^{(1)})$ be a finite quiver.
- Q^ω is the space of infinite paths in Q .

Deaconu-Renault groupoid

- Let $Q = (Q^{(0)}, Q^{(1)})$ be a finite quiver.
- Q^ω is the space of infinite paths in Q .
- Let $T: Q^\omega \rightarrow Q^\omega$ be the **shift** map

$$T(e_0e_1e_2\cdots) = e_1e_2\cdots.$$

Deaconu-Renault groupoid

- Let $Q = (Q^{(0)}, Q^{(1)})$ be a finite quiver.
- Q^ω is the space of infinite paths in Q .
- Let $T: Q^\omega \rightarrow Q^\omega$ be the **shift** map

$$T(e_0e_1e_2\cdots) = e_1e_2\cdots.$$

- The **Deaconu-Renault** groupoid \mathcal{G}_Q has object space Q^ω and arrow space

$$\{x \xrightarrow{m-n} y \mid T^m(x) = T^n(y)\} \subseteq Q^\omega \times \mathbb{Z} \times Q^\omega.$$

Deaconu-Renault groupoid

- Let $Q = (Q^{(0)}, Q^{(1)})$ be a finite quiver.
- Q^ω is the space of infinite paths in Q .
- Let $T: Q^\omega \rightarrow Q^\omega$ be the **shift** map

$$T(e_0e_1e_2\cdots) = e_1e_2\cdots.$$

- The **Deaconu-Renault** groupoid \mathcal{G}_Q has object space Q^ω and arrow space

$$\{x \xrightarrow{m-n} y \mid T^m(x) = T^n(y)\} \subseteq Q^\omega \times \mathbb{Z} \times Q^\omega.$$

- Composition is

$$x \xrightarrow{m-n} y \xrightarrow{m'+n'} z = x \xrightarrow{(m+m')-(n+n')} z$$

Deaconu-Renault groupoid

- Let $Q = (Q^{(0)}, Q^{(1)})$ be a finite quiver.
- Q^ω is the space of infinite paths in Q .
- Let $T: Q^\omega \rightarrow Q^\omega$ be the **shift** map

$$T(e_0 e_1 e_2 \cdots) = e_1 e_2 \cdots .$$

- The **Deaconu-Renault** groupoid \mathcal{G}_Q has object space Q^ω and arrow space

$$\{x \xrightarrow{m-n} y \mid T^m(x) = T^n(y)\} \subseteq Q^\omega \times \mathbb{Z} \times Q^\omega.$$

- Composition is

$$x \xrightarrow{m-n} y \xrightarrow{m'+n'} z = x \xrightarrow{(m+m')-(n+n')} z$$

- $(x \xrightarrow{m-n} y)^{-1} = y \xrightarrow{n-m} x.$

Deaconu-Renault groupoid

- Let $Q = (Q^{(0)}, Q^{(1)})$ be a finite quiver.
- Q^ω is the space of infinite paths in Q .
- Let $T: Q^\omega \rightarrow Q^\omega$ be the **shift** map

$$T(e_0 e_1 e_2 \cdots) = e_1 e_2 \cdots$$

- The **Deaconu-Renault** groupoid \mathcal{G}_Q has object space Q^ω and arrow space

$$\{x \xrightarrow{m-n} y \mid T^m(x) = T^n(y)\} \subseteq Q^\omega \times \mathbb{Z} \times Q^\omega.$$

- Composition is

$$x \xrightarrow{m-n} y \xrightarrow{m'+n'} z = x \xrightarrow{(m+m')-(n+n')} z$$

- $(x \xrightarrow{m-n} y)^{-1} = y \xrightarrow{n-m} x$.
- This ample groupoid gives rise to Cuntz-Krieger C^* -algebras and Leavitt path algebras.

Inverse semigroups

- An **inverse semigroup** is a semigroup S such that, for all $s \in S$, there exists unique $s^* \in S$ such that

$$ss^*s = s \quad \text{and} \quad s^*ss^* = s^*.$$

Inverse semigroups

- An **inverse semigroup** is a semigroup S such that, for all $s \in S$, there exists unique $s^* \in S$ such that

$$ss^*s = s \quad \text{and} \quad s^*ss^* = s^*.$$

- Note that s^*s , ss^* are idempotents.

Inverse semigroups

- An **inverse semigroup** is a semigroup S such that, for all $s \in S$, there exists unique $s^* \in S$ such that

$$ss^*s = s \quad \text{and} \quad s^*ss^* = s^*.$$

- Note that s^*s , ss^* are idempotents.
- Groups are inverse semigroups.

Inverse semigroups

- An **inverse semigroup** is a semigroup S such that, for all $s \in S$, there exists unique $s^* \in S$ such that

$$ss^*s = s \quad \text{and} \quad s^*ss^* = s^*.$$

- Note that s^*s , ss^* are idempotents.
- Groups are inverse semigroups.
- The idempotents $E(S)$ commute and form a subsemigroup.

Inverse semigroups

- An **inverse semigroup** is a semigroup S such that, for all $s \in S$, there exists unique $s^* \in S$ such that

$$ss^*s = s \quad \text{and} \quad s^*ss^* = s^*.$$

- Note that s^*s , ss^* are idempotents.
- Groups are inverse semigroups.
- The idempotents $E(S)$ commute and form a subsemigroup.
- $E(S)$ is a **semilattice** with $e \wedge f = ef$.

Inverse semigroups

- An **inverse semigroup** is a semigroup S such that, for all $s \in S$, there exists unique $s^* \in S$ such that

$$ss^*s = s \quad \text{and} \quad s^*ss^* = s^*.$$

- Note that s^*s , ss^* are idempotents.
- Groups are inverse semigroups.
- The idempotents $E(S)$ commute and form a subsemigroup.
- $E(S)$ is a **semilattice** with $e \wedge f = ef$.
- If G is a group acting on a semilattice E , then the semidirect product $E \rtimes G$ is an inverse semigroup.

Pseudogroups of partial homeomorphisms

- Let X be a topological space.

Pseudogroups of partial homeomorphisms


- Let X be a topological space.
- I_X is the inverse semigroup of all homeomorphisms between open subsets of X .

Pseudogroups of partial homeomorphisms

- Let X be a topological space.
- I_X is the inverse semigroup of all homeomorphisms between open subsets of X .
- Composition is defined where it makes sense:


Pseudogroups of partial homeomorphisms

- Let X be a topological space.
- I_X is the inverse semigroup of all homeomorphisms between open subsets of X .
- Composition is defined where it makes sense:



Pseudogroups of partial homeomorphisms


- Let X be a topological space.
- I_X is the inverse semigroup of all homeomorphisms between open subsets of X .
- Composition is defined where it makes sense:



- The empty partial function is a zero element of I_X .

Pseudogroups of partial homeomorphisms


- Let X be a topological space.
- I_X is the inverse semigroup of all homeomorphisms between open subsets of X .
- Composition is defined where it makes sense:



- The empty partial function is a zero element of I_X .
- If $f: U \rightarrow V$, then $f^* = f^{-1}: V \rightarrow U$.

Pseudogroups of partial homeomorphisms

- Let X be a topological space.
- I_X is the inverse semigroup of all homeomorphisms between open subsets of X .
- Composition is defined where it makes sense:



- The empty partial function is a zero element of I_X .
- If $f: U \rightarrow V$, then $f^* = f^{-1}: V \rightarrow U$.
- Inverse semigroups abstract pseudogroups of partial homeomorphisms from differential geometry.

Actions of inverse semigroups

- S is an inverse semigroup.

Actions of inverse semigroups

- S is an inverse semigroup.
- X is a locally compact Hausdorff space.

Actions of inverse semigroups

- S is an inverse semigroup.
- X is a locally compact Hausdorff space.
- An action $S \curvearrowright X$ is a homomorphism $\theta: S \rightarrow I_X$ with

$$\bigcup_{e \in E(S)} \text{dom}(\theta(e)) = X \quad (1)$$

Actions of inverse semigroups

- S is an inverse semigroup.
- X is a locally compact Hausdorff space.
- An action $S \curvearrowright X$ is a homomorphism $\theta: S \rightarrow I_X$ with

$$\bigcup_{e \in E(S)} \text{dom}(\theta(e)) = X \quad (1)$$

- Since $\text{dom}(\theta(s)) = \text{dom}(\theta(s^*s))$, (1) is equivalent to X being the union of the domains of all elements of S .

Actions of inverse semigroups

- S is an inverse semigroup.
- X is a locally compact Hausdorff space.
- An action $S \curvearrowright X$ is a homomorphism $\theta: S \rightarrow I_X$ with

$$\bigcup_{e \in E(S)} \text{dom}(\theta(e)) = X \quad (1)$$

- Since $\text{dom}(\theta(s)) = \text{dom}(\theta(s^*s))$, (1) is equivalent to X being the union of the domains of all elements of S .
- If S is a group, (1) implies the action is by homeomorphisms.

Actions of inverse semigroups

- S is an inverse semigroup.
- X is a locally compact Hausdorff space.
- An action $S \curvearrowright X$ is a homomorphism $\theta: S \rightarrow I_X$ with

$$\bigcup_{e \in E(S)} \text{dom}(\theta(e)) = X \quad (1)$$

- Since $\text{dom}(\theta(s)) = \text{dom}(\theta(s^*s))$, (1) is equivalent to X being the union of the domains of all elements of S .
- If S is a group, (1) implies the action is by homeomorphisms.
- If $s \in S$ and $e \in E(S)$, then $\theta(se)$ is the restriction of $\theta(s)$ to $\text{dom}(\theta(e))$.

Groupoids of germs

- Suppose $S \curvearrowright X$ is an action of an inverse semigroup.

Groupoids of germs

- Suppose $S \curvearrowright X$ is an action of an inverse semigroup.
- We can form a **groupoid of germs** $S \ltimes X$.

Groupoids of germs

- Suppose $S \curvearrowright X$ is an action of an inverse semigroup.
- We can form a **groupoid of germs** $S \ltimes X$.
- X is the object space and $(S \times X)/\sim$ is the arrow space.

Groupoids of germs

- Suppose $S \curvearrowright X$ is an action of an inverse semigroup.
- We can form a **groupoid of germs** $S \ltimes X$.
- X is the object space and $(S \times X)/\sim$ is the arrow space.
- Think

$$[s, x] = sx \xleftarrow{s} x.$$

Groupoids of germs

- Suppose $S \curvearrowright X$ is an action of an inverse semigroup.
- We can form a **groupoid of germs** $S \ltimes X$.
- X is the object space and $(S \times X)/\sim$ is the arrow space.
- Think

$$[s, x] = sx \xleftarrow{s} x.$$

- Composition and inversion are like in the group action case.

Groupoids of germs

- Suppose $S \curvearrowright X$ is an action of an inverse semigroup.
- We can form a **groupoid of germs** $S \ltimes X$.
- X is the object space and $(S \times X)/\sim$ is the arrow space.
- Think

$$[s, x] = sx \xleftarrow{s} x.$$

- Composition and inversion are like in the group action case.
- $(s, x) \sim (t, y) \iff x = y \text{ and } \exists e \in E(S) \text{ with } x \in \text{dom}(e), se = te.$

Groupoids of germs

- Suppose $S \curvearrowright X$ is an action of an inverse semigroup.
- We can form a **groupoid of germs** $S \ltimes X$.
- X is the object space and $(S \times X)/\sim$ is the arrow space.
- Think

$$[s, x] = sx \xleftarrow{s} x.$$

- Composition and inversion are like in the group action case.
- $(s, x) \sim (t, y) \iff x = y$ and $\exists e \in E(S)$ with $x \in \text{dom}(e)$, $se = te$.
- $S \ltimes X$ is not Hausdorff in general.

Groupoids of germs

- Suppose $S \curvearrowright X$ is an action of an inverse semigroup.
- We can form a **groupoid of germs** $S \ltimes X$.
- X is the object space and $(S \times X)/\sim$ is the arrow space.
- Think

$$[s, x] = sx \xleftarrow{s} x.$$

- Composition and inversion are like in the group action case.
- $(s, x) \sim (t, y) \iff x = y$ and $\exists e \in E(S)$ with $x \in \text{dom}(e)$, $se = te$.
- $S \ltimes X$ is not Hausdorff in general.
- It is ample if X has a basis of compact open sets and the domains of elements of S are compact open.

Paterson's universal groupoid

- Let $\widehat{E(S)} \subseteq \{0, 1\}^{E(S)}$ be the space of non-zero homomorphisms (**characters**) $\chi: E(S) \rightarrow \{0, 1\}$.

Paterson's universal groupoid

- Let $\widehat{E(S)} \subseteq \{0, 1\}^{E(S)}$ be the space of non-zero homomorphisms (**characters**) $\chi: E(S) \rightarrow \{0, 1\}$.
- S acts on $\widehat{E(S)}$ by $s \cdot \chi(e) = \chi(s^*es)$.

Paterson's universal groupoid

- Let $\widehat{E(S)} \subseteq \{0, 1\}^{E(S)}$ be the space of non-zero homomorphisms (**characters**) $\chi: E(S) \rightarrow \{0, 1\}$.
- S acts on $\widehat{E(S)}$ by $s \cdot \chi(e) = \chi(s^*es)$.
- The domain of s consists of those characters with $s \cdot \chi \neq 0$.

Paterson's universal groupoid

- Let $\widehat{E(S)} \subseteq \{0, 1\}^{E(S)}$ be the space of non-zero homomorphisms (**characters**) $\chi: E(S) \rightarrow \{0, 1\}$.
- S acts on $\widehat{E(S)}$ by $s \cdot \chi(e) = \chi(s^*es)$.
- The domain of s consists of those characters with $s \cdot \chi \neq 0$.
- The **universal groupoid** $\mathcal{G}(S)$ is $S \ltimes \widehat{E(S)}$.

Paterson's universal groupoid

- Let $\widehat{E(S)} \subseteq \{0, 1\}^{E(S)}$ be the space of non-zero homomorphisms (**characters**) $\chi: E(S) \rightarrow \{0, 1\}$.
- S acts on $\widehat{E(S)}$ by $s \cdot \chi(e) = \chi(s^*es)$.
- The domain of s consists of those characters with $s \cdot \chi \neq 0$.
- The **universal groupoid** $\mathcal{G}(S)$ is $S \ltimes \widehat{E(S)}$.
- Paterson proved $C^*(\mathcal{G}(S)) \cong C^*(S)$.

Groupoid algebras

- Let \mathbb{k} be a unital comm. ring with the discrete topology.

Groupoid algebras

- Let \mathbb{k} be a unital comm. ring with the discrete topology.
- Let \mathcal{G} be an ample groupoid.

Groupoid algebras

- Let \mathbb{k} be a unital comm. ring with the discrete topology.
- Let \mathcal{G} be an ample groupoid.
- The **groupoid algebra** $\mathbb{k}\mathcal{G}$ is defined as follows.

Groupoid algebras

- Let \mathbb{k} be a unital comm. ring with the discrete topology.
- Let \mathcal{G} be an ample groupoid.
- The **groupoid algebra** $\mathbb{k}\mathcal{G}$ is defined as follows.
- The underlying vector space consists of the continuous \mathbb{k} -valued functions on \mathcal{G} with compact support.

Groupoid algebras

- Let \mathbb{k} be a unital comm. ring with the discrete topology.
- Let \mathcal{G} be an ample groupoid.
- The **groupoid algebra** $\mathbb{k}\mathcal{G}$ is defined as follows.
- The underlying vector space consists of the continuous \mathbb{k} -valued functions on \mathcal{G} with compact support.



Warning

↪ This definition needs adjustment for the non-Hausdorff case.

Groupoid algebras

- Let \mathbb{k} be a unital comm. ring with the discrete topology.
- Let \mathcal{G} be an ample groupoid.
- The **groupoid algebra** $\mathbb{k}\mathcal{G}$ is defined as follows.
- The underlying vector space consists of the continuous \mathbb{k} -valued functions on \mathcal{G} with compact support.
- Ample implies there are many such functions.

Groupoid algebras

- Let \mathbb{k} be a unital comm. ring with the discrete topology.
- Let \mathcal{G} be an ample groupoid.
- The **groupoid algebra** $\mathbb{k}\mathcal{G}$ is defined as follows.
- The underlying vector space consists of the continuous \mathbb{k} -valued functions on \mathcal{G} with compact support.
- Ample implies there are many such functions.
- The product is convolution:

$$f * g(x) = \sum_{d(x)=d(z)} f(xz^{-1})g(z).$$

Groupoid algebras

- Let \mathbb{k} be a unital comm. ring with the discrete topology.
- Let \mathcal{G} be an ample groupoid.
- The **groupoid algebra** $\mathbb{k}\mathcal{G}$ is defined as follows.
- The underlying vector space consists of the continuous \mathbb{k} -valued functions on \mathcal{G} with compact support.
- Ample implies there are many such functions.
- The product is convolution:

$$f * g(x) = \sum_{d(x)=d(z)} f(xz^{-1})g(z).$$

- The sum is finite because fibers of d are closed and discrete and f, g have compact support.

Examples

- If G is a discrete group, $\mathbb{k}G$ is the usual group algebra and similarly for discrete groupoids.

Examples

- If G is a discrete group, $\mathbb{k}G$ is the usual group algebra and similarly for discrete groupoids.
- If X is a space, $\mathbb{k}X$ is the ring of continuous \mathbb{k} -valued functions with compact support.

Examples

- If G is a discrete group, $\mathbb{k}G$ is the usual group algebra and similarly for discrete groupoids.
- If X is a space, $\mathbb{k}X$ is the ring of continuous \mathbb{k} -valued functions with compact support.
- The Leavitt path algebras are the algebras of Deaconu-Renault groupoids.

Examples

- If G is a discrete group, $\mathbb{k}G$ is the usual group algebra and similarly for discrete groupoids.
- If X is a space, $\mathbb{k}X$ is the ring of continuous \mathbb{k} -valued functions with compact support.
- The Leavitt path algebras are the algebras of Deaconu-Renault groupoids.
- If $G \curvearrowright X$ is a group action, $\mathbb{k}(G \ltimes X)$ is the cross product $\mathbb{k}X \rtimes G$.

Examples

- If G is a discrete group, $\mathbb{k}G$ is the usual group algebra and similarly for discrete groupoids.
- If X is a space, $\mathbb{k}X$ is the ring of continuous \mathbb{k} -valued functions with compact support.
- The Leavitt path algebras are the algebras of Deaconu-Renault groupoids.
- If $G \curvearrowright X$ is a group action, $\mathbb{k}(G \ltimes X)$ is the cross product $\mathbb{k}X \rtimes G$.

Theorem (BS)

Let S be an inverse semigroup and \mathbb{k} a comm. ring with 1. Then $\mathbb{k}S \cong \mathbb{k}\mathcal{G}(S)$.

First properties

- $C^*(\mathcal{G})$ is the completion of $\mathbb{C}\mathcal{G}$ (Stone-Weierstrass).

First properties

- $C^*(\mathcal{G})$ is the completion of $\mathbb{C}\mathcal{G}$ (Stone-Weierstrass).
- This lets us recover $C^*(S) \cong C^*(\mathcal{G}(S))$ for an inverse semigroup S .

First properties

- $C^*(\mathcal{G})$ is the completion of $\mathbb{C}\mathcal{G}$ (Stone-Weierstrass).
- This lets us recover $C^*(S) \cong C^*(\mathcal{G}(S))$ for an inverse semigroup S .
- $\mathbb{k}\mathcal{G}$ is unital iff \mathcal{G}_0 is compact.

First properties

- $C^*(\mathcal{G})$ is the completion of $\mathbb{C}\mathcal{G}$ (Stone-Weierstrass).
- This lets us recover $C^*(S) \cong C^*(\mathcal{G}(S))$ for an inverse semigroup S .
- $\mathbb{k}\mathcal{G}$ is unital iff \mathcal{G}_0 is compact.
- $\mathbb{k}\mathcal{G}$ has local units (is a directed union of unital subrings).

First properties

- $C^*(\mathcal{G})$ is the completion of $\mathbb{C}\mathcal{G}$ (Stone-Weierstrass).
- This lets us recover $C^*(S) \cong C^*(\mathcal{G}(S))$ for an inverse semigroup S .
- $\mathbb{k}\mathcal{G}$ is unital iff \mathcal{G}_0 is compact.
- $\mathbb{k}\mathcal{G}$ has local units (is a directed union of unital subrings).
- Abrams, Ahn and Marki developed a successful Morita theory for rings with local units.

First properties

- $C^*(\mathcal{G})$ is the completion of $\mathbb{C}\mathcal{G}$ (Stone-Weierstrass).
- This lets us recover $C^*(S) \cong C^*(\mathcal{G}(S))$ for an inverse semigroup S .
- $\mathbb{k}\mathcal{G}$ is unital iff \mathcal{G}_0 is compact.
- $\mathbb{k}\mathcal{G}$ has local units (is a directed union of unital subrings).
- Abrams, Ahn and Marki developed a successful Morita theory for rings with local units.
- $\mathbb{k}\mathcal{G}$ carries an involution $f^*(g) = f(g^{-1})$.

Isotropy

- The **isotropy group** G_x of $x \in \mathcal{G}^{(0)}$ consists of all loops $x \xrightarrow{g} x$.

Isotropy

- The **isotropy group** G_x of $x \in \mathcal{G}^{(0)}$ consists of all loops $x \xrightarrow{g} x$.
- The **isotropy bundle** $\mathcal{G}_{iso} = \bigcup_{x \in X} G_x$.

Isotropy

- The **isotropy group** G_x of $x \in \mathcal{G}^{(0)}$ consists of all loops $x \xrightarrow{g} x$.
- The **isotropy bundle** $\mathcal{G}_{iso} = \bigcup_{x \in X} G_x$.
- \mathcal{G} is **effective** if $\text{Int}(\mathcal{G}_{iso}) = \mathcal{G}^{(0)}$.

Isotropy

- The **isotropy group** G_x of $x \in \mathcal{G}^{(0)}$ consists of all loops $x \xrightarrow{g} x$.
- The **isotropy bundle** $\mathcal{G}_{iso} = \bigcup_{x \in X} G_x$.
- \mathcal{G} is **effective** if $\text{Int}(\mathcal{G}_{iso}) = \mathcal{G}^{(0)}$.
- The Deaconu-Renault groupoid is effective iff the Leavitt path algebra satisfies condition (L).

Isotropy

- The **isotropy group** G_x of $x \in \mathcal{G}^{(0)}$ consists of all loops $x \xrightarrow{g} x$.
- The **isotropy bundle** $\mathcal{G}_{iso} = \bigcup_{x \in X} G_x$.
- \mathcal{G} is **effective** if $\text{Int}(\mathcal{G}_{iso}) = \mathcal{G}^{(0)}$.
- The Deaconu-Renault groupoid is effective iff the Leavitt path algebra satisfies condition (L).
- $G \ltimes X$ is effective iff $G \curvearrowright X$ is topologically free.

Isotropy

- The **isotropy group** G_x of $x \in \mathcal{G}^{(0)}$ consists of all loops $x \xrightarrow{g} x$.
- The **isotropy bundle** $\mathcal{G}_{iso} = \bigcup_{x \in X} G_x$.
- \mathcal{G} is **effective** if $\text{Int}(\mathcal{G}_{iso}) = \mathcal{G}^{(0)}$.
- The Deaconu-Renault groupoid is effective iff the Leavitt path algebra satisfies condition (L).
- $G \ltimes X$ is effective iff $G \curvearrowright X$ is topologically free.
- $f \in \mathbb{k}\mathcal{G}$ is a **class function** if:

Isotropy

- The **isotropy group** G_x of $x \in \mathcal{G}^{(0)}$ consists of all loops $x \xrightarrow{g} x$.
- The **isotropy bundle** $\mathcal{G}_{iso} = \bigcup_{x \in X} G_x$.
- \mathcal{G} is **effective** if $\text{Int}(\mathcal{G}_{iso}) = \mathcal{G}^{(0)}$.
- The Deaconu-Renault groupoid is effective iff the Leavitt path algebra satisfies condition (L).
- $G \ltimes X$ is effective iff $G \curvearrowright X$ is topologically free.
- $f \in \mathbb{k}\mathcal{G}$ is a **class function** if:
 1. $\text{supp}(f) \subseteq \mathcal{G}_{iso}$

Isotropy

- The **isotropy group** G_x of $x \in \mathcal{G}^{(0)}$ consists of all loops $x \xrightarrow{g} x$.
- The **isotropy bundle** $\mathcal{G}_{iso} = \bigcup_{x \in X} G_x$.
- \mathcal{G} is **effective** if $\text{Int}(\mathcal{G}_{iso}) = \mathcal{G}^{(0)}$.
- The Deaconu-Renault groupoid is effective iff the Leavitt path algebra satisfies condition (L).
- $G \ltimes X$ is effective iff $G \curvearrowright X$ is topologically free.
- $f \in \mathbb{k}\mathcal{G}$ is a **class function** if:
 1. $\text{supp}(f) \subseteq \mathcal{G}_{iso}$
 2. $\begin{array}{c} h \\ \circlearrowleft \\ \bullet \xrightarrow{g} \bullet \end{array}$ implies $f(ghg^{-1}) = f(h)$:

Isotropy

- The **isotropy group** G_x of $x \in \mathcal{G}^{(0)}$ consists of all loops $x \xrightarrow{g} x$.
- The **isotropy bundle** $\mathcal{G}_{iso} = \bigcup_{x \in X} G_x$.
- \mathcal{G} is **effective** if $\text{Int}(\mathcal{G}_{iso}) = \mathcal{G}^{(0)}$.
- The Deaconu-Renault groupoid is effective iff the Leavitt path algebra satisfies condition (L).
- $G \ltimes X$ is effective iff $G \curvearrowright X$ is topologically free.
- $f \in \mathbb{k}\mathcal{G}$ is a **class function** if:
 1. $\text{supp}(f) \subseteq \mathcal{G}_{iso}$
 2. $\begin{array}{c} h \\ \circlearrowleft \\ \bullet \xrightarrow{g} \bullet \end{array}$ implies $f(ghg^{-1}) = f(h)$:

Theorem (BS)

The class functions form the center of $\mathbb{k}\mathcal{G}$.

Orbits and minimality

- The **orbit** \mathcal{O}_x of $x \in \mathcal{G}^{(0)}$ consists of all y such that there exists $x \xrightarrow{g} y$.

Orbits and minimality

- The **orbit** \mathcal{O}_x of $x \in \mathcal{G}^{(0)}$ consists of all y such that there exists $x \xrightarrow{g} y$.
- If $G \curvearrowright X$ is a group action, then the orbits of $G \ltimes X$ are the orbits of G on X .

Orbits and minimality

- The **orbit** \mathcal{O}_x of $x \in \mathcal{G}^{(0)}$ consists of all y such that there exists $x \xrightarrow{g} y$.
- If $G \curvearrowright X$ is a group action, then the orbits of $G \ltimes X$ are the orbits of G on X .
- \mathcal{G} is **minimal** if every orbit is dense.

Orbits and minimality

- The **orbit** \mathcal{O}_x of $x \in \mathcal{G}^{(0)}$ consists of all y such that there exists $x \xrightarrow{g} y$.
- If $G \curvearrowright X$ is a group action, then the orbits of $G \ltimes X$ are the orbits of G on X .
- \mathcal{G} is **minimal** if every orbit is dense.

Theorem (BS)

Let \mathcal{G} be an effective, Hausdorff ample groupoid with a dense orbit. Then

$$Z(\mathbb{k}\mathcal{G}) = \begin{cases} \mathbb{k}, & \text{if } \mathcal{G}^{(0)} \text{ is compact} \\ 0, & \text{else.} \end{cases}$$

Orbits and minimality

- The **orbit** \mathcal{O}_x of $x \in \mathcal{G}^{(0)}$ consists of all y such that there exists $x \xrightarrow{g} y$.
- If $G \curvearrowright X$ is a group action, then the orbits of $G \ltimes X$ are the orbits of G on X .
- \mathcal{G} is **minimal** if every orbit is dense.

Theorem (BS)

Let \mathcal{G} be an effective, Hausdorff ample groupoid with a dense orbit. Then

$$Z(\mathbb{k}\mathcal{G}) = \begin{cases} \mathbb{k}, & \text{if } \mathcal{G}^{(0)} \text{ is compact} \\ 0, & \text{else.} \end{cases}$$

- This generalizes an earlier result of L. O. Clark and C. Edie-Michelle for the minimal case.

Simplicity

Theorem (L. O. Clark, C. Edie-Michelle)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field. Then $\mathbb{k}\mathcal{G}$ is simple if and only if \mathcal{G} is effective and minimal.

Simplicity

Theorem (L. O. Clark, C. Edie-Michelle)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field. Then $\mathbb{k}\mathcal{G}$ is simple if and only if \mathcal{G} is effective and minimal.

- First proved by J. H. Brown, L. O. Clark, C. Farthing and A. Sims over \mathbb{C} .

Simplicity

Theorem (L. O. Clark, C. Edie-Michelle)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field. Then $\mathbb{k}\mathcal{G}$ is simple if and only if \mathcal{G} is effective and minimal.

- First proved by J. H. Brown, L. O. Clark, C. Farthing and A. Sims over \mathbb{C} .
- Translates into simplicity criterion for Leavitt path algebras.

Simplicity

Theorem (L. O. Clark, C. Edie-Michelle)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field. Then $\mathbb{k}\mathcal{G}$ is simple if and only if \mathcal{G} is effective and minimal.

- First proved by J. H. Brown, L. O. Clark, C. Farthing and A. Sims over \mathbb{C} .
- Translates into simplicity criterion for Leavitt path algebras.
- I can use to characterize inverse semigroups S with simple contracted semigroup algebras provided $\mathcal{G}(S)$ is Hausdorff.

Simplicity

Theorem (L. O. Clark, C. Edie-Michelle)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field. Then $\mathbb{k}\mathcal{G}$ is simple if and only if \mathcal{G} is effective and minimal.

- First proved by J. H. Brown, L. O. Clark, C. Farthing and A. Sims over \mathbb{C} .
- Translates into simplicity criterion for Leavitt path algebras.
- I can use to characterize inverse semigroups S with simple contracted semigroup algebras provided $\mathcal{G}(S)$ is Hausdorff.
- This answers in part a question of Munn.

Simplicity

Theorem (L. O. Clark, C. Edie-Michelle)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field. Then $\mathbb{k}\mathcal{G}$ is simple if and only if \mathcal{G} is effective and minimal.

- First proved by J. H. Brown, L. O. Clark, C. Farthing and A. Sims over \mathbb{C} .
- Translates into simplicity criterion for Leavitt path algebras.
- I can use to characterize inverse semigroups S with simple contracted semigroup algebras provided $\mathcal{G}(S)$ is Hausdorff.
- This answers in part a question of Munn.
- A sufficient condition for Hausdorff is being 0-E-unitary:

Simplicity

Theorem (L. O. Clark, C. Edie-Michelle)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field. Then $\mathbb{k}\mathcal{G}$ is simple if and only if \mathcal{G} is effective and minimal.

- First proved by J. H. Brown, L. O. Clark, C. Farthing and A. Sims over \mathbb{C} .
- Translates into simplicity criterion for Leavitt path algebras.
- I can use to characterize inverse semigroups S with simple contracted semigroup algebras provided $\mathcal{G}(S)$ is Hausdorff.
- This answers in part a question of Munn.
- A sufficient condition for Hausdorff is being 0-E-unitary:
 - $se, e \in E(S) \setminus \{0\} \implies s \in E(S)$.

Primitivity

- Recall: a ring is **primitive** if it has a faithful simple module.

Primitivity

- Recall: a ring is **primitive** if it has a faithful simple module.
- Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field.

Primitivity

- Recall: a ring is **primitive** if it has a faithful simple module.
- Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field.

Theorem (BS)

If \mathcal{G} is effective, then $\mathbb{k}\mathcal{G}$ is primitive iff \mathcal{G} has a dense orbit.

Primitivity

- Recall: a ring is **primitive** if it has a faithful simple module.
- Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field.

Theorem (BS)

If \mathcal{G} is effective, then $\mathbb{k}\mathcal{G}$ is primitive iff \mathcal{G} has a dense orbit.

Theorem (BS)

Suppose that \mathcal{G} has a dense orbit \mathcal{O}_x with $\mathbb{k}G_x$ primitive. Then $\mathbb{k}\mathcal{G}$ is primitive. The converse holds if x is isolated.

Primitivity

- Recall: a ring is **primitive** if it has a faithful simple module.
- Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field.

Theorem (BS)

If \mathcal{G} is effective, then $\mathbb{k}\mathcal{G}$ is primitive iff \mathcal{G} has a dense orbit.

Theorem (BS)

Suppose that \mathcal{G} has a dense orbit \mathcal{O}_x with $\mathbb{k}G_x$ primitive. Then $\mathbb{k}\mathcal{G}$ is primitive. The converse holds if x is isolated.

- Recall that G_x is the isotropy group at x .

Primitivity

- Recall: a ring is **primitive** if it has a faithful simple module.
- Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field.

Theorem (BS)

If \mathcal{G} is effective, then $\mathbb{k}\mathcal{G}$ is primitive iff \mathcal{G} has a dense orbit.

Theorem (BS)

Suppose that \mathcal{G} has a dense orbit \mathcal{O}_x with $\mathbb{k}G_x$ primitive. Then $\mathbb{k}\mathcal{G}$ is primitive. The converse holds if x is isolated.

- Recall that G_x is the isotropy group at x .
- A dense orbit is always necessary for primitivity.

Primitivity

- Recall: a ring is **primitive** if it has a faithful simple module.
- Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a field.

Theorem (BS)

If \mathcal{G} is effective, then $\mathbb{k}\mathcal{G}$ is primitive iff \mathcal{G} has a dense orbit.

Theorem (BS)

Suppose that \mathcal{G} has a dense orbit \mathcal{O}_x with $\mathbb{k}G_x$ primitive. Then $\mathbb{k}\mathcal{G}$ is primitive. The converse holds if x is isolated.

- Recall that G_x is the isotropy group at x .
- A dense orbit is always necessary for primitivity.
- There is a non-Hausdorff version.

Semiprimitivity: the effective case

- Recall: a ring is **semiprimitive** if it has a faithful semisimple module.

Semiprimitivity: the effective case

- Recall: a ring is **semiprimitive** if it has a faithful semisimple module.

Theorem (BS)

Let \mathcal{G} be a Hausdorff effective ample groupoid and \mathbb{k} a semiprimitive comm. ring with 1. Then $\mathbb{k}\mathcal{G}$ is semiprimitive.

Semiprimitivity

Theorem (BS)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a semiprimitive comm. ring with 1. If $\mathbb{k}G_x$ is semiprimitive for all x in some dense subset $X \subseteq \mathcal{G}^{(0)}$, then $\mathbb{k}\mathcal{G}$ is semiprimitive. The converse holds if X consists of isolated points.

Semiprimitivity

Theorem (BS)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a semiprimitive comm. ring with 1. If $\mathbb{k}G_x$ is semiprimitive for all x in some dense subset $X \subseteq \mathcal{G}^{(0)}$, then $\mathbb{k}\mathcal{G}$ is semiprimitive. The converse holds if X consists of isolated points.

- A non-Hausdorff version holds.

Semiprimitivity

Theorem (BS)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a semiprimitive comm. ring with 1. If $\mathbb{k}G_x$ is semiprimitive for all x in some dense subset $X \subseteq \mathcal{G}^{(0)}$, then $\mathbb{k}\mathcal{G}$ is semiprimitive. The converse holds if X consists of isolated points.

- A non-Hausdorff version holds.
- Isotropy groups for Deaconu-Renault groupoids are trivial or \mathbb{Z} , recovering semiprimitivity of Leavitt path algebras.

Semiprimitivity

Theorem (BS)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a semiprimitive comm. ring with 1. If $\mathbb{k}G_x$ is semiprimitive for all x in some dense subset $X \subseteq \mathcal{G}^{(0)}$, then $\mathbb{k}\mathcal{G}$ is semiprimitive. The converse holds if X consists of isolated points.

- A non-Hausdorff version holds.
- Isotropy groups for Deaconu-Renault groupoids are trivial or \mathbb{Z} , recovering semiprimitivity of Leavitt path algebras.

Corollary

Let \mathcal{G} be an ample groupoid and \mathbb{k} be a field of characteristic 0 such that \mathbb{k}/\mathbb{Q} is not algebraic. Then $\mathbb{k}\mathcal{G}$ is semiprimitive.

Semiprimitivity

Theorem (BS)

Let \mathcal{G} be a Hausdorff ample groupoid and \mathbb{k} a semiprimitive comm. ring with 1. If $\mathbb{k}G_x$ is semiprimitive for all x in some dense subset $X \subseteq \mathcal{G}^{(0)}$, then $\mathbb{k}\mathcal{G}$ is semiprimitive. The converse holds if X consists of isolated points.

- A non-Hausdorff version holds.
- Isotropy groups for Deaconu-Renault groupoids are trivial or \mathbb{Z} , recovering semiprimitivity of Leavitt path algebras.

Corollary

Let \mathcal{G} be an ample groupoid and \mathbb{k} be a field of characteristic 0 such that \mathbb{k}/\mathbb{Q} is not algebraic. Then $\mathbb{k}\mathcal{G}$ is semiprimitive.

Proof.

Apply Amitsur '59 with $X = \mathcal{G}^{(0)}$.



Applications to inverse semigroup algebras

Corollary

Let \mathbb{k} be a semiprimitive comm. ring with 1 and S an inverse semigroup. If $\mathbb{k}G_e$ is semiprimitive for every maximal subgroup G_e of S , then $\mathbb{k}S$ is semiprimitive. The converse holds if $E(S)$ is pseudofinite.

Applications to inverse semigroup algebras

Corollary

Let \mathbb{k} be a semiprimitive comm. ring with 1 and S an inverse semigroup. If $\mathbb{k}G_e$ is semiprimitive for every maximal subgroup G_e of S , then $\mathbb{k}S$ is semiprimitive. The converse holds if $E(S)$ is pseudofinite.

- Sufficiency was proved by Domanov (1976).

Applications to inverse semigroup algebras

Corollary

Let \mathbb{k} be a semiprimitive comm. ring with 1 and S an inverse semigroup. If $\mathbb{k}G_e$ is semiprimitive for every maximal subgroup G_e of S , then $\mathbb{k}S$ is semiprimitive. The converse holds if $E(S)$ is pseudofinite.

- Sufficiency was proved by Domanov (1976).
- The converse was proved by Munn (1987).

Applications to inverse semigroup algebras

Corollary

Let \mathbb{k} be a semiprimitive comm. ring with 1 and S an inverse semigroup. If $\mathbb{k}G_e$ is semiprimitive for every maximal subgroup G_e of S , then $\mathbb{k}S$ is semiprimitive. The converse holds if $E(S)$ is pseudofinite.

- Sufficiency was proved by Domanov (1976).
- The converse was proved by Munn (1987).
- G_e is the group of units of eSe for e idempotent.

Applications to inverse semigroup algebras

Corollary

Let \mathbb{k} be a semiprimitive comm. ring with 1 and S an inverse semigroup. If $\mathbb{k}G_e$ is semiprimitive for every maximal subgroup G_e of S , then $\mathbb{k}S$ is semiprimitive. The converse holds if $E(S)$ is pseudofinite.

- Sufficiency was proved by Domanov (1976).
- The converse was proved by Munn (1987).
- G_e is the group of units of eSe for e idempotent.
- $E(S)$ is pseudofinite if the characteristic functions of principal filters are discrete in $\widehat{E(S)}$.

Applications to inverse semigroup algebras

Corollary

Let \mathbb{k} be a semiprimitive comm. ring with 1 and S an inverse semigroup. If $\mathbb{k}G_e$ is semiprimitive for every maximal subgroup G_e of S , then $\mathbb{k}S$ is semiprimitive. The converse holds if $E(S)$ is pseudofinite.

- Sufficiency was proved by Domanov (1976).
- The converse was proved by Munn (1987).
- G_e is the group of units of eSe for e idempotent.
- $E(S)$ is pseudofinite if the characteristic functions of principal filters are discrete in $\widehat{E(S)}$.
- This is a topological reformulation of the original definition.

Unitary modules

- Let R be a ring with local units.

Unitary modules

- Let R be a ring with local units.
- An R -module M is **unitary** if $RM = M$.

Unitary modules

- Let R be a ring with local units.
- An R -module M is **unitary** if $RM = M$.
- The category of unitary R -modules is denoted **$R\text{-mod}$** .

Unitary modules

- Let R be a ring with local units.
- An R -module M is **unitary** if $RM = M$.
- The category of unitary R -modules is denoted **R -mod**.
- R and S are **Morita equivalent** if R -mod is equivalent to S -mod.

Unitary modules

- Let R be a ring with local units.
- An R -module M is **unitary** if $RM = M$.
- The category of unitary R -modules is denoted $R\text{-mod}$.
- R and S are **Morita equivalent** if $R\text{-mod}$ is equivalent to $S\text{-mod}$.
- Morita equivalence can also be formulated in terms of Morita contexts.

Sheaves of \mathcal{G} -modules

- Let \mathcal{G} be an ample groupoid and \mathbb{k} a comm. ring with 1.

Sheaves of \mathcal{G} -modules

- Let \mathcal{G} be an ample groupoid and \mathbb{k} a comm. ring with 1.
- A \mathcal{G} -sheaf of \mathbb{k} -modules is a sheaf of \mathbb{k} -modules over $\mathcal{G}^{(0)}$ with an action of \mathcal{G} .

Sheaves of \mathcal{G} -modules

- Let \mathcal{G} be an ample groupoid and \mathbb{k} a comm. ring with 1.
- A \mathcal{G} -sheaf of \mathbb{k} -modules is a sheaf of \mathbb{k} -modules over $\mathcal{G}^{(0)}$ with an action of \mathcal{G} .
- Roughly, this means that if $x \xrightarrow{g} y$ is in $\mathcal{G}^{(1)}$, then g takes the stalk over x to the stalk over y by a \mathbb{k} -module isomorphism.

Sheaves of \mathcal{G} -modules

- Let \mathcal{G} be an ample groupoid and \mathbb{k} a comm. ring with 1.
- A \mathcal{G} -sheaf of \mathbb{k} -modules is a sheaf of \mathbb{k} -modules over $\mathcal{G}^{(0)}$ with an action of \mathcal{G} .
- Roughly, this means that if $x \xrightarrow{g} y$ is in $\mathcal{G}^{(1)}$, then g takes the stalk over x to the stalk over y by a \mathbb{k} -module isomorphism.
- There are also some continuity conditions.

Sheaves of \mathcal{G} -modules

- Let \mathcal{G} be an ample groupoid and \mathbb{k} a comm. ring with 1.
- A \mathcal{G} -sheaf of \mathbb{k} -modules is a sheaf of \mathbb{k} -modules over $\mathcal{G}^{(0)}$ with an action of \mathcal{G} .
- Roughly, this means that if $x \xrightarrow{g} y$ is in $\mathcal{G}^{(1)}$, then g takes the stalk over x to the stalk over y by a \mathbb{k} -module isomorphism.
- There are also some continuity conditions.
- If \mathcal{G} is discrete, this amounts to a functor $\mathcal{G} \rightarrow \mathbb{k}\text{-mod}$.

Sheaves of \mathcal{G} -modules

- Let \mathcal{G} be an ample groupoid and \mathbb{k} a comm. ring with 1.
- A \mathcal{G} -sheaf of \mathbb{k} -modules is a sheaf of \mathbb{k} -modules over $\mathcal{G}^{(0)}$ with an action of \mathcal{G} .
- Roughly, this means that if $x \xrightarrow{g} y$ is in $\mathcal{G}^{(1)}$, then g takes the stalk over x to the stalk over y by a \mathbb{k} -module isomorphism.
- There are also some continuity conditions.
- If \mathcal{G} is discrete, this amounts to a functor $\mathcal{G} \rightarrow \mathbb{k}\text{-mod}$.

Theorem (BS)

$\mathbb{k}\mathcal{G}\text{-mod}$ is equivalent to the category of \mathcal{G} -sheaves of \mathbb{k} -modules.

Morita equivalence

- If Z is a locally compact space and $f: Z \rightarrow \mathcal{G}^{(0)}$ is continuous, there is a pullback groupoid $\mathcal{G}[Z]$.

Morita equivalence

- If Z is a locally compact space and $f: Z \rightarrow \mathcal{G}^{(0)}$ is continuous, there is a pullback groupoid $\mathcal{G}[Z]$.
- $\mathcal{G}[Z]^{(0)} = Z$, $\mathcal{G}[Z]^{(1)} = \{z \xrightarrow{g} z' \mid f(z) \xrightarrow{g} f(z')\}$.

Morita equivalence

- If Z is a locally compact space and $f: Z \rightarrow \mathcal{G}^{(0)}$ is continuous, there is a pullback groupoid $\mathcal{G}[Z]$.
- $\mathcal{G}[Z]^{(0)} = Z$, $\mathcal{G}[Z]^{(1)} = \{z \xrightarrow{g} z' \mid f(z) \xrightarrow{g} f(z')\}$.
- Groupoids \mathcal{G} and \mathcal{H} are **Morita equivalent** if there is a locally compact space Z and continuous open surjections $p: Z \rightarrow \mathcal{G}^{(0)}$ and $q: Z \rightarrow \mathcal{H}^{(0)}$ such that $\mathcal{G}[Z] \cong \mathcal{H}[Z]$.

Morita equivalence

- If Z is a locally compact space and $f: Z \rightarrow \mathcal{G}^{(0)}$ is continuous, there is a pullback groupoid $\mathcal{G}[Z]$.
- $\mathcal{G}[Z]^{(0)} = Z$, $\mathcal{G}[Z]^{(1)} = \{z \xrightarrow{g} z' \mid f(z) \xrightarrow{g} f(z')\}$.
- Groupoids \mathcal{G} and \mathcal{H} are **Morita equivalent** if there is a locally compact space Z and continuous open surjections $p: Z \rightarrow \mathcal{G}^{(0)}$ and $q: Z \rightarrow \mathcal{H}^{(0)}$ such that $\mathcal{G}[Z] \cong \mathcal{H}[Z]$.

Theorem (L. O. Clark, A. Sims)

If \mathcal{G} and \mathcal{H} are Morita equivalent ample groupoids, then $\mathbb{k}\mathcal{G}$ is Morita equivalent to $\mathbb{k}\mathcal{H}$.

Morita equivalence

- If Z is a locally compact space and $f: Z \rightarrow \mathcal{G}^{(0)}$ is continuous, there is a pullback groupoid $\mathcal{G}[Z]$.
- $\mathcal{G}[Z]^{(0)} = Z$, $\mathcal{G}[Z]^{(1)} = \{z \xrightarrow{g} z' \mid f(z) \xrightarrow{g} f(z')\}$.
- Groupoids \mathcal{G} and \mathcal{H} are **Morita equivalent** if there is a locally compact space Z and continuous open surjections $p: Z \rightarrow \mathcal{G}^{(0)}$ and $q: Z \rightarrow \mathcal{H}^{(0)}$ such that $\mathcal{G}[Z] \cong \mathcal{H}[Z]$.

Theorem (L. O. Clark, A. Sims)

If \mathcal{G} and \mathcal{H} are Morita equivalent ample groupoids, then $\mathbb{k}\mathcal{G}$ is Morita equivalent to $\mathbb{k}\mathcal{H}$.

- Gives a uniform explanation for Morita equivalences of graph C^* -algebras and Leavitt path algebras.

Morita equivalence

- If Z is a locally compact space and $f: Z \rightarrow \mathcal{G}^{(0)}$ is continuous, there is a pullback groupoid $\mathcal{G}[Z]$.
- $\mathcal{G}[Z]^{(0)} = Z$, $\mathcal{G}[Z]^{(1)} = \{z \xrightarrow{g} z' \mid f(z) \xrightarrow{g} f(z')\}$.
- Groupoids \mathcal{G} and \mathcal{H} are **Morita equivalent** if there is a locally compact space Z and continuous open surjections $p: Z \rightarrow \mathcal{G}^{(0)}$ and $q: Z \rightarrow \mathcal{H}^{(0)}$ such that $\mathcal{G}[Z] \cong \mathcal{H}[Z]$.

Theorem (L. O. Clark, A. Sims)

If \mathcal{G} and \mathcal{H} are Morita equivalent ample groupoids, then $\mathbb{k}\mathcal{G}$ is Morita equivalent to $\mathbb{k}\mathcal{H}$.

- Gives a uniform explanation for Morita equivalences of graph C^* -algebras and Leavitt path algebras.
- The original proof constructed a Morita context.

Morita equivalence

- If Z is a locally compact space and $f: Z \rightarrow \mathcal{G}^{(0)}$ is continuous, there is a pullback groupoid $\mathcal{G}[Z]$.
- $\mathcal{G}[Z]^{(0)} = Z$, $\mathcal{G}[Z]^{(1)} = \{z \xrightarrow{g} z' \mid f(z) \xrightarrow{g} f(z')\}$.
- Groupoids \mathcal{G} and \mathcal{H} are **Morita equivalent** if there is a locally compact space Z and continuous open surjections $p: Z \rightarrow \mathcal{G}^{(0)}$ and $q: Z \rightarrow \mathcal{H}^{(0)}$ such that $\mathcal{G}[Z] \cong \mathcal{H}[Z]$.

Theorem (L. O. Clark, A. Sims)

If \mathcal{G} and \mathcal{H} are Morita equivalent ample groupoids, then $\mathbb{k}\mathcal{G}$ is Morita equivalent to $\mathbb{k}\mathcal{H}$.

- Gives a uniform explanation for Morita equivalences of graph C^* -algebras and Leavitt path algebras.
- The original proof constructed a Morita context.
- A result of Moerdijk implies if \mathcal{G}, \mathcal{H} are Morita equivalent, then so are their categories of sheaves.

Schützenberger representations


- Fix $x \in \mathcal{G}^{(0)}$.

Schützenberger representations

- Fix $x \in \mathcal{G}^{(0)}$.
- Let $L_x = d^{-1}(x)$ be the arrows emanating from x .


Schützenberger representations

- Fix $x \in \mathcal{G}^{(0)}$.
- Let $L_x = d^{-1}(x)$ be the arrows emanating from x .
- The isotropy group G_x acts freely on the right of L_x .



Schützenberger representations


- Fix $x \in \mathcal{G}^{(0)}$.
- Let $L_x = d^{-1}(x)$ be the arrows emanating from x .
- The isotropy group G_x acts freely on the right of L_x .



- So $\mathbb{k}L_x$ is a free $\mathbb{k}G_x$ -module.

Schützenberger representations


- Fix $x \in \mathcal{G}^{(0)}$.
- Let $L_x = d^{-1}(x)$ be the arrows emanating from x .
- The isotropy group G_x acts freely on the right of L_x .



- So $\mathbb{k}L_x$ is a free $\mathbb{k}G_x$ -module.
- It is in fact a $\mathbb{k}\mathcal{G}$ - $\mathbb{k}G_x$ -bimodule.

Schützenberger representations

- Fix $x \in \mathcal{G}^{(0)}$.
- Let $L_x = d^{-1}(x)$ be the arrows emanating from x .
- The isotropy group G_x acts freely on the right of L_x .



- So $\mathbb{k}L_x$ is a free $\mathbb{k}G_x$ -module.
- It is in fact a $\mathbb{k}\mathcal{G}$ - $\mathbb{k}G_x$ -bimodule.
- If $f \in \mathbb{k}\mathcal{G}$ and $t \in L_x$, then

$$f \cdot t = \sum_{d(s)=r(t)} f(s)st.$$

Induction and restriction functors

- There is an exact functor $\mathrm{Ind}_x: \mathbb{k}G_x\text{-mod} \rightarrow \mathbb{k}\mathcal{G}\text{-mod}$ given by

$$M \longmapsto \mathbb{k}L_x \otimes_{\mathbb{k}G_x} M.$$

Induction and restriction functors

- There is an exact functor $\mathrm{Ind}_x: \mathbb{k}G_x\text{-mod} \rightarrow \mathbb{k}\mathcal{G}\text{-mod}$ given by

$$M \longmapsto \mathbb{k}L_x \otimes_{\mathbb{k}G_x} M.$$

- It has a right adjoint $\mathrm{Res}_x: \mathbb{k}\mathcal{G}\text{-mod} \rightarrow \mathbb{k}G_x\text{-mod}$.

Induction and restriction functors

- There is an exact functor $\mathrm{Ind}_x: \mathbb{k}G_x\text{-mod} \rightarrow \mathbb{k}\mathcal{G}\text{-mod}$ given by

$$M \longmapsto \mathbb{k}L_x \otimes_{\mathbb{k}G_x} M.$$

- It has a right adjoint $\mathrm{Res}_x: \mathbb{k}\mathcal{G}\text{-mod} \rightarrow \mathbb{k}G_x\text{-mod}$.
- Ind_x preserves simple modules.

Induction and restriction functors

- There is an exact functor $\mathrm{Ind}_x: \mathbb{k}G_x\text{-mod} \rightarrow \mathbb{k}\mathcal{G}\text{-mod}$ given by

$$M \longmapsto \mathbb{k}L_x \otimes_{\mathbb{k}G_x} M.$$

- It has a right adjoint $\mathrm{Res}_x: \mathbb{k}\mathcal{G}\text{-mod} \rightarrow \mathbb{k}G_x\text{-mod}$.
- Ind_x preserves simple modules.
- Res_x sends simple modules to simple modules or 0.

Induction and restriction functors

- There is an exact functor $\mathrm{Ind}_x: \mathbb{k}G_x\text{-mod} \rightarrow \mathbb{k}\mathcal{G}\text{-mod}$ given by

$$M \longmapsto \mathbb{k}L_x \otimes_{\mathbb{k}G_x} M.$$

- It has a right adjoint $\mathrm{Res}_x: \mathbb{k}\mathcal{G}\text{-mod} \rightarrow \mathbb{k}G_x\text{-mod}$.
- Ind_x preserves simple modules.
- Res_x sends simple modules to simple modules or 0.

Theorem (BS)

Let \mathbb{k} be a field and \mathcal{G} an ample groupoid. Then the finite dimensional simple $\mathbb{k}\mathcal{G}$ -modules are the $\mathrm{Ind}_x(M)$ with $|\mathcal{O}_x| < \infty$ and M a finite dimensional simple $\mathbb{k}G_x$ -module.

A question of Exel

Question (Exel)

Let \mathbb{k} be a field and \mathcal{G} a Hausdorff ample groupoid. Is every primitive ideal of $\mathbb{k}\mathcal{G}$ a kernel of an induced module $\mathrm{Ind}_x(M)$ with M a simple $\mathbb{k}G_x$ -module?

A question of Exel

Question (Exel)

Let \mathbb{k} be a field and \mathcal{G} a Hausdorff ample groupoid. Is every primitive ideal of $\mathbb{k}\mathcal{G}$ a kernel of an induced module $\text{Ind}_x(M)$ with M a simple $\mathbb{k}G_x$ -module?

- A C^* -algebraic analogue is true if \mathcal{G} is amenable.

A question of Exel

Question (Exel)

Let \mathbb{k} be a field and \mathcal{G} a Hausdorff ample groupoid. Is every primitive ideal of $\mathbb{k}\mathcal{G}$ a kernel of an induced module $\text{Ind}_x(M)$ with M a simple $\mathbb{k}G_x$ -module?

- A C^* -algebraic analogue is true if \mathcal{G} is amenable.
- It is true for Leavitt path algebras.

Further work

- There are still many ample groupoid C^* -algebras whose algebraic analogues have not been studied.

Further work

- There are still many ample groupoid C^* -algebras whose algebraic analogues have not been studied.
- Algebras associated to tilings.

Further work

- There are still many ample groupoid C^* -algebras whose algebraic analogues have not been studied.
- Algebras associated to tilings.
- Algebras associated to self-similar/automaton groups.

Further work

- There are still many ample groupoid C^* -algebras whose algebraic analogues have not been studied.
- Algebras associated to tilings.
- Algebras associated to self-similar/automaton groups.
- Algebras associated to Cantor dynamical systems.

Further work

- There are still many ample groupoid C^* -algebras whose algebraic analogues have not been studied.
- Algebras associated to tilings.
- Algebras associated to self-similar/automaton groups.
- Algebras associated to Cantor dynamical systems.
- Algebras associated to higher rank graphs.

Further work

- There are still many ample groupoid C^* -algebras whose algebraic analogues have not been studied.
- Algebras associated to tilings.
- Algebras associated to self-similar/automaton groups.
- Algebras associated to Cantor dynamical systems.
- Algebras associated to higher rank graphs.
- Algebras associated to partial group actions.

The end

Thank you for your attention!