

# Symbolic Dynamics, Profinite Groups and Profinite Monoids

Benjamin Steinberg

Carleton University

University of Leipzig

May, 2009

# An ultrametric completion of the free monoid

- $A^*$  denotes the free monoid on  $A$ .
- We always assume  $2 \leq |A| < \infty$ .
- An **ultrametric** can be defined on  $A^*$  by putting

$$d(u, v) = |A|^{-|u \wedge v|}$$

where  $u \wedge v$  is the longest common prefix of  $u$  and  $v$ .

- The completion is  $A^* \cup A^\omega$ , which can be viewed as a regular rooted tree together with its boundary.
- The boundary  $A^\omega$  is the realm of symbolic dynamics.

# An ultrametric completion of the free monoid

- $A^*$  denotes the free monoid on  $A$ .
- We always assume  $2 \leq |A| < \infty$ .
- An **ultrametric** can be defined on  $A^*$  by putting

$$d(u, v) = |A|^{-|u \wedge v|}$$

where  $u \wedge v$  is the longest common prefix of  $u$  and  $v$ .

- The completion is  $A^* \cup A^\omega$ , which can be viewed as a regular rooted tree together with its boundary.
- The boundary  $A^\omega$  is the realm of **symbolic dynamics**.

# An ultrametric completion of the free monoid

- $A^*$  denotes the free monoid on  $A$ .
- We always assume  $2 \leq |A| < \infty$ .
- An **ultrametric** can be defined on  $A^*$  by putting

$$d(u, v) = |A|^{-|u \wedge v|}$$

where  $u \wedge v$  is the longest common prefix of  $u$  and  $v$ .

- The completion is  $A^* \cup A^\omega$ , which can be viewed as a regular rooted tree together with its boundary.
- The boundary  $A^\omega$  is the realm of **symbolic dynamics**.

# An ultrametric completion of the free monoid

- $A^*$  denotes the free monoid on  $A$ .
- We always assume  $2 \leq |A| < \infty$ .
- An **ultrametric** can be defined on  $A^*$  by putting

$$d(u, v) = |A|^{-|u \wedge v|}$$

where  $u \wedge v$  is the longest common prefix of  $u$  and  $v$ .

- The completion is  $A^* \cup A^\omega$ , which can be viewed as a regular rooted tree together with its boundary.
- The boundary  $A^\omega$  is the realm of **symbolic dynamics**.

# An ultrametric completion of the free monoid

- $A^*$  denotes the free monoid on  $A$ .
- We always assume  $2 \leq |A| < \infty$ .
- An **ultrametric** can be defined on  $A^*$  by putting

$$d(u, v) = |A|^{-|u \wedge v|}$$

where  $u \wedge v$  is the longest common prefix of  $u$  and  $v$ .

- The completion is  $A^* \cup A^\omega$ , which can be viewed as a regular rooted tree together with its boundary.
- The boundary  $A^\omega$  is the realm of **symbolic dynamics**.

# Symbolic dynamics

- The **shift** map  $\sigma: A^\omega \rightarrow A^\omega$  is given by

$$\sigma(a_0a_1\cdots) = a_1a_2\cdots.$$

- A **subshift** is a closed, non-empty, shift-invariant subspace of  $A^\omega$ .
- Subshifts  $\mathcal{X} \subseteq A^\omega$  and  $\mathcal{Y} \subseteq B^\omega$  are **conjugate** if there is a homeomorphism  $\psi: \mathcal{X} \rightarrow \mathcal{Y}$  so that

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\sigma} & \mathcal{X} \\ \psi \downarrow & & \downarrow \psi \\ \mathcal{Y} & \xrightarrow{\sigma} & \mathcal{Y} \end{array}$$

commutes.

# Symbolic dynamics

- The **shift** map  $\sigma: A^\omega \rightarrow A^\omega$  is given by

$$\sigma(a_0a_1\cdots) = a_1a_2\cdots.$$

- A **subshift** is a closed, non-empty, shift-invariant subspace of  $A^\omega$ .
- Subshifts  $\mathcal{X} \subseteq A^\omega$  and  $\mathcal{Y} \subseteq B^\omega$  are **conjugate** if there is a homeomorphism  $\psi: \mathcal{X} \rightarrow \mathcal{Y}$  so that

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\sigma} & \mathcal{X} \\ \psi \downarrow & & \downarrow \psi \\ \mathcal{Y} & \xrightarrow{\sigma} & \mathcal{Y} \end{array}$$

commutes.



# Symbolic dynamics

- The **shift** map  $\sigma: A^\omega \rightarrow A^\omega$  is given by

$$\sigma(a_0a_1\cdots) = a_1a_2\cdots.$$

- A **subshift** is a closed, non-empty, shift-invariant subspace of  $A^\omega$ .
- Subshifts  $\mathcal{X} \subseteq A^\omega$  and  $\mathcal{Y} \subseteq B^\omega$  are **conjugate** if there is a homeomorphism  $\psi: \mathcal{X} \rightarrow \mathcal{Y}$  so that

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\sigma} & \mathcal{X} \\ \psi \downarrow & & \downarrow \psi \\ \mathcal{Y} & \xrightarrow{\sigma} & \mathcal{Y} \end{array}$$

commutes.

# Irreducible subshifts

- Let  $\mathcal{X} \subseteq A^\omega$  be a subshift.
- $\mathcal{X}$  is said to be **irreducible** if it has a dense orbit.
- Let  $L(\mathcal{X}) \subseteq A^*$  denote the language of all finite factors of elements of  $\mathcal{X}$ .
- It turns out  $\mathcal{X}$  is irreducible if and only if, for all  $u, v \in L(\mathcal{X})$ , there exists  $w \in A^*$  so that  $uwv \in L(\mathcal{X})$ .
- In this talk we consider only irreducible subshifts.
- The map  $\mathcal{X} \mapsto L(\mathcal{X})$  is injective.
- Indeed,  $\mathcal{X} = \overline{\partial L(\mathcal{X})}$  (it is the boundary of the subtree spanned by  $L(\mathcal{X})$ ).

# Irreducible subshifts

- Let  $\mathcal{X} \subseteq A^\omega$  be a subshift.
- $\mathcal{X}$  is said to be **irreducible** if it has a dense orbit.
- Let  $L(\mathcal{X}) \subseteq A^*$  denote the language of all finite factors of elements of  $\mathcal{X}$ .
- It turns out  $\mathcal{X}$  is irreducible if and only if, for all  $u, v \in L(\mathcal{X})$ , there exists  $w \in A^*$  so that  $uwv \in L(\mathcal{X})$ .
- In this talk we consider only irreducible subshifts.
- The map  $\mathcal{X} \mapsto L(\mathcal{X})$  is injective.
- Indeed,  $\mathcal{X} = \overline{\partial L(\mathcal{X})}$  (it is the boundary of the subtree spanned by  $L(\mathcal{X})$ ).

# Irreducible subshifts

- Let  $\mathcal{X} \subseteq A^\omega$  be a subshift.
- $\mathcal{X}$  is said to be **irreducible** if it has a dense orbit.
- Let  $L(\mathcal{X}) \subseteq A^*$  denote the language of all finite factors of elements of  $\mathcal{X}$ .
- It turns out  $\mathcal{X}$  is irreducible if and only if, for all  $u, v \in L(\mathcal{X})$ , there exists  $w \in A^*$  so that  $uwv \in L(\mathcal{X})$ .
- In this talk we consider only irreducible subshifts.
- The map  $\mathcal{X} \mapsto L(\mathcal{X})$  is injective.
- Indeed,  $\mathcal{X} = \overline{\partial L(\mathcal{X})}$  (it is the boundary of the subtree spanned by  $L(\mathcal{X})$ ).

# Irreducible subshifts

- Let  $\mathcal{X} \subseteq A^\omega$  be a subshift.
- $\mathcal{X}$  is said to be **irreducible** if it has a dense orbit.
- Let  $L(\mathcal{X}) \subseteq A^*$  denote the language of all finite factors of elements of  $\mathcal{X}$ .
- It turns out  $\mathcal{X}$  is irreducible if and only if, for all  $u, v \in L(\mathcal{X})$ , there exists  $w \in A^*$  so that  $uwv \in L(\mathcal{X})$ .
- In this talk we consider only irreducible subshifts.
- The map  $\mathcal{X} \mapsto L(\mathcal{X})$  is injective.
- Indeed,  $\mathcal{X} = \overline{\partial L(\mathcal{X})}$  (it is the boundary of the subtree spanned by  $L(\mathcal{X})$ ).

# Irreducible subshifts

- Let  $\mathcal{X} \subseteq A^\omega$  be a subshift.
- $\mathcal{X}$  is said to be **irreducible** if it has a dense orbit.
- Let  $L(\mathcal{X}) \subseteq A^*$  denote the language of all finite factors of elements of  $\mathcal{X}$ .
- It turns out  $\mathcal{X}$  is irreducible if and only if, for all  $u, v \in L(\mathcal{X})$ , there exists  $w \in A^*$  so that  $uwv \in L(\mathcal{X})$ .
- In this talk we consider only irreducible subshifts.
- The map  $\mathcal{X} \mapsto L(\mathcal{X})$  is injective.
- Indeed,  $\mathcal{X} = \overline{\partial L(\mathcal{X})}$  (it is the boundary of the subtree spanned by  $L(\mathcal{X})$ ).

# Irreducible subshifts

- Let  $\mathcal{X} \subseteq A^\omega$  be a subshift.
- $\mathcal{X}$  is said to be **irreducible** if it has a dense orbit.
- Let  $L(\mathcal{X}) \subseteq A^*$  denote the language of all finite factors of elements of  $\mathcal{X}$ .
- It turns out  $\mathcal{X}$  is irreducible if and only if, for all  $u, v \in L(\mathcal{X})$ , there exists  $w \in A^*$  so that  $uwv \in L(\mathcal{X})$ .
- In this talk we consider only irreducible subshifts.
- The map  $\mathcal{X} \mapsto L(\mathcal{X})$  is injective.
- Indeed,  $\mathcal{X} = \overline{\partial L(\mathcal{X})}$  (it is the boundary of the subtree spanned by  $L(\mathcal{X})$ ).

# Irreducible subshifts

- Let  $\mathcal{X} \subseteq A^\omega$  be a subshift.
- $\mathcal{X}$  is said to be **irreducible** if it has a dense orbit.
- Let  $L(\mathcal{X}) \subseteq A^*$  denote the language of all finite factors of elements of  $\mathcal{X}$ .
- It turns out  $\mathcal{X}$  is irreducible if and only if, for all  $u, v \in L(\mathcal{X})$ , there exists  $w \in A^*$  so that  $uwv \in L(\mathcal{X})$ .
- In this talk we consider only irreducible subshifts.
- The map  $\mathcal{X} \mapsto L(\mathcal{X})$  is injective.
- Indeed,  $\mathcal{X} = \overline{\partial L(\mathcal{X})}$  (it is the boundary of the subtree spanned by  $L(\mathcal{X})$ ).



# Minimal subshifts

- A **minimal subshift** must be the closure of the orbit of any of its elements under the shift.
- It follows immediately that minimal subshifts are irreducible.
- A word  $w \in A^\omega$  generates a minimal subshift if and only if  $w$  is **uniformly recurrent**.
- This means that if  $v$  is a finite factor of  $w$ , then there exists  $N > 0$  so that each factor of  $w$  of length  $N$  contains  $v$  as a factor: the “bounded gaps property.”
- If  $u \in A^*$ , then the subshift generated by

$$u^\omega = uuu \cdots$$

is finite and minimal. Such a minimal subshift is called **periodic**.

# Minimal subshifts

- A **minimal subshift** must be the closure of the orbit of any of its elements under the shift.
- It follows immediately that minimal subshifts are irreducible.
- A word  $w \in A^\omega$  generates a minimal subshift if and only if  $w$  is **uniformly recurrent**.
- This means that if  $v$  is a finite factor of  $w$ , then there exists  $N > 0$  so that each factor of  $w$  of length  $N$  contains  $v$  as a factor: the “**bounded gaps property**.”
- If  $u \in A^*$ , then the subshift generated by

$$u^\omega = uuu \cdots$$

is finite and minimal. Such a minimal subshift is called **periodic**.

# Minimal subshifts

- A **minimal subshift** must be the closure of the orbit of any of its elements under the shift.
- It follows immediately that minimal subshifts are irreducible.
- A word  $w \in A^\omega$  generates a minimal subshift if and only if  $w$  is **uniformly recurrent**.
- This means that if  $v$  is a finite factor of  $w$ , then there exists  $N > 0$  so that each factor of  $w$  of length  $N$  contains  $v$  as a factor: the “**bounded gaps property**.”
- If  $u \in A^*$ , then the subshift generated by

$$u^\omega = uuu \cdots$$

is finite and minimal. Such a minimal subshift is called **periodic**.

# Minimal subshifts

- A **minimal subshift** must be the closure of the orbit of any of its elements under the shift.
- It follows immediately that minimal subshifts are irreducible.
- A word  $w \in A^\omega$  generates a minimal subshift if and only if  $w$  is **uniformly recurrent**.
- This means that if  $v$  is a finite factor of  $w$ , then there exists  $N > 0$  so that each factor of  $w$  of length  $N$  contains  $v$  as a factor: the “**bounded gaps property**.”
- If  $u \in A^*$ , then the subshift generated by

$$u^\omega = uuu \cdots$$

is finite and minimal. Such a minimal subshift is called **periodic**.

# Minimal subshifts

- A **minimal subshift** must be the closure of the orbit of any of its elements under the shift.
- It follows immediately that minimal subshifts are irreducible.
- A word  $w \in A^\omega$  generates a minimal subshift if and only if  $w$  is **uniformly recurrent**.
- This means that if  $v$  is a finite factor of  $w$ , then there exists  $N > 0$  so that each factor of  $w$  of length  $N$  contains  $v$  as a factor: the “**bounded gaps property**.”
- If  $u \in A^*$ , then the subshift generated by

$$u^\omega = uuu \cdots$$

is finite and minimal. Such a minimal subshift is called **periodic**.

# Uniform recurrence and primitive substitutions

- The famous **Morse-Thue** cube-free word is uniformly recurrent. It is the fixed point obtained by iterating the substitution

$$a \mapsto ab, b \mapsto ba$$

starting from  $a$ .

- $a$
- $ab$
- $abba$
- $abbabaab$
- $abbabaabbaababba \dots$
- A substitution  $f: A^* \rightarrow A^*$  is primitive if there exists  $N > 0$  so that each letter of  $A$  appears in  $f^N(a)$ , all  $a \in A$ .
- If  $f$  is a primitive substitution with  $a$  the first letter of  $f(a)$ , then  $\lim f^n(a)$  is a uniformly recurrent word.

# Uniform recurrence and primitive substitutions

- The famous **Morse-Thue** cube-free word is uniformly recurrent. It is the fixed point obtained by iterating the substitution

$$a \mapsto ab, b \mapsto ba$$

starting from  $a$ .

- $a$
- $ab$
- $abba$
- $abbabaab$
- $abbabaabbaababba \dots$
- A substitution  $f: A^* \rightarrow A^*$  is primitive if there exists  $N > 0$  so that each letter of  $A$  appears in  $f^N(a)$ , all  $a \in A$ .
- If  $f$  is a primitive substitution with  $a$  the first letter of  $f(a)$ , then  $\lim f^n(a)$  is a uniformly recurrent word.

# Uniform recurrence and primitive substitutions

- The famous **Morse-Thue** cube-free word is uniformly recurrent. It is the fixed point obtained by iterating the substitution

$$a \mapsto ab, b \mapsto ba$$

starting from  $a$ .

- $a$
- $ab$
- $abba$
- $abbabaab$
- $abbabaabbaababba \dots$
- A substitution  $f: A^* \rightarrow A^*$  is primitive if there exists  $N > 0$  so that each letter of  $A$  appears in  $f^N(a)$ , all  $a \in A$ .
- If  $f$  is a primitive substitution with  $a$  the first letter of  $f(a)$ , then  $\lim f^n(a)$  is a uniformly recurrent word.



# Uniform recurrence and primitive substitutions

- The famous **Morse-Thue** cube-free word is uniformly recurrent. It is the fixed point obtained by iterating the substitution

$$a \mapsto ab, b \mapsto ba$$

starting from  $a$ .

- $a$
- $ab$
- $abba$
- $abbabaab$
- $abbabaabbaababba \dots$
- A substitution  $f: A^* \rightarrow A^*$  is **primitive** if there exists  $N > 0$  so that each letter of  $A$  appears in  $f^N(a)$ , all  $a \in A$ .
- If  $f$  is a primitive substitution with  $a$  the first letter of  $f(a)$ , then  $\lim f^n(a)$  is a uniformly recurrent word.

# Uniform recurrence and primitive substitutions

- The famous **Morse-Thue** cube-free word is uniformly recurrent. It is the fixed point obtained by iterating the substitution

$$a \mapsto ab, b \mapsto ba$$

starting from  $a$ .

- $a$
- $ab$
- $abba$
- $abbabaab$
- $abbabaabbaababba \dots$
- A substitution  $f: A^* \rightarrow A^*$  is **primitive** if there exists  $N > 0$  so that each letter of  $A$  appears in  $f^N(a)$ , all  $a \in A$ .
- If  $f$  is a primitive substitution with  $a$  the first letter of  $f(a)$ , then  $\lim f^n(a)$  is a uniformly recurrent word.

# Uniform recurrence and primitive substitutions

- The famous **Morse-Thue** cube-free word is uniformly recurrent. It is the fixed point obtained by iterating the substitution

$$a \mapsto ab, b \mapsto ba$$

starting from  $a$ .

- $a$
- $ab$
- $abba$
- $abbabaab$
- $abbabaabbaababba \dots$
- A substitution  $f: A^* \rightarrow A^*$  is **primitive** if there exists  $N > 0$  so that each letter of  $A$  appears in  $f^N(a)$ , all  $a \in A$ .
- If  $f$  is a primitive substitution with  $a$  the first letter of  $f(a)$ , then  $\lim f^n(a)$  is a uniformly recurrent word.

# Uniform recurrence and primitive substitutions

- The famous **Morse-Thue** cube-free word is uniformly recurrent. It is the fixed point obtained by iterating the substitution

$$a \mapsto ab, b \mapsto ba$$

starting from  $a$ .

- $a$
- $ab$
- $abba$
- $abbabaab$
- $abbabaabbaababba \dots$
- A substitution  $f: A^* \rightarrow A^*$  is **primitive** if there exists  $N > 0$  so that each letter of  $A$  appears in  $f^N(a)$ , all  $a \in A$ .
- If  $f$  is a primitive substitution with  $a$  the first letter of  $f(a)$ , then  $\lim f^n(a)$  is a uniformly recurrent word.

# Uniform recurrence and primitive substitutions

- The famous **Morse-Thue** cube-free word is uniformly recurrent. It is the fixed point obtained by iterating the substitution

$$a \mapsto ab, b \mapsto ba$$

starting from  $a$ .

- $a$
- $ab$
- $abba$
- $abbabaab$
- $abbabaabbaababba \dots$
- A substitution  $f: A^* \rightarrow A^*$  is **primitive** if there exists  $N > 0$  so that each letter of  $A$  appears in  $f^N(a)$ , all  $a \in A$ .
- If  $f$  is a primitive substitution with  $a$  the first letter of  $f(a)$ , then  $\lim f^n(a)$  is a uniformly recurrent word.

# Sofic shifts

- A subshift  $\mathcal{X} \subseteq A^\omega$  is said to be of **finite type** if there is a finite set  $F$  of **forbidden factors** defining  $\mathcal{X}$ . That is,  $L(\mathcal{X}) = A^* \setminus A^*FA^*$ .
- In this case,  $L(\mathcal{X})$  is a **regular language**, i.e., recognized by a finite automaton.
- Weiss defined  $\mathcal{X}$  to be a **sofic shift** if  $L(\mathcal{X})$  is regular.
- Sofic shifts are precisely the quotients (in the appropriate category) of subshifts of finite type.
- Irreducible sofic shifts can always be recognized by a strongly connected automaton all of whose states are initial and final.
- A minimal sofic shift must be periodic (follow a cycle in the automaton).

# Sofic shifts

- A subshift  $\mathcal{X} \subseteq A^\omega$  is said to be of **finite type** if there is a finite set  $F$  of **forbidden factors** defining  $\mathcal{X}$ . That is,  $L(\mathcal{X}) = A^* \setminus A^*FA^*$ .
- In this case,  $L(\mathcal{X})$  is a **regular language**, i.e., recognized by a finite automaton.
- Weiss defined  $\mathcal{X}$  to be a **sofic shift** if  $L(\mathcal{X})$  is regular.
- Sofic shifts are precisely the quotients (in the appropriate category) of subshifts of finite type.
- Irreducible sofic shifts can always be recognized by a strongly connected automaton all of whose states are initial and final.
- A minimal sofic shift must be periodic (follow a cycle in the automaton).

# Sofic shifts

- A subshift  $\mathcal{X} \subseteq A^\omega$  is said to be of **finite type** if there is a finite set  $F$  of **forbidden factors** defining  $\mathcal{X}$ . That is,  $L(\mathcal{X}) = A^* \setminus A^*FA^*$ .
- In this case,  $L(\mathcal{X})$  is a **regular language**, i.e., recognized by a finite automaton.
- Weiss defined  $\mathcal{X}$  to be a **sofic shift** if  $L(\mathcal{X})$  is regular.
- Sofic shifts are precisely the quotients (in the appropriate category) of subshifts of finite type.
- Irreducible sofic shifts can always be recognized by a strongly connected automaton all of whose states are initial and final.
- A minimal sofic shift must be periodic (follow a cycle in the automaton).



# Sofic shifts

- A subshift  $\mathcal{X} \subseteq A^\omega$  is said to be of **finite type** if there is a finite set  $F$  of **forbidden factors** defining  $\mathcal{X}$ . That is,  $L(\mathcal{X}) = A^* \setminus A^*FA^*$ .
- In this case,  $L(\mathcal{X})$  is a **regular language**, i.e., recognized by a finite automaton.
- Weiss defined  $\mathcal{X}$  to be a **sofic shift** if  $L(\mathcal{X})$  is regular.
- Sofic shifts are precisely the quotients (in the appropriate category) of subshifts of finite type.
- Irreducible sofic shifts can always be recognized by a strongly connected automaton all of whose states are initial and final.
- A minimal sofic shift must be periodic (follow a cycle in the automaton).

# Sofic shifts


- A subshift  $\mathcal{X} \subseteq A^\omega$  is said to be of **finite type** if there is a finite set  $F$  of **forbidden factors** defining  $\mathcal{X}$ . That is,  $L(\mathcal{X}) = A^* \setminus A^*FA^*$ .
- In this case,  $L(\mathcal{X})$  is a **regular language**, i.e., recognized by a finite automaton.
- Weiss defined  $\mathcal{X}$  to be a **sofic shift** if  $L(\mathcal{X})$  is regular.
- Sofic shifts are precisely the quotients (in the appropriate category) of subshifts of finite type.
- Irreducible sofic shifts can always be recognized by a strongly connected automaton all of whose states are initial and final.
- A minimal sofic shift must be periodic (follow a cycle in the automaton).

# Sofic shifts

- A subshift  $\mathcal{X} \subseteq A^\omega$  is said to be of **finite type** if there is a finite set  $F$  of **forbidden factors** defining  $\mathcal{X}$ . That is,  $L(\mathcal{X}) = A^* \setminus A^*FA^*$ .
- In this case,  $L(\mathcal{X})$  is a **regular language**, i.e., recognized by a finite automaton.
- Weiss defined  $\mathcal{X}$  to be a **sofic shift** if  $L(\mathcal{X})$  is regular.
- Sofic shifts are precisely the quotients (in the appropriate category) of subshifts of finite type.
- Irreducible sofic shifts can always be recognized by a strongly connected automaton all of whose states are initial and final.
- A minimal sofic shift must be periodic (follow a cycle in the automaton).

# The even shift

- An automaton recognizing the **even shift**:



- The even shift consists of all infinite words with an even number of occurrences of **b** between consecutive occurrences of **a**.

# A hidden profinite monoid

- Multiplication on  $A^*$  is uniformly continuous in the ultrametric defining the completion  $A^* \cup A^\omega$ .
- Thus  $A^* \cup A^\omega$  has the structure of an  $A$ -generated profinite monoid.
- The product is given by

$$u \cdot v = \begin{cases} uv & u \in A^* \\ u & u \in A^\omega. \end{cases}$$

- It follows that  $A^* \cup A^\omega$  is a continuous homomorphic image of the profinite completion  $\widehat{A^*}$  of  $A^*$ , i.e., of the free profinite monoid on  $A$ .
- Almeida used this to “lift” symbolic dynamics to the free profinite monoid.

# A hidden profinite monoid

- Multiplication on  $A^*$  is uniformly continuous in the ultrametric defining the completion  $A^* \cup A^\omega$ .
- Thus  $A^* \cup A^\omega$  has the structure of an  $A$ -generated profinite monoid.
- The product is given by

$$u \cdot v = \begin{cases} uv & u \in A^* \\ u & u \in A^\omega. \end{cases}$$

- It follows that  $A^* \cup A^\omega$  is a continuous homomorphic image of the profinite completion  $\widehat{A^*}$  of  $A^*$ , i.e., of the free profinite monoid on  $A$ .
- Almeida used this to “lift” symbolic dynamics to the free profinite monoid.

# A hidden profinite monoid

- Multiplication on  $A^*$  is uniformly continuous in the ultrametric defining the completion  $A^* \cup A^\omega$ .
- Thus  $A^* \cup A^\omega$  has the structure of an  $A$ -generated profinite monoid.
- The product is given by

$$u \cdot v = \begin{cases} uv & u \in A^* \\ u & u \in A^\omega. \end{cases}$$

- It follows that  $A^* \cup A^\omega$  is a continuous homomorphic image of the profinite completion  $\widehat{A^*}$  of  $A^*$ , i.e., of the free profinite monoid on  $A$ .
- Almeida used this to “lift” symbolic dynamics to the free profinite monoid.

# A hidden profinite monoid

- Multiplication on  $A^*$  is uniformly continuous in the ultrametric defining the completion  $A^* \cup A^\omega$ .
- Thus  $A^* \cup A^\omega$  has the structure of an  $A$ -generated profinite monoid.
- The product is given by

$$u \cdot v = \begin{cases} uv & u \in A^* \\ u & u \in A^\omega. \end{cases}$$

- It follows that  $A^* \cup A^\omega$  is a continuous homomorphic image of the profinite completion  $\widehat{A^*}$  of  $A^*$ , i.e., of the free profinite monoid on  $A$ .
- Almeida used this to “lift” symbolic dynamics to the free profinite monoid.



# A hidden profinite monoid

- Multiplication on  $A^*$  is uniformly continuous in the ultrametric defining the completion  $A^* \cup A^\omega$ .
- Thus  $A^* \cup A^\omega$  has the structure of an  $A$ -generated profinite monoid.
- The product is given by

$$u \cdot v = \begin{cases} uv & u \in A^* \\ u & u \in A^\omega. \end{cases}$$

- It follows that  $A^* \cup A^\omega$  is a continuous homomorphic image of the profinite completion  $\widehat{A^*}$  of  $A^*$ , i.e., of the free profinite monoid on  $A$ .
- Almeida used this to “lift” symbolic dynamics to the free profinite monoid.

# The profinite completion of the free monoid

- For  $u \neq v \in A^*$ , define  $\sigma(u, v)$  to be the minimum size of a finite monoid separating  $u$  from  $v$ .
- $A^*$  is residually finite, so  $\sigma(u, v)$  is well defined.
- The profinite ultrametric on  $A^*$  is defined by

$$d(u, v) = |A|^{-\sigma(u, v)}.$$

- The completion is the free profinite monoid  $\widehat{A^*}$ .
- Set  $\partial\widehat{A^*} = \widehat{A^*} \setminus A^*$ ; it is an ideal of  $\widehat{A^*}$ .
- $\widehat{A^*}$  plays a crucial role in automata theory because it is the Zariski spectrum of the Boolean ring of regular languages.

# The profinite completion of the free monoid

- For  $u \neq v \in A^*$ , define  $\sigma(u, v)$  to be the minimum size of a finite monoid separating  $u$  from  $v$ .
- $A^*$  is residually finite, so  $\sigma(u, v)$  is well defined.
- The **profinite ultrametric** on  $A^*$  is defined by

$$d(u, v) = |A|^{-\sigma(u, v)}.$$

- The completion is the **free profinite monoid**  $\widehat{A^*}$ .
- Set  $\partial\widehat{A^*} = \widehat{A^*} \setminus A^*$ ; it is an ideal of  $\widehat{A^*}$ .
- $\widehat{A^*}$  plays a crucial role in automata theory because it is the Zariski spectrum of the Boolean ring of regular languages.

# The profinite completion of the free monoid

- For  $u \neq v \in A^*$ , define  $\sigma(u, v)$  to be the minimum size of a finite monoid separating  $u$  from  $v$ .
- $A^*$  is residually finite, so  $\sigma(u, v)$  is well defined.
- The **profinite ultrametric** on  $A^*$  is defined by

$$d(u, v) = |A|^{-\sigma(u, v)}.$$

- The completion is the **free profinite monoid**  $\widehat{A^*}$ .
- Set  $\partial\widehat{A^*} = \widehat{A^*} \setminus A^*$ ; it is an ideal of  $\widehat{A^*}$ .
- $\widehat{A^*}$  plays a crucial role in automata theory because it is the **Zariski spectrum** of the Boolean ring of regular languages.

# The profinite completion of the free monoid

- For  $u \neq v \in A^*$ , define  $\sigma(u, v)$  to be the minimum size of a finite monoid separating  $u$  from  $v$ .
- $A^*$  is residually finite, so  $\sigma(u, v)$  is well defined.
- The **profinite ultrametric** on  $A^*$  is defined by

$$d(u, v) = |A|^{-\sigma(u, v)}.$$

- The completion is the **free profinite monoid**  $\widehat{A^*}$ .
- Set  $\partial\widehat{A^*} = \widehat{A^*} \setminus A^*$ ; it is an ideal of  $\widehat{A^*}$ .
- $\widehat{A^*}$  plays a crucial role in automata theory because it is the **Zariski spectrum** of the Boolean ring of regular languages.

# The profinite completion of the free monoid

- For  $u \neq v \in A^*$ , define  $\sigma(u, v)$  to be the minimum size of a finite monoid separating  $u$  from  $v$ .
- $A^*$  is residually finite, so  $\sigma(u, v)$  is well defined.
- The **profinite ultrametric** on  $A^*$  is defined by

$$d(u, v) = |A|^{-\sigma(u, v)}.$$

- The completion is the **free profinite monoid**  $\widehat{A^*}$ .
- Set  $\partial\widehat{A^*} = \widehat{A^*} \setminus A^*$ ; it is an ideal of  $\widehat{A^*}$ .
- $\widehat{A^*}$  plays a crucial role in automata theory because it is the **Zariski spectrum** of the Boolean ring of regular languages.

# The profinite completion of the free monoid

- For  $u \neq v \in A^*$ , define  $\sigma(u, v)$  to be the minimum size of a finite monoid separating  $u$  from  $v$ .
- $A^*$  is residually finite, so  $\sigma(u, v)$  is well defined.
- The **profinite ultrametric** on  $A^*$  is defined by

$$d(u, v) = |A|^{-\sigma(u, v)}.$$

- The completion is the **free profinite monoid**  $\widehat{A^*}$ .
- Set  $\partial\widehat{A^*} = \widehat{A^*} \setminus A^*$ ; it is an ideal of  $\widehat{A^*}$ .
- $\widehat{A^*}$  plays a crucial role in automata theory because it is the **Zariski spectrum** of the Boolean ring of regular languages.

# The Almeida dictionary

- Let  $\mathcal{X} \subseteq A^\omega$  be an irreducible subshift.
- Almeida established the following result:

## Theorem (Almeida)

- *The map  $\mathcal{X} \mapsto \overline{L(\mathcal{X})} \cap \partial \widehat{A}^*$  is injective.*
- *Among all principal ideals  $\widehat{A}^* u \widehat{A}^*$  intersecting  $\overline{L(\mathcal{X})}$  there is a unique minimal one, denoted  $I(\mathcal{X})$ .*
- *$I(\mathcal{X})$  can be generated by an idempotent  $e = e^2$ .*
- *The subshift  $\mathcal{X}$  is minimal if and only if  $I(\mathcal{X})$  is maximal among principal ideals in  $\partial \widehat{A}^*$ .*
- *Every maximal principal ideal in  $\partial \widehat{A}^*$  is of the form  $I(\mathcal{X})$  for a unique minimal subshift  $\mathcal{X}$ .*



# The Almeida dictionary

- Let  $\mathcal{X} \subseteq A^\omega$  be an irreducible subshift.
- Almeida established the following result:

## Theorem (Almeida)

- *The map  $\mathcal{X} \mapsto \overline{L(\mathcal{X})} \cap \partial \widehat{A}^*$  is injective.*
- *Among all principal ideals  $\widehat{A}^* u \widehat{A}^*$  intersecting  $\overline{L(\mathcal{X})}$  there is a unique minimal one, denoted  $I(\mathcal{X})$ .*
- *$I(\mathcal{X})$  can be generated by an idempotent  $e = e^2$ .*
- *The subshift  $\mathcal{X}$  is minimal if and only if  $I(\mathcal{X})$  is maximal among principal ideals in  $\partial \widehat{A}^*$ .*
- *Every maximal principal ideal in  $\partial \widehat{A}^*$  is of the form  $I(\mathcal{X})$  for a unique minimal subshift  $\mathcal{X}$ .*

# The Almeida dictionary

- Let  $\mathcal{X} \subseteq A^\omega$  be an irreducible subshift.
- Almeida established the following result:

## Theorem (Almeida)

- *The map  $\mathcal{X} \mapsto \overline{L(\mathcal{X})} \cap \partial \widehat{A}^*$  is injective.*
- *Among all principal ideals  $\widehat{A}^* u \widehat{A}^*$  intersecting  $\overline{L(\mathcal{X})}$  there is a unique minimal one, denoted  $I(\mathcal{X})$ .*
- *$I(\mathcal{X})$  can be generated by an idempotent  $e = e^2$ .*
- *The subshift  $\mathcal{X}$  is minimal if and only if  $I(\mathcal{X})$  is maximal among principal ideals in  $\partial \widehat{A}^*$ .*
- *Every maximal principal ideal in  $\partial \widehat{A}^*$  is of the form  $I(\mathcal{X})$  for a unique minimal subshift  $\mathcal{X}$ .*

# The Almeida dictionary

- Let  $\mathcal{X} \subseteq A^\omega$  be an irreducible subshift.
- Almeida established the following result:

## Theorem (Almeida)

- *The map  $\mathcal{X} \mapsto \overline{L(\mathcal{X})} \cap \partial \widehat{A}^*$  is injective.*
- *Among all principal ideals  $\widehat{A}^* u \widehat{A}^*$  intersecting  $\overline{L(\mathcal{X})}$  there is a unique minimal one, denoted  $I(\mathcal{X})$ .*
- *$I(\mathcal{X})$  can be generated by an idempotent  $e = e^2$ .*
- *The subshift  $\mathcal{X}$  is minimal if and only if  $I(\mathcal{X})$  is maximal among principal ideals in  $\partial \widehat{A}^*$ .*
- *Every maximal principal ideal in  $\partial \widehat{A}^*$  is of the form  $I(\mathcal{X})$  for a unique minimal subshift  $\mathcal{X}$ .*

# The Almeida dictionary

- Let  $\mathcal{X} \subseteq A^\omega$  be an irreducible subshift.
- Almeida established the following result:

## Theorem (Almeida)

- *The map  $\mathcal{X} \mapsto \overline{L(\mathcal{X})} \cap \partial \widehat{A}^*$  is injective.*
- *Among all principal ideals  $\widehat{A}^* u \widehat{A}^*$  intersecting  $\overline{L(\mathcal{X})}$  there is a unique minimal one, denoted  $I(\mathcal{X})$ .*
- *$I(\mathcal{X})$  can be generated by an idempotent  $e = e^2$ .*
- *The subshift  $\mathcal{X}$  is minimal if and only if  $I(\mathcal{X})$  is maximal among principal ideals in  $\partial \widehat{A}^*$ .*
- *Every maximal principal ideal in  $\partial \widehat{A}^*$  is of the form  $I(\mathcal{X})$  for a unique minimal subshift  $\mathcal{X}$ .*

# The Almeida dictionary

- Let  $\mathcal{X} \subseteq A^\omega$  be an irreducible subshift.
- Almeida established the following result:

## Theorem (Almeida)

- *The map  $\mathcal{X} \mapsto \overline{L(\mathcal{X})} \cap \partial \widehat{A}^*$  is injective.*
- *Among all principal ideals  $\widehat{A}^* u \widehat{A}^*$  intersecting  $\overline{L(\mathcal{X})}$  there is a unique minimal one, denoted  $I(\mathcal{X})$ .*
- *$I(\mathcal{X})$  can be generated by an idempotent  $e = e^2$ .*
- *The subshift  $\mathcal{X}$  is minimal if and only if  $I(\mathcal{X})$  is maximal among principal ideals in  $\partial \widehat{A}^*$ .*
- *Every maximal principal ideal in  $\partial \widehat{A}^*$  is of the form  $I(\mathcal{X})$  for a unique minimal subshift  $\mathcal{X}$ .*

# The Almeida dictionary

- Let  $\mathcal{X} \subseteq A^\omega$  be an irreducible subshift.
- Almeida established the following result:

## Theorem (Almeida)

- *The map  $\mathcal{X} \mapsto \overline{L(\mathcal{X})} \cap \partial \widehat{A}^*$  is injective.*
- *Among all principal ideals  $\widehat{A}^* u \widehat{A}^*$  intersecting  $\overline{L(\mathcal{X})}$  there is a unique minimal one, denoted  $I(\mathcal{X})$ .*
- *$I(\mathcal{X})$  can be generated by an idempotent  $e = e^2$ .*
- *The subshift  $\mathcal{X}$  is minimal if and only if  $I(\mathcal{X})$  is maximal among principal ideals in  $\partial \widehat{A}^*$ .*
- *Every maximal principal ideal in  $\partial \widehat{A}^*$  is of the form  $I(\mathcal{X})$  for a unique minimal subshift  $\mathcal{X}$ .*

# Maximal subgroups of a profinite monoid

- Suppose  $M$  is a profinite monoid and  $e \in M$  is an idempotent.
- Then  $eMe$  is a profinite monoid with identity  $e$ .
- The group of units  $G_e$  of  $eMe$  is a profinite group known as the **maximal subgroup at  $e$** .
- If  $e, f$  are idempotents, then  $MeM = MfM$  implies  $eMe \cong fMf$  and hence  $G_e \cong G_f$ .
- Thus to each principal ideal generated by an idempotent, we can associate a unique maximal subgroup (up to isomorphism).
- I've shown that any ideal of  $\widehat{A}^*$  generated by an idempotent is prime; in particular, the ideal  $I(\mathcal{X})$  associated to an irreducible shift  $\mathcal{X}$  is prime.

# Maximal subgroups of a profinite monoid

- Suppose  $M$  is a profinite monoid and  $e \in M$  is an idempotent.
- Then  $eMe$  is a profinite monoid with identity  $e$ .
- The group of units  $G_e$  of  $eMe$  is a profinite group known as the **maximal subgroup at  $e$** .
- If  $e, f$  are idempotents, then  $MeM = MfM$  implies  $eMe \cong fMf$  and hence  $G_e \cong G_f$ .
- Thus to each principal ideal generated by an idempotent, we can associate a **unique maximal subgroup** (up to isomorphism).
- I've shown that any ideal of  $\widehat{A}^*$  generated by an idempotent is prime; in particular, the ideal  $I(\mathcal{X})$  associated to an irreducible shift  $\mathcal{X}$  is prime.



# Maximal subgroups of a profinite monoid

- Suppose  $M$  is a profinite monoid and  $e \in M$  is an idempotent.
- Then  $eMe$  is a profinite monoid with identity  $e$ .
- The group of units  $G_e$  of  $eMe$  is a profinite group known as the **maximal subgroup at  $e$** .
- If  $e, f$  are idempotents, then  $MeM = MfM$  implies  $eMe \cong fMf$  and hence  $G_e \cong G_f$ .
- Thus to each principal ideal generated by an idempotent, we can associate a **unique** maximal subgroup (up to isomorphism).
- I've shown that any ideal of  $\widehat{A}^*$  generated by an idempotent is **prime**; in particular, the ideal  $I(\mathcal{X})$  associated to an irreducible shift  $\mathcal{X}$  is prime.

# Maximal subgroups of a profinite monoid

- Suppose  $M$  is a profinite monoid and  $e \in M$  is an idempotent.
- Then  $eMe$  is a profinite monoid with identity  $e$ .
- The group of units  $G_e$  of  $eMe$  is a profinite group known as the **maximal subgroup at  $e$** .
- If  $e, f$  are idempotents, then  $MeM = MfM$  implies  $eMe \cong fMf$  and hence  $G_e \cong G_f$ .
- Thus to each principal ideal generated by an idempotent, we can associate a **unique** maximal subgroup (up to isomorphism).
- I've shown that any ideal of  $\widehat{A}^*$  generated by an idempotent is **prime**; in particular, the ideal  $I(\mathcal{X})$  associated to an irreducible shift  $\mathcal{X}$  is prime.

# Maximal subgroups of a profinite monoid

- Suppose  $M$  is a profinite monoid and  $e \in M$  is an idempotent.
- Then  $eMe$  is a profinite monoid with identity  $e$ .
- The group of units  $G_e$  of  $eMe$  is a profinite group known as the **maximal subgroup at  $e$** .
- If  $e, f$  are idempotents, then  $MeM = MfM$  implies  $eMe \cong fMf$  and hence  $G_e \cong G_f$ .
- Thus to each principal ideal generated by an idempotent, we can associate a **unique** maximal subgroup (up to isomorphism).
- I've shown that any ideal of  $\widehat{A}^*$  generated by an idempotent is **prime**; in particular, the ideal  $I(\mathcal{X})$  associated to an irreducible shift  $\mathcal{X}$  is prime.

# Maximal subgroups of a profinite monoid

- Suppose  $M$  is a profinite monoid and  $e \in M$  is an idempotent.
- Then  $eMe$  is a profinite monoid with identity  $e$ .
- The group of units  $G_e$  of  $eMe$  is a profinite group known as the **maximal subgroup at  $e$** .
- If  $e, f$  are idempotents, then  $MeM = MfM$  implies  $eMe \cong fMf$  and hence  $G_e \cong G_f$ .
- Thus to each principal ideal generated by an idempotent, we can associate a **unique** maximal subgroup (up to isomorphism).
- I've shown that any ideal of  $\widehat{A}^*$  generated by an idempotent is **prime**; in particular, the ideal  $I(\mathcal{X})$  associated to an irreducible shift  $\mathcal{X}$  is prime.

# The profinite group associated to an irreducible subshift

- Since the ideal  $I(\mathcal{X})$  associated to an irreducible subshift  $\mathcal{X}$  is generated by an idempotent, it has a unique maximal subgroup  $G(\mathcal{X})$  associated to it.
- Almeida defined this to be the **profinite group associated to the subshift  $\mathcal{X}$** .
- He announced that it is a conjugacy invariant of the sofic shift.
- A proof was first published by his student, A. Costa.
- So there is a profinite group invariant associated to an irreducible shift via the free profinite monoid!

# The profinite group associated to an irreducible subshift

- Since the ideal  $I(\mathcal{X})$  associated to an irreducible subshift  $\mathcal{X}$  is generated by an idempotent, it has a unique maximal subgroup  $G(\mathcal{X})$  associated to it.
- Almeida defined this to be the **profinite group associated to the subshift  $\mathcal{X}$** .
- He announced that it is a conjugacy invariant of the sofic shift.
- A proof was first published by his student, A. Costa.
- So there is a profinite group invariant associated to an irreducible shift via the free profinite monoid!

# The profinite group associated to an irreducible subshift

- Since the ideal  $I(\mathcal{X})$  associated to an irreducible subshift  $\mathcal{X}$  is generated by an idempotent, it has a unique maximal subgroup  $G(\mathcal{X})$  associated to it.
- Almeida defined this to be the **profinite group associated to the subshift  $\mathcal{X}$** .
- He announced that it is a conjugacy invariant of the sofic shift.
- A proof was first published by his student, A. Costa.
- So there is a profinite group invariant associated to an irreducible shift via the free profinite monoid!

# The profinite group associated to an irreducible subshift

- Since the ideal  $I(\mathcal{X})$  associated to an irreducible subshift  $\mathcal{X}$  is generated by an idempotent, it has a unique maximal subgroup  $G(\mathcal{X})$  associated to it.
- Almeida defined this to be the **profinite group associated to the subshift  $\mathcal{X}$** .
- He announced that it is a conjugacy invariant of the sofic shift.
- A proof was first published by his student, A. Costa.
- So there is a profinite group invariant associated to an irreducible shift via the free profinite monoid!



# The profinite group associated to an irreducible subshift

- Since the ideal  $I(\mathcal{X})$  associated to an irreducible subshift  $\mathcal{X}$  is generated by an idempotent, it has a unique maximal subgroup  $G(\mathcal{X})$  associated to it.
- Almeida defined this to be the **profinite group associated to the subshift  $\mathcal{X}$** .
- He announced that it is a conjugacy invariant of the sofic shift.
- A proof was first published by his student, A. Costa.
- So there is a profinite group invariant associated to an irreducible shift via the free profinite monoid!

# Maximal subgroups of $\widehat{A}^*$

- Let  $\widehat{F}_A$  be a **free profinite group** on  $A$ .
- There is a natural surjective map  $\varphi: \widehat{A}^* \rightarrow \widehat{F}_A$ .
- If  $e$  is an idempotent of the minimal ideal  $I$ , then  $e\widehat{A}^*e = G_e$  and  $\varphi(G_e) = \widehat{F}_A$ .
- So  $\varphi$  splits and hence all projective profinite groups embed in free profinite monoids (observation of Almeida and Volkov).
- Margolis and I observed that the maximal subgroup of the minimal ideal maps onto any countably based profinite group.

## Question (Margolis 97)

- ① *Is every maximal subgroup of  $\widehat{A}^*$  a free profinite group, or at least projective?*
- *Is the maximal subgroup of the minimal ideal of  $\widehat{A}^*$  a free profinite group?*

# Maximal subgroups of $\widehat{A}^*$

- Let  $\widehat{F}_A$  be a **free profinite group** on  $A$ .
- There is a natural surjective map  $\varphi: \widehat{A}^* \rightarrow \widehat{F}_A$ .
- If  $e$  is an idempotent of the minimal ideal  $I$ , then  $e\widehat{A}^*e = G_e$  and  $\varphi(G_e) = \widehat{F}_A$ .
- So  $\varphi$  splits and hence all projective profinite groups embed in free profinite monoids (observation of Almeida and Volkov).
- Margolis and I observed that the maximal subgroup of the minimal ideal maps onto any countably based profinite group.

## Question (Margolis 97)

- ① *Is every maximal subgroup of  $\widehat{A}^*$  a free profinite group, or at least projective?*
- *Is the maximal subgroup of the minimal ideal of  $\widehat{A}^*$  a free profinite group?*

# Maximal subgroups of $\widehat{A^*}$

- Let  $\widehat{F}_A$  be a **free profinite group** on  $A$ .
- There is a natural surjective map  $\varphi: \widehat{A^*} \rightarrow \widehat{F}_A$ .
- If  $e$  is an idempotent of the minimal ideal  $I$ , then  $e\widehat{A^*}e = G_e$  and  $\varphi(G_e) = \widehat{F}_A$ .
- So  $\varphi$  splits and hence all projective profinite groups embed in free profinite monoids (observation of Almeida and Volkov).
- Margolis and I observed that the maximal subgroup of the minimal ideal maps onto any countably based profinite group.

## Question (Margolis 97)

- 1 *Is every maximal subgroup of  $\widehat{A^*}$  a free profinite group, or at least projective?*
- 2 *Is the maximal subgroup of the minimal ideal of  $\widehat{A^*}$  a free profinite group?*

# Maximal subgroups of $\widehat{A^*}$

- Let  $\widehat{F}_A$  be a **free profinite group** on  $A$ .
- There is a natural surjective map  $\varphi: \widehat{A^*} \rightarrow \widehat{F}_A$ .
- If  $e$  is an idempotent of the minimal ideal  $I$ , then  $e\widehat{A^*}e = G_e$  and  $\varphi(G_e) = \widehat{F}_A$ .
- So  $\varphi$  splits and hence all projective profinite groups embed in free profinite monoids (observation of Almeida and Volkov).
- Margolis and I observed that the maximal subgroup of the minimal ideal maps onto any countably based profinite group.

## Question (Margolis 97)

- 1 *Is every maximal subgroup of  $\widehat{A^*}$  a free profinite group, or at least projective?*
- 2 *Is the maximal subgroup of the minimal ideal of  $\widehat{A^*}$  a free profinite group?*

# Maximal subgroups of $\widehat{A}^*$

- Let  $\widehat{F}_A$  be a **free profinite group** on  $A$ .
- There is a natural surjective map  $\varphi: \widehat{A}^* \rightarrow \widehat{F}_A$ .
- If  $e$  is an idempotent of the minimal ideal  $I$ , then  $e\widehat{A}^*e = G_e$  and  $\varphi(G_e) = \widehat{F}_A$ .
- So  $\varphi$  splits and hence all projective profinite groups embed in free profinite monoids (observation of Almeida and Volkov).
- Margolis and I observed that the maximal subgroup of the minimal ideal maps onto any countably based profinite group.

## Question (Margolis 97)

- 1 *Is every maximal subgroup of  $\widehat{A}^*$  a free profinite group, or at least projective?*
- 2 *Is the maximal subgroup of the minimal ideal of  $\widehat{A}^*$  a free profinite group?*

# Maximal subgroups of $\widehat{A^*}$

- Let  $\widehat{F}_A$  be a **free profinite group** on  $A$ .
- There is a natural surjective map  $\varphi: \widehat{A^*} \rightarrow \widehat{F}_A$ .
- If  $e$  is an idempotent of the minimal ideal  $I$ , then  $e\widehat{A^*}e = G_e$  and  $\varphi(G_e) = \widehat{F}_A$ .
- So  $\varphi$  splits and hence all projective profinite groups embed in free profinite monoids (observation of Almeida and Volkov).
- Margolis and I observed that the maximal subgroup of the minimal ideal maps onto any countably based profinite group.

## Question (Margolis 97)

- 1 *Is every maximal subgroup of  $\widehat{A^*}$  a free profinite group, or at least projective?*
- 2 *Is the maximal subgroup of the minimal ideal of  $\widehat{A^*}$  a free profinite group?*

# Maximal subgroups of $\widehat{A^*}$

- Let  $\widehat{F}_A$  be a **free profinite group** on  $A$ .
- There is a natural surjective map  $\varphi: \widehat{A^*} \rightarrow \widehat{F}_A$ .
- If  $e$  is an idempotent of the minimal ideal  $I$ , then  $e\widehat{A^*}e = G_e$  and  $\varphi(G_e) = \widehat{F}_A$ .
- So  $\varphi$  splits and hence all projective profinite groups embed in free profinite monoids (observation of Almeida and Volkov).
- Margolis and I observed that the maximal subgroup of the minimal ideal maps onto any countably based profinite group.

## Question (Margolis 97)

- 1 *Is every maximal subgroup of  $\widehat{A^*}$  a free profinite group, or at least projective?*
- 2 *Is the maximal subgroup of the minimal ideal of  $\widehat{A^*}$  a free profinite group?*



# Projective profinite groups

- A profinite group  $G$  is **projective** if given a diagram

$$\begin{array}{ccc} & & G \\ & & \downarrow \varphi \\ A & \xrightarrow{\alpha} & B \end{array}$$

of epimorphisms of profinite groups, there exists a homomorphism  $\lambda: G \rightarrow A$  so that the diagram commutes.

- Projective profinite groups turn out to be precisely the closed subgroups of free profinite groups.

# Projective profinite groups

- A profinite group  $G$  is **projective** if given a diagram


$$\begin{array}{ccc} & & G \\ & & \downarrow \varphi \\ A & \xrightarrow{\alpha} & B \end{array}$$

of epimorphisms of profinite groups, there exists a **homomorphism**  $\lambda: G \rightarrow A$  so that the diagram commutes.

- Projective profinite groups turn out to be precisely the closed subgroups of free profinite groups.

# Projective profinite groups

- A profinite group  $G$  is **projective** if given a diagram




of epimorphisms of profinite groups, there exists a **homomorphism**  $\lambda: G \rightarrow A$  so that the diagram commutes.

- Projective profinite groups turn out to be precisely the closed subgroups of free profinite groups.

# Projective profinite groups

- A profinite group  $G$  is **projective** if given a diagram



of epimorphisms of profinite groups, there exists a **homomorphism**  $\lambda: G \rightarrow A$  so that the diagram commutes.

- Projective profinite groups turn out to be precisely the closed subgroups of free profinite groups.

# Profinite groups associated to minimal subshifts

- Almeida and Volkov showed that the group associated to a **periodic** subshift is free procyclic.
- Almeida used techniques from symbolic dynamics to study the group associated to a primitive substitution.

## Theorem (Almeida)

*Let  $f: A^* \rightarrow A^*$  be a primitive substitution which is invertible over the free group  $F_A$ . Then the profinite group associated to the corresponding minimal subshift is a free profinite group on  $A$ .*

- For example, the **Fibonacci word** is the fixed point of the invertible substitution  $a \mapsto ab, b \mapsto a$ . The associated maximal subgroup is then free of rank 2.
- More generally, the group associated to any Sturmian system is free of rank 2.

# Profinite groups associated to minimal subshifts

- Almeida and Volkov showed that the group associated to a **periodic** subshift is free procyclic.
- Almeida used techniques from symbolic dynamics to study the group associated to a primitive substitution.

## Theorem (Almeida)

*Let  $f: A^* \rightarrow A^*$  be a primitive substitution which is invertible over the free group  $F_A$ . Then the profinite group associated to the corresponding minimal subshift is a free profinite group on  $A$ .*

- For example, the **Fibonacci word** is the fixed point of the invertible substitution  $a \mapsto ab, b \mapsto a$ . The associated maximal subgroup is then free of rank 2.
- More generally, the group associated to any Sturmian system is free of rank 2.

# Profinite groups associated to minimal subshifts

- Almeida and Volkov showed that the group associated to a **periodic** subshift is free procyclic.
- Almeida used techniques from symbolic dynamics to study the group associated to a primitive substitution.

## Theorem (Almeida)

*Let  $f: A^* \rightarrow A^*$  be a primitive substitution which is invertible over the free group  $F_A$ . Then the profinite group associated to the corresponding minimal subshift is a free profinite group on  $A$ .*

- For example, the **Fibonacci word** is the fixed point of the invertible substitution  $a \mapsto ab, b \mapsto a$ . The associated maximal subgroup is then free of rank 2.
- More generally, the group associated to any Sturmian system is free of rank 2.

# Profinite groups associated to minimal subshifts

- Almeida and Volkov showed that the group associated to a **periodic** subshift is free procyclic.
- Almeida used techniques from symbolic dynamics to study the group associated to a primitive substitution.

## Theorem (Almeida)

*Let  $f: A^* \rightarrow A^*$  be a primitive substitution which is invertible over the free group  $F_A$ . Then the profinite group associated to the corresponding minimal subshift is a free profinite group on  $A$ .*

- For example, the **Fibonacci word** is the fixed point of the invertible substitution  $a \mapsto ab, b \mapsto a$ . The associated maximal subgroup is then free of rank 2.
- More generally, the group associated to any Sturmian system is free of rank 2.



# Profinite groups associated to minimal subshifts

- Almeida and Volkov showed that the group associated to a **periodic** subshift is free procyclic.
- Almeida used techniques from symbolic dynamics to study the group associated to a primitive substitution.

## Theorem (Almeida)

*Let  $f: A^* \rightarrow A^*$  be a primitive substitution which is invertible over the free group  $F_A$ . Then the profinite group associated to the corresponding minimal subshift is a free profinite group on  $A$ .*

- For example, the **Fibonacci word** is the fixed point of the invertible substitution  $a \mapsto ab, b \mapsto a$ . The associated maximal subgroup is then free of rank 2.
- More generally, the group associated to any Sturmian system is free of rank 2.

# Non-free subgroups

- Almeida showed the profinite group associated to the substitution  $a \mapsto a^3b, b \mapsto ab$  is not free.
- Almeida presented this work at the Fields workshop on profinite groups organized by me and Ribes in 2005.
- Lubotzky asked during the questions after Almeida's talk whether all maximal groups of  $\widehat{A^*}$  are projective.
- Recently, Almeida and Costa showed that the profinite group associated to the Morse-Thue infinite word is not free profinite.

# Non-free subgroups

- Almeida showed the profinite group associated to the substitution  $a \mapsto a^3b, b \mapsto ab$  is not free.
- Almeida presented this work at the Fields workshop on profinite groups organized by me and Ribes in 2005.
- Lubotzky asked during the questions after Almeida's talk whether all maximal groups of  $\widehat{A^*}$  are projective.
- Recently, Almeida and Costa showed that the profinite group associated to the Morse-Thue infinite word is not free profinite.

# Non-free subgroups

- Almeida showed the profinite group associated to the substitution  $a \mapsto a^3b, b \mapsto ab$  is not free.
- Almeida presented this work at the Fields workshop on profinite groups organized by me and Ribes in 2005.
- Lubotzky asked during the questions after Almeida's talk whether all maximal groups of  $\widehat{A^*}$  are projective.
- Recently, Almeida and Costa showed that the profinite group associated to the Morse-Thue infinite word is not free profinite.

# Non-free subgroups

- Almeida showed the profinite group associated to the substitution  $a \mapsto a^3b, b \mapsto ab$  is not free.
- Almeida presented this work at the Fields workshop on profinite groups organized by me and Ribes in 2005.
- Lubotzky asked during the questions after Almeida's talk whether all maximal groups of  $\widehat{A^*}$  are projective.
- Recently, Almeida and Costa showed that the profinite group associated to the Morse-Thue infinite word is not free profinite.

# Maximal subgroups are projective

## Theorem (Rhodes, BS)

*The closed subgroups of  $\widehat{A}^*$  are precisely the projective profinite groups.*

- The proof uses wreath products and Schützenberger's generalization of the Krasner-Kaloujnine embedding to monoids in order to extend maps from the maximal subgroup to the whole free profinite monoid.
- Ribes later pointed us to a similar proof scheme by Cossey, Kegel and Kovács for the case of free profinite groups.
- Ribes and I have used the same ideas to give simple algebraic proofs of the Nielsen-Schreier and Kurosh Theorems via wreath products (in both the abstract and profinite settings).

# Maximal subgroups are projective

## Theorem (Rhodes, BS)

*The closed subgroups of  $\widehat{A}^*$  are precisely the projective profinite groups.*

- The proof uses wreath products and Schützenberger's generalization of the Krasner-Kaloujnine embedding to monoids in order to extend maps from the maximal subgroup to the whole free profinite monoid.
- Ribes later pointed us to a similar proof scheme by Cossey, Kegel and Kovács for the case of free profinite groups.
- Ribes and I have used the same ideas to give simple algebraic proofs of the Nielsen-Schreier and Kurosh Theorems via wreath products (in both the abstract and profinite settings).

# Maximal subgroups are projective

## Theorem (Rhodes, BS)

*The closed subgroups of  $\widehat{A}^*$  are precisely the projective profinite groups.*

- The proof uses wreath products and Schützenberger's generalization of the Krasner-Kaloujnine embedding to monoids in order to extend maps from the maximal subgroup to the whole free profinite monoid.
- Ribes later pointed us to a similar proof scheme by Cossey, Kegel and Kovács for the case of free profinite groups.
- Ribes and I have used the same ideas to give simple algebraic proofs of the Nielsen-Schreier and Kurosh Theorems via wreath products (in both the abstract and profinite settings).



# Maximal subgroups are projective

## Theorem (Rhodes, BS)

*The closed subgroups of  $\widehat{A}^*$  are precisely the projective profinite groups.*

- The proof uses wreath products and Schützenberger's generalization of the Krasner-Kaloujnine embedding to monoids in order to extend maps from the maximal subgroup to the whole free profinite monoid.
- Ribes later pointed us to a similar proof scheme by Cossey, Kegel and Kovács for the case of free profinite groups.
- Ribes and I have used the same ideas to give simple algebraic proofs of the Nielsen-Schreier and Kurosh Theorems via wreath products (in both the abstract and profinite settings).

# Torsion in free profinite monoids

- Any element  $s$  of finite order in  $\widehat{A}^*$  must satisfy  $s^n = s^{n+m}$  for some  $n, m \geq 1$ .
- But then  $C = \{s^n, \dots, s^{n+m-1}\}$  is a finite cyclic subgroup with identity  $s^k$  some  $n \leq k \leq n+m-1$ .
- But then, since idempotents generate a prime ideal, it follows that  $s$  and  $s^k$  generate the same ideal.
- Standard profinite semigroup theory then implies  $s \in C$ .
- But projective profinite groups are torsion-free so  $C = \{s^k\}$ .

Theorem (Rhodes, BS)

*Every element of finite order in  $\widehat{A}^*$  is an idempotent.*

# Torsion in free profinite monoids

- Any element  $s$  of finite order in  $\widehat{A}^*$  must satisfy  $s^n = s^{n+m}$  for some  $n, m \geq 1$ .
- But then  $C = \{s^n, \dots, s^{n+m-1}\}$  is a finite cyclic subgroup with identity  $s^k$  some  $n \leq k \leq n+m-1$ .
- But then, since idempotents generate a prime ideal, it follows that  $s$  and  $s^k$  generate the same ideal.
- Standard profinite semigroup theory then implies  $s \in C$ .
- But projective profinite groups are torsion-free so  $C = \{s^k\}$ .

Theorem (Rhodes, BS)

*Every element of finite order in  $\widehat{A}^*$  is an idempotent.*

# Torsion in free profinite monoids

- Any element  $s$  of finite order in  $\widehat{A}^*$  must satisfy  $s^n = s^{n+m}$  for some  $n, m \geq 1$ .
- But then  $C = \{s^n, \dots, s^{n+m-1}\}$  is a finite cyclic subgroup with identity  $s^k$  some  $n \leq k \leq n+m-1$ .
- But then, since idempotents generate a prime ideal, it follows that  $s$  and  $s^k$  generate the same ideal.
- Standard profinite semigroup theory then implies  $s \in C$ .
- But projective profinite groups are torsion-free so  $C = \{s^k\}$ .

Theorem (Rhodes, BS)

*Every element of finite order in  $\widehat{A}^*$  is an idempotent.*

# Torsion in free profinite monoids

- Any element  $s$  of finite order in  $\widehat{A}^*$  must satisfy  $s^n = s^{n+m}$  for some  $n, m \geq 1$ .
- But then  $C = \{s^n, \dots, s^{n+m-1}\}$  is a finite cyclic subgroup with identity  $s^k$  some  $n \leq k \leq n + m - 1$ .
- But then, since idempotents generate a prime ideal, it follows that  $s$  and  $s^k$  generate the same ideal.
- Standard profinite semigroup theory then implies  $s \in C$ .
- But projective profinite groups are torsion-free so  $C = \{s^k\}$ .

Theorem (Rhodes, BS)

*Every element of finite order in  $\widehat{A}^*$  is an idempotent.*

# Torsion in free profinite monoids

- Any element  $s$  of finite order in  $\widehat{A}^*$  must satisfy  $s^n = s^{n+m}$  for some  $n, m \geq 1$ .
- But then  $C = \{s^n, \dots, s^{n+m-1}\}$  is a finite cyclic subgroup with identity  $s^k$  some  $n \leq k \leq n + m - 1$ .
- But then, since idempotents generate a prime ideal, it follows that  $s$  and  $s^k$  generate the same ideal.
- Standard profinite semigroup theory then implies  $s \in C$ .
- But projective profinite groups are torsion-free so  $C = \{s^k\}$ .

Theorem (Rhodes, BS)

*Every element of finite order in  $\widehat{A}^*$  is an idempotent.*

# Torsion in free profinite monoids

- Any element  $s$  of finite order in  $\widehat{A}^*$  must satisfy  $s^n = s^{n+m}$  for some  $n, m \geq 1$ .
- But then  $C = \{s^n, \dots, s^{n+m-1}\}$  is a finite cyclic subgroup with identity  $s^k$  some  $n \leq k \leq n + m - 1$ .
- But then, since idempotents generate a prime ideal, it follows that  $s$  and  $s^k$  generate the same ideal.
- Standard profinite semigroup theory then implies  $s \in C$ .
- But projective profinite groups are torsion-free so  $C = \{s^k\}$ .

## Theorem (Rhodes, BS)

*Every element of finite order in  $\widehat{A}^*$  is an idempotent.*

# The free profinite group on a set converging to 1

- A subset  $Y$  of a profinite group  $G$  is a **set of generators converging to 1** if:
  - $\overline{\langle Y \rangle} = G$ ;
  - Each neighborhood of 1 contains all but finitely many elements of  $Y$ .
- One can define a free profinite group  $\hat{F}_Y$  on a set  $Y$  of generators converging to 1. The cardinality of  $Y$  is called the rank of  $\hat{F}_Y$ .
- A free profinite group on a topological space  $X$  is also free on a set of generators converging to 1 of the same cardinality as the Boolean algebra of clopen subsets of  $X$ .



# The free profinite group on a set converging to 1

- A subset  $Y$  of a profinite group  $G$  is a **set of generators converging to 1** if:
  - $\overline{\langle Y \rangle} = G$ ;
  - Each neighborhood of 1 contains all but finitely many elements of  $Y$ .
- One can define a free profinite group  $\widehat{F}_Y$  on a set  $Y$  of generators converging to 1. The cardinality of  $Y$  is called the **rank** of  $\widehat{F}_Y$ .
- A free profinite group on a topological space  $X$  is also free on a set of generators converging to 1 of the same cardinality as the Boolean algebra of clopen subsets of  $X$ .

# The free profinite group on a set converging to 1

- A subset  $Y$  of a profinite group  $G$  is a **set of generators converging to 1** if:
  - $\overline{\langle Y \rangle} = G$ ;
  - Each neighborhood of 1 contains all but finitely many elements of  $Y$ .
- One can define a free profinite group  $\widehat{F}_Y$  on a set  $Y$  of generators converging to 1. The cardinality of  $Y$  is called the **rank** of  $\widehat{F}_Y$ .
- A free profinite group on a topological space  $X$  is also free on a set of generators converging to 1 of the same cardinality as the Boolean algebra of clopen subsets of  $X$ .

# The free profinite group on a set converging to 1

- A subset  $Y$  of a profinite group  $G$  is a **set of generators converging to 1** if:
  - $\overline{\langle Y \rangle} = G$ ;
  - Each neighborhood of 1 contains all but finitely many elements of  $Y$ .
- One can define a free profinite group  $\widehat{F}_Y$  on a set  $Y$  of generators converging to 1. The cardinality of  $Y$  is called the **rank** of  $\widehat{F}_Y$ .
- A free profinite group on a topological space  $X$  is also free on a set of generators converging to 1 of the same cardinality as the Boolean algebra of clopen subsets of  $X$ .

# The free profinite group on a set converging to 1

- A subset  $Y$  of a profinite group  $G$  is a **set of generators converging to 1** if:
  - $\overline{\langle Y \rangle} = G$ ;
  - Each neighborhood of 1 contains all but finitely many elements of  $Y$ .
- One can define a free profinite group  $\widehat{F}_Y$  on a set  $Y$  of generators converging to 1. The cardinality of  $Y$  is called the **rank** of  $\widehat{F}_Y$ .
- A free profinite group on a topological space  $X$  is also free on a set of generators converging to 1 of the same cardinality as the Boolean algebra of clopen subsets of  $X$ .

# The maximal subgroup of the minimal ideal is free

## Theorem (BS)

*The maximal subgroup of the minimal ideal of  $\widehat{A}^*$  is a free profinite group of countable rank.*

- The proof relies on **Iwasawa's criterion**: a countably based profinite group  $G$  is free of countable rank if and only if given a diagram

$$\begin{array}{ccc} & G & \\ & \downarrow \varphi & \\ A & \xrightarrow{\alpha} & B \end{array}$$

of epimorphisms ( $A$  and  $B$  are finite), there exists an epimorphism  $\lambda: G \twoheadrightarrow A$  so that the diagram commutes.

# The maximal subgroup of the minimal ideal is free

## Theorem (BS)

*The maximal subgroup of the minimal ideal of  $\widehat{A}^*$  is a free profinite group of countable rank.*

- The proof relies on **Iwasawa's criterion**: a countably based profinite group  $G$  is free of countable rank if and only if given a diagram

$$\begin{array}{ccc} & G & \\ & \downarrow \varphi & \\ A & \xrightarrow{\alpha} & B \end{array}$$

of epimorphisms ( $A$  and  $B$  are finite), there exists an **epimorphism**  $\lambda: G \twoheadrightarrow A$  so that the diagram commutes.

# The maximal subgroup of the minimal ideal is free

## Theorem (BS)

*The maximal subgroup of the minimal ideal of  $\widehat{A}^*$  is a free profinite group of countable rank.*

- The proof relies on **Iwasawa's criterion**: a countably based profinite group  $G$  is free of countable rank if and only if given a diagram

$$\begin{array}{ccc} & G & \\ & \downarrow \varphi & \\ A & \xrightarrow{\alpha} & B \end{array}$$


of epimorphisms ( $A$  and  $B$  are finite), there exists an **epimorphism**  $\lambda: G \twoheadrightarrow A$  so that the diagram commutes.

# The maximal subgroup of the minimal ideal is free

## Theorem (BS)

*The maximal subgroup of the minimal ideal of  $\widehat{A}^*$  is a free profinite group of countable rank.*

- The proof relies on **Iwasawa's criterion**: a countably based profinite group  $G$  is free of countable rank if and only if given a diagram



of epimorphisms ( $A$  and  $B$  are finite), there exists an **epimorphism**  $\lambda: G \twoheadrightarrow A$  so that the diagram commutes.



# The proof idea

- Again wreath products play a role: this time iterated wreath products.
- The idea is based on Bernhard Neumann's proof that every countable semigroup embeds in a 2-generated semigroup, and variations on this theme.
- The most relevant variant for us embeds any countable group as the maximal subgroup of the minimal ideal of a 2-generated monoid with cyclic group of units.
- Ideas from Krohn-Rhodes Theory and the Synthesis Theorem also play a role.

# The proof idea

- Again wreath products play a role: this time iterated wreath products.
- The idea is based on Bernhard Neumann's proof that every countable semigroup embeds in a 2-generated semigroup, and variations on this theme.
- The most relevant variant for us embeds any countable group as the maximal subgroup of the minimal ideal of a 2-generated monoid with cyclic group of units.
- Ideas from Krohn-Rhodes Theory and the Synthesis Theorem also play a role.

# The proof idea

- Again wreath products play a role: this time iterated wreath products.
- The idea is based on Bernhard Neumann's proof that every countable semigroup embeds in a 2-generated semigroup, and variations on this theme.
- The most relevant variant for us embeds any countable group as the maximal subgroup of the minimal ideal of a 2-generated monoid with cyclic group of units.
- Ideas from Krohn-Rhodes Theory and the Synthesis Theorem also play a role.

# The proof idea

- Again wreath products play a role: this time iterated wreath products.
- The idea is based on Bernhard Neumann's proof that every countable semigroup embeds in a 2-generated semigroup, and variations on this theme.
- The most relevant variant for us embeds any countable group as the maximal subgroup of the minimal ideal of a 2-generated monoid with cyclic group of units.
- Ideas from Krohn-Rhodes Theory and the Synthesis Theorem also play a role.

## Another free profinite subgroup

- Recall that we have a canonical projection  $\varphi: \widehat{A^*} \rightarrow \widehat{F_A}$  where  $\widehat{F_A}$  is the free profinite group generated by  $A$ .
- Moreover,  $\varphi$  restricts to an epimorphism  $\varphi: G \rightarrow \widehat{F_A}$  where  $G$  is the maximal subgroup of the minimal ideal  $I$ .
- Let  $K = \ker \varphi$ .

### Theorem (BS)

*The subgroup  $K$  is a free profinite group of countable rank.*

- The proof uses Melnikov's characterization of free normal subgroups of a free profinite group.
- The key point is that  $K$  can map onto every finite group.

# Another free profinite subgroup

- Recall that we have a canonical projection  $\varphi: \widehat{A^*} \rightarrow \widehat{F_A}$  where  $\widehat{F_A}$  is the free profinite group generated by  $A$ .
- Moreover,  $\varphi$  restricts to an epimorphism  $\varphi: G \rightarrow \widehat{F_A}$  where  $G$  is the maximal subgroup of the minimal ideal  $I$ .
- Let  $K = \ker \varphi$ .

## Theorem (BS)

*The subgroup  $K$  is a free profinite group of countable rank.*

- The proof uses Melnikov's characterization of free normal subgroups of a free profinite group.
- The key point is that  $K$  can map onto every finite group.

## Another free profinite subgroup

- Recall that we have a canonical projection  $\varphi: \widehat{A^*} \rightarrow \widehat{F_A}$  where  $\widehat{F_A}$  is the free profinite group generated by  $A$ .
- Moreover,  $\varphi$  restricts to an epimorphism  $\varphi: G \rightarrow \widehat{F_A}$  where  $G$  is the maximal subgroup of the minimal ideal  $I$ .
- Let  $K = \ker \varphi$ .

### Theorem (BS)

*The subgroup  $K$  is a free profinite group of countable rank.*

- The proof uses Melnikov's characterization of free normal subgroups of a free profinite group.
- The key point is that  $K$  can map onto every finite group.

## Another free profinite subgroup

- Recall that we have a canonical projection  $\varphi: \widehat{A^*} \rightarrow \widehat{F_A}$  where  $\widehat{F_A}$  is the free profinite group generated by  $A$ .
- Moreover,  $\varphi$  restricts to an epimorphism  $\varphi: G \rightarrow \widehat{F_A}$  where  $G$  is the maximal subgroup of the minimal ideal  $I$ .
- Let  $K = \ker \varphi$ .

### Theorem (BS)

*The subgroup  $K$  is a free profinite group of countable rank.*

- The proof uses Melnikov's characterization of free normal subgroups of a free profinite group.
- The key point is that  $K$  can map onto every finite group.



## Another free profinite subgroup

- Recall that we have a canonical projection  $\varphi: \widehat{A^*} \rightarrow \widehat{F_A}$  where  $\widehat{F_A}$  is the free profinite group generated by  $A$ .
- Moreover,  $\varphi$  restricts to an epimorphism  $\varphi: G \rightarrow \widehat{F_A}$  where  $G$  is the maximal subgroup of the minimal ideal  $I$ .
- Let  $K = \ker \varphi$ .

### Theorem (BS)

*The subgroup  $K$  is a free profinite group of countable rank.*

- The proof uses Melnikov's characterization of free normal subgroups of a free profinite group.
- The key point is that  $K$  can map onto every finite group.

## Another free profinite subgroup

- Recall that we have a canonical projection  $\varphi: \widehat{A}^* \rightarrow \widehat{F}_A$  where  $\widehat{F}_A$  is the free profinite group generated by  $A$ .
- Moreover,  $\varphi$  restricts to an epimorphism  $\varphi: G \rightarrow \widehat{F}_A$  where  $G$  is the maximal subgroup of the minimal ideal  $I$ .
- Let  $K = \ker \varphi$ .

### Theorem (BS)

*The subgroup  $K$  is a free profinite group of countable rank.*

- The proof uses Melnikov's characterization of free normal subgroups of a free profinite group.
- The key point is that  $K$  can map onto every finite group.

# Profinite groups associated to irreducible sofic subshifts

- The minimal ideal of  $\widehat{A^*}$  is the principal ideal associated to the **full shift**  $A^\omega$ , which is an irreducible sofic shift.
- It is then natural to ask whether the result for the minimal ideal extends to all irreducible sofic shifts.
- Minimal sofic shifts are periodic and hence have free procyclic associated groups.
- So the interesting case is the non-minimal case.

## Theorem (Costa, BS)

*The profinite group associated to a non-minimal irreducible sofic shift is a free profinite group of countable rank.*

- The proof makes use of the conjugacy invariance of the profinite group associated to an irreducible shift.

# Profinite groups associated to irreducible sofic subshifts

- The minimal ideal of  $\widehat{A^*}$  is the principal ideal associated to the **full shift**  $A^\omega$ , which is an irreducible sofic shift.
- It is then natural to ask whether the result for the minimal ideal extends to all irreducible sofic shifts.
- Minimal sofic shifts are periodic and hence have free procyclic associated groups.
- So the interesting case is the non-minimal case.

## Theorem (Costa, BS)

*The profinite group associated to a non-minimal irreducible sofic shift is a free profinite group of countable rank.*

- The proof makes use of the conjugacy invariance of the profinite group associated to an irreducible shift.

# Profinite groups associated to irreducible sofic subshifts

- The minimal ideal of  $\widehat{A^*}$  is the principal ideal associated to the **full shift**  $A^\omega$ , which is an irreducible sofic shift.
- It is then natural to ask whether the result for the minimal ideal extends to all irreducible sofic shifts.
- Minimal sofic shifts are periodic and hence have free procyclic associated groups.
- So the interesting case is the non-minimal case.

## Theorem (Costa, BS)

*The profinite group associated to a non-minimal irreducible sofic shift is a free profinite group of countable rank.*

- The proof makes use of the conjugacy invariance of the profinite group associated to an irreducible shift.

# Profinite groups associated to irreducible sofic subshifts

- The minimal ideal of  $\widehat{A^*}$  is the principal ideal associated to the **full shift**  $A^\omega$ , which is an irreducible sofic shift.
- It is then natural to ask whether the result for the minimal ideal extends to all irreducible sofic shifts.
- Minimal sofic shifts are periodic and hence have free procyclic associated groups.
- So the interesting case is the non-minimal case.

## Theorem (Costa, BS)

*The profinite group associated to a non-minimal irreducible sofic shift is a free profinite group of countable rank.*

- The proof makes use of the conjugacy invariance of the profinite group associated to an irreducible shift.

# Profinite groups associated to irreducible sofic subshifts

- The minimal ideal of  $\widehat{A^*}$  is the principal ideal associated to the **full shift**  $A^\omega$ , which is an irreducible sofic shift.
- It is then natural to ask whether the result for the minimal ideal extends to all irreducible sofic shifts.
- Minimal sofic shifts are periodic and hence have free procyclic associated groups.
- So the interesting case is the non-minimal case.

## Theorem (Costa, BS)

*The profinite group associated to a non-minimal irreducible sofic shift is a free profinite group of countable rank.*

- The proof makes use of the conjugacy invariance of the profinite group associated to an irreducible shift.

# Profinite groups associated to irreducible sofic subshifts

- The minimal ideal of  $\widehat{A^*}$  is the principal ideal associated to the **full shift**  $A^\omega$ , which is an irreducible sofic shift.
- It is then natural to ask whether the result for the minimal ideal extends to all irreducible sofic shifts.
- Minimal sofic shifts are periodic and hence have free procyclic associated groups.
- So the interesting case is the non-minimal case.

## Theorem (Costa, BS)

*The profinite group associated to a non-minimal irreducible sofic shift is a free profinite group of countable rank.*

- The proof makes use of the conjugacy invariance of the profinite group associated to an irreducible shift.



# Sofic shifts and generalized group mapping semigroups

- A key observation in the proof is that the following objects are the same:
  - Syntactic monoids of languages of the form  $L(\mathcal{X})$  with  $\mathcal{X}$  an irreducible sofic shift
  - Generalized group mapping monoids with an aperiodic 0-minimal ideal.
- A semigroup  $S$  is generalized group mapping if it has a (necessarily unique and regular) 0-minimal ideal  $I$  on which it acts faithfully on both the left and right.
- The so-called Fischer cover of an irreducible sofic shift is just the Schützenberger graph of the aperiodic 0-minimal ideal.
- Generalized group mapping semigroups were introduced by Rhodes in his work on the complexity of finite semigroups.
- The theory of lifting regular  $\mathcal{J}$ -classes, again due to Rhodes, plays a salient role in the proof of our theorem.

# Sofic shifts and generalized group mapping semigroups

- A key observation in the proof is that the following objects are the same:
  - **Syntactic monoids** of languages of the form  $L(\mathcal{X})$  with  $\mathcal{X}$  an irreducible sofic shift
  - **Generalized group mapping monoids** with an aperiodic 0-minimal ideal.
- A semigroup  $S$  is **generalized group mapping** if it has a (necessarily unique and regular) 0-minimal ideal  $I$  on which it acts faithfully on both the left and right.
- The so-called **Fischer cover** of an irreducible sofic shift is just the **Schützenberger graph** of the aperiodic 0-minimal ideal.
- Generalized group mapping semigroups were introduced by Rhodes in his work on the complexity of finite semigroups.
- The theory of lifting regular  $\mathcal{J}$ -classes, again due to Rhodes, plays a salient role in the proof of our theorem.

# Sofic shifts and generalized group mapping semigroups

- A key observation in the proof is that the following objects are the same:
  - **Syntactic monoids** of languages of the form  $L(\mathcal{X})$  with  $\mathcal{X}$  an irreducible sofic shift
  - **Generalized group mapping monoids** with an aperiodic 0-minimal ideal.
- A semigroup  $S$  is **generalized group mapping** if it has a (necessarily unique and regular) 0-minimal ideal  $I$  on which it acts faithfully on both the left and right.
- The so-called **Fischer cover** of an irreducible sofic shift is just the **Schützenberger graph** of the aperiodic 0-minimal ideal.
- Generalized group mapping semigroups were introduced by Rhodes in his work on the complexity of finite semigroups.
- The theory of lifting regular  $\mathcal{J}$ -classes, again due to Rhodes, plays a salient role in the proof of our theorem.

# Sofic shifts and generalized group mapping semigroups

- A key observation in the proof is that the following objects are the same:
  - **Syntactic monoids** of languages of the form  $L(\mathcal{X})$  with  $\mathcal{X}$  an irreducible sofic shift
  - **Generalized group mapping monoids** with an aperiodic 0-minimal ideal.
- A semigroup  $S$  is **generalized group mapping** if it has a (necessarily unique and regular) 0-minimal ideal  $I$  on which it acts faithfully on both the left and right.
- The so-called **Fischer cover** of an irreducible sofic shift is just the **Schützenberger graph** of the aperiodic 0-minimal ideal.
- Generalized group mapping semigroups were introduced by Rhodes in his work on the complexity of finite semigroups.
- The theory of lifting regular  $\mathcal{J}$ -classes, again due to Rhodes, plays a salient role in the proof of our theorem.

# Sofic shifts and generalized group mapping semigroups

- A key observation in the proof is that the following objects are the same:
  - **Syntactic monoids** of languages of the form  $L(\mathcal{X})$  with  $\mathcal{X}$  an irreducible sofic shift
  - **Generalized group mapping monoids** with an aperiodic 0-minimal ideal.
- A semigroup  $S$  is **generalized group mapping** if it has a (necessarily unique and regular) 0-minimal ideal  $I$  on which it acts faithfully on both the left and right.
- The so-called **Fischer cover** of an irreducible sofic shift is just the **Schützenberger graph** of the aperiodic 0-minimal ideal.
- Generalized group mapping semigroups were introduced by Rhodes in his work on the complexity of finite semigroups.
- The theory of lifting regular  $\mathcal{J}$ -classes, again due to Rhodes, plays a salient role in the proof of our theorem.

# Sofic shifts and generalized group mapping semigroups

- A key observation in the proof is that the following objects are the same:
  - **Syntactic monoids** of languages of the form  $L(\mathcal{X})$  with  $\mathcal{X}$  an irreducible sofic shift
  - **Generalized group mapping monoids** with an aperiodic 0-minimal ideal.
- A semigroup  $S$  is **generalized group mapping** if it has a (necessarily unique and regular) 0-minimal ideal  $I$  on which it acts faithfully on both the left and right.
- The so-called **Fischer cover** of an irreducible sofic shift is just the **Schützenberger graph** of the aperiodic 0-minimal ideal.
- Generalized group mapping semigroups were introduced by Rhodes in his work on the complexity of finite semigroups.
- The theory of lifting regular  $\mathcal{J}$ -classes, again due to Rhodes, plays a salient role in the proof of our theorem.

# Sofic shifts and generalized group mapping semigroups

- A key observation in the proof is that the following objects are the same:
  - **Syntactic monoids** of languages of the form  $L(\mathcal{X})$  with  $\mathcal{X}$  an irreducible sofic shift
  - **Generalized group mapping monoids** with an aperiodic 0-minimal ideal.
- A semigroup  $S$  is **generalized group mapping** if it has a (necessarily unique and regular) 0-minimal ideal  $I$  on which it acts faithfully on both the left and right.
- The so-called **Fischer cover** of an irreducible sofic shift is just the **Schützenberger graph** of the aperiodic 0-minimal ideal.
- Generalized group mapping semigroups were introduced by Rhodes in his work on the complexity of finite semigroups.
- The theory of lifting regular  $\mathcal{J}$ -classes, again due to Rhodes, plays a salient role in the proof of our theorem.

# Open questions

- Which projective profinite groups can be maximal subgroups of a free profinite monoid (Zaleskii)?
- Are there any trivial maximal subgroups of  $\widehat{A}^*$  other than the group of units?
- Can a free pro- $p$  group be a maximal subgroup of  $\widehat{A}^*$  (Zaleskii)?
- What makes the profinite group associated to a subshift finitely generated?
- What are the possible finite subsemigroups of a free profinite monoid?
  - We know that each element of such a semigroup is idempotent.
  - It can be shown that the principal ideals must form a chain using that ideals generated by idempotents are prime.



# Open questions

- Which projective profinite groups can be maximal subgroups of a free profinite monoid (Zaleskii)?
- Are there any trivial maximal subgroups of  $\widehat{A}^*$  other than the group of units?
- Can a free pro- $p$  group be a maximal subgroup of  $\widehat{A}^*$  (Zaleskii)?
- What makes the profinite group associated to a subshift finitely generated?
- What are the possible finite subsemigroups of a free profinite monoid?
  - We know that each element of such a semigroup is idempotent.
  - It can be shown that the principal ideals must form a chain using that ideals generated by idempotents are prime.

# Open questions

- Which projective profinite groups can be maximal subgroups of a free profinite monoid (Zaleskii)?
- Are there any trivial maximal subgroups of  $\widehat{A}^*$  other than the group of units?
- Can a free pro- $p$  group be a maximal subgroup of  $\widehat{A}^*$  (Zaleskii)?
- What makes the profinite group associated to a subshift finitely generated?
- What are the possible finite subsemigroups of a free profinite monoid?
  - We know that each element of such a semigroup is idempotent.
  - It can be shown that the principal ideals must form a chain using that ideals generated by idempotents are prime.

# Open questions

- Which projective profinite groups can be maximal subgroups of a free profinite monoid (Zaleskii)?
- Are there any trivial maximal subgroups of  $\widehat{A}^*$  other than the group of units?
- Can a free pro- $p$  group be a maximal subgroup of  $\widehat{A}^*$  (Zaleskii)?
- What makes the profinite group associated to a subshift finitely generated?
- What are the possible finite subsemigroups of a free profinite monoid?
  - We know that each element of such a semigroup is idempotent.
  - It can be shown that the principal ideals must form a chain using that ideals generated by idempotents are prime.

# Open questions

- Which projective profinite groups can be maximal subgroups of a free profinite monoid (Zaleskii)?
- Are there any trivial maximal subgroups of  $\widehat{A}^*$  other than the group of units?
- Can a free pro- $p$  group be a maximal subgroup of  $\widehat{A}^*$  (Zaleskii)?
- What makes the profinite group associated to a subshift finitely generated?
- What are the possible finite subsemigroups of a free profinite monoid?
  - We know that each element of such a semigroup is idempotent.
  - It can be shown that the principal ideals must form a chain using that ideals generated by idempotents are prime.

# Open questions

- Which projective profinite groups can be maximal subgroups of a free profinite monoid (Zaleskii)?
- Are there any trivial maximal subgroups of  $\widehat{A}^*$  other than the group of units?
- Can a free pro- $p$  group be a maximal subgroup of  $\widehat{A}^*$  (Zaleskii)?
- What makes the profinite group associated to a subshift finitely generated?
- What are the possible finite subsemigroups of a free profinite monoid?
  - We know that each element of such a semigroup is idempotent.
  - It can be shown that the principal ideals must form a chain using that ideals generated by idempotents are prime.

# Open questions

- Which projective profinite groups can be maximal subgroups of a free profinite monoid (Zaleskii)?
- Are there any trivial maximal subgroups of  $\widehat{A}^*$  other than the group of units?
- Can a free pro- $p$  group be a maximal subgroup of  $\widehat{A}^*$  (Zaleskii)?
- What makes the profinite group associated to a subshift finitely generated?
- What are the possible finite subsemigroups of a free profinite monoid?
  - We know that each element of such a semigroup is idempotent.
  - It can be shown that the principal ideals must form a chain using that ideals generated by idempotents are prime.

The end

THANK YOU FOR YOUR  
ATTENTION!