

Étale groupoids and inverse semigroups

David Milan¹ Benjamin Steinberg²

¹University of Texas at Tyler, ²Carleton University

bsteinbg@math.carleton.ca

<http://www.mathstat.carleton.ca/~bsteinbg>

July 14, 2010

The Exel paradigm

- Ruy Exel suggested the following paradigm for the study of “combinatorial C^* -algebras.”
 - Start with a combinatorial object (e.g., a graph or a tiling).
 - Construct an inverse semigroup.
 - From the semigroup construct an étale groupoid.
 - From the groupoid construct a C^* -algebra.
- One can hope to use the algebra to understand the combinatorial object (as in the case of tilings);
- Or use the combinatorial object to understand the algebra (as in the case of graphs).

The Exel paradigm

- Ruy Exel suggested the following paradigm for the study of “combinatorial C^* -algebras.”
 - Start with a combinatorial object (e.g., a graph or a tiling).
 - Construct an inverse semigroup.
 - From the semigroup construct an étale groupoid.
 - From the groupoid construct a C^* -algebra.
- One can hope to use the algebra to understand the combinatorial object (as in the case of tilings);
- Or use the combinatorial object to understand the algebra (as in the case of graphs).

The Exel paradigm

- Ruy Exel suggested the following paradigm for the study of “combinatorial C^* -algebras.”
 - Start with a combinatorial object (e.g., a graph or a tiling).
 - Construct an inverse semigroup.
 - From the semigroup construct an étale groupoid.
 - From the groupoid construct a C^* -algebra.
- One can hope to use the algebra to understand the combinatorial object (as in the case of tilings);
- Or use the combinatorial object to understand the algebra (as in the case of graphs).

The Exel paradigm

- Ruy Exel suggested the following paradigm for the study of “combinatorial C^* -algebras.”
 - Start with a combinatorial object (e.g., a graph or a tiling).
 - Construct an inverse semigroup.
 - From the semigroup construct an étale groupoid.
 - From the groupoid construct a C^* -algebra.
- One can hope to use the algebra to understand the combinatorial object (as in the case of tilings);
- Or use the combinatorial object to understand the algebra (as in the case of graphs).

The Exel paradigm

- Ruy Exel suggested the following paradigm for the study of “combinatorial C^* -algebras.”
 - Start with a combinatorial object (e.g., a graph or a tiling).
 - Construct an inverse semigroup.
 - From the semigroup construct an étale groupoid.
 - From the groupoid construct a C^* -algebra.
- One can hope to use the algebra to understand the combinatorial object (as in the case of tilings);
- Or use the combinatorial object to understand the algebra (as in the case of graphs).

The Exel paradigm

- Ruy Exel suggested the following paradigm for the study of “combinatorial C^* -algebras.”
 - Start with a combinatorial object (e.g., a graph or a tiling).
 - Construct an inverse semigroup.
 - From the semigroup construct an étale groupoid.
 - From the groupoid construct a C^* -algebra.
- One can hope to use the algebra to understand the combinatorial object (as in the case of tilings);
- Or use the combinatorial object to understand the algebra (as in the case of graphs).

The Exel paradigm

- Ruy Exel suggested the following paradigm for the study of “combinatorial C^* -algebras.”
 - Start with a combinatorial object (e.g., a graph or a tiling).
 - Construct an inverse semigroup.
 - From the semigroup construct an étale groupoid.
 - From the groupoid construct a C^* -algebra.
- One can hope to use the algebra to understand the combinatorial object (as in the case of tilings);
- Or use the combinatorial object to understand the algebra (as in the case of graphs).

- It is by now well known that natural notions for étale groupoids translate into natural notions for C^* -algebras.
- For instance, Morita equivalent groupoids have strongly Morita equivalent algebras.
- The (partial action) semidirect product of a group with a space yields a (partial action) cross product of the group with a commutative C^* -algebra.
- The algebra of a groupoid of germs of an inverse semigroup action on a space translates to an inverse semigroup cross product algebra.

- It is by now well known that natural notions for étale groupoids translate into natural notions for C^* -algebras.
- For instance, **Morita equivalent** groupoids have **strongly Morita equivalent** algebras.
- The (partial action) **semidirect product** of a group with a space yields a (partial action) **cross product** of the group with a commutative C^* -algebra.
- The algebra of a **groupoid of germs** of an inverse semigroup action on a space translates to an inverse semigroup **cross product** algebra.

- It is by now well known that natural notions for étale groupoids translate into natural notions for C^* -algebras.
- For instance, Morita equivalent groupoids have strongly Morita equivalent algebras.
- The (partial action) semidirect product of a group with a space yields a (partial action) cross product of the group with a commutative C^* -algebra.
- The algebra of a groupoid of germs of an inverse semigroup action on a space translates to an inverse semigroup cross product algebra.

- It is by now well known that natural notions for étale groupoids translate into natural notions for C^* -algebras.
- For instance, **Morita equivalent** groupoids have **strongly Morita equivalent** algebras.
- The (partial action) **semidirect product** of a group with a space yields a (partial action) **cross product** of the group with a commutative C^* -algebra.
- The algebra of a **groupoid of germs** of an inverse semigroup action on a space translates to an inverse semigroup **cross product** algebra.

The dictionary II

- It seems to be less well known what happens when passing from an inverse semigroup to a groupoid.
- Does Morita equivalence pass through to the groupoids?
- Do (partial action) semidirect products become (partial action) semidirect products?
- Functoriality seems not to have been addressed.
- It is also not universally agreed upon which groupoid to assign to an inverse semigroup.
- There are Paterson's universal groupoid, Kellendonk's groupoid and Exel's tight groupoid.
- Here we will focus on Paterson's groupoid, as the others are reductions. All of our results so far are valid for any of these groupoids.

The dictionary II

- It seems to be less well known what happens when passing from an inverse semigroup to a groupoid.
- Does Morita equivalence pass through to the groupoids?
- Do (partial action) semidirect products become (partial action) semidirect products?
- Functoriality seems not to have been addressed.
- It is also not universally agreed upon which groupoid to assign to an inverse semigroup.
- There are Paterson's universal groupoid, Kellendonk's groupoid and Exel's tight groupoid.
- Here we will focus on Paterson's groupoid, as the others are reductions. All of our results so far are valid for any of these groupoids.

The dictionary II

- It seems to be less well known what happens when passing from an inverse semigroup to a groupoid.
- Does Morita equivalence pass through to the groupoids?
- Do (partial action) semidirect products become (partial action) semidirect products?
- Functoriality seems not to have been addressed.
- It is also not universally agreed upon which groupoid to assign to an inverse semigroup.
- There are Paterson's universal groupoid, Kellendonk's groupoid and Exel's tight groupoid.
- Here we will focus on Paterson's groupoid, as the others are reductions. All of our results so far are valid for any of these groupoids.

The dictionary II

- It seems to be less well known what happens when passing from an inverse semigroup to a groupoid.
- Does Morita equivalence pass through to the groupoids?
- Do (partial action) semidirect products become (partial action) semidirect products?
- Functoriality seems not to have been addressed.
- It is also not universally agreed upon which groupoid to assign to an inverse semigroup.
- There are Paterson's universal groupoid, Kellendonk's groupoid and Exel's tight groupoid.
- Here we will focus on Paterson's groupoid, as the others are reductions. All of our results so far are valid for any of these groupoids.

The dictionary II

- It seems to be less well known what happens when passing from an inverse semigroup to a groupoid.
- Does Morita equivalence pass through to the groupoids?
- Do (partial action) semidirect products become (partial action) semidirect products?
- Functoriality seems not to have been addressed.
- It is also not universally agreed upon which groupoid to assign to an inverse semigroup.
- There are Paterson's universal groupoid, Kellendonk's groupoid and Exel's tight groupoid.
- Here we will focus on Paterson's groupoid, as the others are reductions. All of our results so far are valid for any of these groupoids.

The dictionary II

- It seems to be less well known what happens when passing from an inverse semigroup to a groupoid.
- Does Morita equivalence pass through to the groupoids?
- Do (partial action) semidirect products become (partial action) semidirect products?
- Functoriality seems not to have been addressed.
- It is also not universally agreed upon which groupoid to assign to an inverse semigroup.
- There are Paterson's universal groupoid, Kellendonk's groupoid and Exel's tight groupoid.
- Here we will focus on Paterson's groupoid, as the others are reductions. All of our results so far are valid for any of these groupoids.

- It seems to be less well known what happens when passing from an inverse semigroup to a groupoid.
- Does Morita equivalence pass through to the groupoids?
- Do (partial action) semidirect products become (partial action) semidirect products?
- Functoriality seems not to have been addressed.
- It is also not universally agreed upon which groupoid to assign to an inverse semigroup.
- There are Paterson's universal groupoid, Kellendonk's groupoid and Exel's tight groupoid.
- Here we will focus on Paterson's groupoid, as the others are reductions. All of our results so far are valid for any of these groupoids.

A primer on inverse semigroups

- Inverse semigroups are the abstract counterparts of pseudogroups of partial homeomorphisms of a space.
- If X is a topological space, then I_X is the pseudogroup of all homeomorphisms between open subsets of X .
- Composition is defined where it makes sense: if $f: U \rightarrow V$ and $g: U' \rightarrow V'$, then

$$f \circ g: g^{-1}(U \cap V') \rightarrow f(U \cap V').$$

- Inverse semigroups can also be viewed as $*$ -semigroups of partial isometries of a Hilbert space.

A primer on inverse semigroups

- Inverse semigroups are the abstract counterparts of pseudogroups of partial homeomorphisms of a space.
- If X is a topological space, then I_X is the pseudogroup of all homeomorphisms between open subsets of X .
- Composition is defined where it makes sense: if $f: U \rightarrow V$ and $g: U' \rightarrow V'$, then

$$f \circ g: g^{-1}(U \cap V') \rightarrow f(U \cap V').$$

- Inverse semigroups can also be viewed as $*$ -semigroups of partial isometries of a Hilbert space.

A primer on inverse semigroups

- Inverse semigroups are the abstract counterparts of pseudogroups of partial homeomorphisms of a space.
- If X is a topological space, then I_X is the pseudogroup of all homeomorphisms between open subsets of X .
- Composition is defined where it makes sense: if $f: U \rightarrowtail V$ and $g: U' \rightarrowtail V'$, then

$$f \circ g: g^{-1}(U \cap V') \rightarrowtail f(U \cap V').$$

- Inverse semigroups can also be viewed as $*$ -semigroups of partial isometries of a Hilbert space.

A primer on inverse semigroups

- Inverse semigroups are the abstract counterparts of pseudogroups of partial homeomorphisms of a space.
- If X is a topological space, then I_X is the pseudogroup of all homeomorphisms between open subsets of X .
- Composition is defined where it makes sense: if $f: U \rightarrowtail V$ and $g: U' \rightarrowtail V'$, then

$$f \circ g: g^{-1}(U \cap V') \rightarrowtail f(U \cap V').$$

- Inverse semigroups can also be viewed as $*$ -semigroups of partial isometries of a Hilbert space.

More on inverse semigroups

- Formally, an **inverse semigroup** is a semigroup S such that for all $s \in S$, there is a unique element s^* such that

$$ss^*s = s, \quad s^*ss^* = s^*$$

- .
- Munn and Penrose proved that the idempotents of S commute and hence $E(S)$ is a subsemigroup.
- $E(S)$ is ordered by $e \leq f$ iff $ef = e$ and then $ef = e \wedge f$.
- So $E(S)$ is a **meet semilattice**.
- The order extends to S by putting $s \leq t$ if $s = te$ for some idempotent e .
- For I_X , the ordering is restriction.

More on inverse semigroups

- Formally, an **inverse semigroup** is a semigroup S such that for all $s \in S$, there is a unique element s^* such that

$$ss^*s = s, \quad s^*ss^* = s^*$$

- Munn and Penrose proved that the idempotents of S commute and hence $E(S)$ is a subsemigroup.
- $E(S)$ is ordered by $e \leq f$ iff $ef = e$ and then $ef = e \wedge f$.
- So $E(S)$ is a **meet semilattice**.
- The order extends to S by putting $s \leq t$ if $s = te$ for some idempotent e .
- For I_X , the ordering is restriction.

More on inverse semigroups

- Formally, an **inverse semigroup** is a semigroup S such that for all $s \in S$, there is a unique element s^* such that

$$ss^*s = s, \quad s^*ss^* = s^*$$

- .
- Munn and Penrose proved that the idempotents of S commute and hence $E(S)$ is a subsemigroup.
- $E(S)$ is ordered by $e \leq f$ iff $ef = e$ and then $ef = e \wedge f$.
- So $E(S)$ is a **meet semilattice**.
- The order extends to S by putting $s \leq t$ if $s = te$ for some idempotent e .
- For I_X , the ordering is restriction.

More on inverse semigroups

- Formally, an **inverse semigroup** is a semigroup S such that for all $s \in S$, there is a unique element s^* such that

$$ss^*s = s, \quad s^*ss^* = s^*$$

- .
- Munn and Penrose proved that the idempotents of S commute and hence $E(S)$ is a subsemigroup.
- $E(S)$ is ordered by $e \leq f$ iff $ef = e$ and then $ef = e \wedge f$.
- So $E(S)$ is a **meet semilattice**.
- The order extends to S by putting $s \leq t$ if $s = te$ for some idempotent e .
- For I_X , the ordering is restriction.

More on inverse semigroups

- Formally, an **inverse semigroup** is a semigroup S such that for all $s \in S$, there is a unique element s^* such that

$$ss^*s = s, \quad s^*ss^* = s^*$$

- .
- Munn and Penrose proved that the idempotents of S commute and hence $E(S)$ is a subsemigroup.
- $E(S)$ is ordered by $e \leq f$ iff $ef = e$ and then $ef = e \wedge f$.
- So $E(S)$ is a **meet semilattice**.
- The order extends to S by putting $s \leq t$ if $s = te$ for some idempotent e .
- For I_X , the ordering is restriction.

More on inverse semigroups

- Formally, an **inverse semigroup** is a semigroup S such that for all $s \in S$, there is a unique element s^* such that

$$ss^*s = s, \quad s^*ss^* = s^*$$

.

- Munn and Penrose proved that the idempotents of S commute and hence $E(S)$ is a subsemigroup.
- $E(S)$ is ordered by $e \leq f$ iff $ef = e$ and then $ef = e \wedge f$.
- So $E(S)$ is a **meet semilattice**.
- The order extends to S by putting $s \leq t$ if $s = te$ for some idempotent e .
- For I_X , the ordering is restriction.

The maximal group image

- Identifying two elements of S with a common lower bound (i.e., the same germ) yields a congruence.
- The quotient is a group $G(S)$ called the **maximal group image**.
- The canonical projection $\sigma: S \twoheadrightarrow G(S)$ is universal to groups.
- Clearly $E(S) \subseteq \sigma^{-1}(1)$.
- S is called E -unitary if $E(S) = \sigma^{-1}(1)$.
- S is called F -inverse if $eSf \cap \sigma^{-1}(g)$ has a maximum element for all $e, f \in E(S)$ and $g \in G(S)$.
- F -inverse semigroups are E -unitary.

The maximal group image

- Identifying two elements of S with a common lower bound (i.e., the same germ) yields a congruence.
- The quotient is a group $G(S)$ called the **maximal group image**.
- The canonical projection $\sigma: S \twoheadrightarrow G(S)$ is universal to groups.
- Clearly $E(S) \subseteq \sigma^{-1}(1)$.
- S is called **E -unitary** if $E(S) = \sigma^{-1}(1)$.
- S is called **F -inverse** if $eSf \cap \sigma^{-1}(g)$ has a maximum element for all $e, f \in E(S)$ and $g \in G(S)$.
- **F -inverse semigroups are E -unitary.**

The maximal group image

- Identifying two elements of S with a common lower bound (i.e., the same germ) yields a congruence.
- The quotient is a group $G(S)$ called the **maximal group image**.
- The canonical projection $\sigma: S \twoheadrightarrow G(S)$ is universal to groups.
- Clearly $E(S) \subseteq \sigma^{-1}(1)$.
- S is called **E -unitary** if $E(S) = \sigma^{-1}(1)$.
- S is called **F -inverse** if $eSf \cap \sigma^{-1}(g)$ has a maximum element for all $e, f \in E(S)$ and $g \in G(S)$.
- F -inverse semigroups are E -unitary.

The maximal group image

- Identifying two elements of S with a common lower bound (i.e., the same germ) yields a congruence.
- The quotient is a group $G(S)$ called the **maximal group image**.
- The canonical projection $\sigma: S \twoheadrightarrow G(S)$ is universal to groups.
- Clearly $E(S) \subseteq \sigma^{-1}(1)$.
- S is called **E -unitary** if $E(S) = \sigma^{-1}(1)$.
- S is called **F -inverse** if $eSf \cap \sigma^{-1}(g)$ has a maximum element for all $e, f \in E(S)$ and $g \in G(S)$.
- F -inverse semigroups are E -unitary.

The maximal group image

- Identifying two elements of S with a common lower bound (i.e., the same germ) yields a congruence.
- The quotient is a group $G(S)$ called the **maximal group image**.
- The canonical projection $\sigma: S \twoheadrightarrow G(S)$ is universal to groups.
- Clearly $E(S) \subseteq \sigma^{-1}(1)$.
- S is called **E -unitary** if $E(S) = \sigma^{-1}(1)$.
- S is called **F -inverse** if $eSf \cap \sigma^{-1}(g)$ has a maximum element for all $e, f \in E(S)$ and $g \in G(S)$.
- F -inverse semigroups are E -unitary.

The maximal group image

- Identifying two elements of S with a common lower bound (i.e., the same germ) yields a congruence.
- The quotient is a group $G(S)$ called the **maximal group image**.
- The canonical projection $\sigma: S \twoheadrightarrow G(S)$ is universal to groups.
- Clearly $E(S) \subseteq \sigma^{-1}(1)$.
- S is called **E -unitary** if $E(S) = \sigma^{-1}(1)$.
- S is called **F -inverse** if $eSf \cap \sigma^{-1}(g)$ has a maximum element for all $e, f \in E(S)$ and $g \in G(S)$.
- F -inverse semigroups are E -unitary.

The maximal group image

- Identifying two elements of S with a common lower bound (i.e., the same germ) yields a congruence.
- The quotient is a group $G(S)$ called the **maximal group image**.
- The canonical projection $\sigma: S \twoheadrightarrow G(S)$ is universal to groups.
- Clearly $E(S) \subseteq \sigma^{-1}(1)$.
- S is called **E -unitary** if $E(S) = \sigma^{-1}(1)$.
- S is called **F -inverse** if $eSf \cap \sigma^{-1}(g)$ has a maximum element for all $e, f \in E(S)$ and $g \in G(S)$.
- F -inverse semigroups are E -unitary.

- If E is a meet semilattice, then E is an inverse semigroup of “projections.”
- Meet semilattices are to inverse semigroups as spaces are to étale groupoids.
- A group is an inverse semigroup.
- If a group G acts on a semilattice E , the semidirect product $E \rtimes G$ is an F -inverse semigroup.
- If G acts partially on E , then $E \rtimes G$ is E -unitary and by a theorem of McAlister all E -unitary inverse semigroups arise in this way.

- If E is a meet semilattice, then E is an inverse semigroup of “projections.”
- Meet semilattices are to inverse semigroups as spaces are to étale groupoids.
- A group is an inverse semigroup.
- If a group G acts on a semilattice E , the semidirect product $E \rtimes G$ is an F -inverse semigroup.
- If G acts partially on E , then $E \rtimes G$ is E -unitary and by a theorem of McAlister all E -unitary inverse semigroups arise in this way.

Extreme examples

- If E is a meet semilattice, then E is an inverse semigroup of “projections.”
- Meet semilattices are to inverse semigroups as spaces are to étale groupoids.
- A group is an inverse semigroup.
- If a group G acts on a semilattice E , the semidirect product $E \rtimes G$ is an F -inverse semigroup.
- If G acts partially on E , then $E \rtimes G$ is E -unitary and by a theorem of McAlister all E -unitary inverse semigroups arise in this way.

- If E is a meet semilattice, then E is an inverse semigroup of “projections.”
- Meet semilattices are to inverse semigroups as spaces are to étale groupoids.
- A group is an inverse semigroup.
- If a group G acts on a semilattice E , the semidirect product $E \rtimes G$ is an F -inverse semigroup.
- If G acts partially on E , then $E \rtimes G$ is E -unitary and by a theorem of McAlister all E -unitary inverse semigroups arise in this way.

- If E is a meet semilattice, then E is an inverse semigroup of “projections.”
- Meet semilattices are to inverse semigroups as spaces are to étale groupoids.
- A group is an inverse semigroup.
- If a group G acts on a semilattice E , the semidirect product $E \rtimes G$ is an F -inverse semigroup.
- If G acts partially on E , then $E \rtimes G$ is E -unitary and by a theorem of McAlister all E -unitary inverse semigroups arise in this way.

More interesting examples

- The **bicyclic** monoid is the inverse semigroup B generated by a unilateral shift and its adjoint.
- Its semilattice of idempotents is (\mathbb{N}, \geq) .
- It is E -unitary with maximal group image is \mathbb{Z} (given by the Fredholm index).
- The **polycyclic** (or Cuntz) inverse semigroup is given by the presentation $P_X = \langle X \mid x^*y = \delta_{x,y} \rangle$ (as an inverse semigroup with 0).
- It is a (congruence) simple inverse semigroup and can be identified with the inverse semigroup generated by the generators of the Cuntz algebra.
- Similarly, there is a graph inverse semigroup associated to any directed graph which is analogous to the graph C^* -algebra.

More interesting examples

- The **bicyclic** monoid is the inverse semigroup B generated by a unilateral shift and its adjoint.
- Its semilattice of idempotents is (\mathbb{N}, \geq) .
- It is E -unitary with maximal group image is \mathbb{Z} (given by the Fredholm index).
- The **polycyclic** (or Cuntz) inverse semigroup is given by the presentation $P_X = \langle X \mid x^*y = \delta_{x,y} \rangle$ (as an inverse semigroup with 0).
- It is a (congruence) simple inverse semigroup and can be identified with the inverse semigroup generated by the generators of the Cuntz algebra.
- Similarly, there is a graph inverse semigroup associated to any directed graph which is analogous to the graph C^* -algebra.

More interesting examples

- The **bicyclic** monoid is the inverse semigroup B generated by a unilateral shift and its adjoint.
- Its semilattice of idempotents is (\mathbb{N}, \geq) .
- It is E -unitary with maximal group image is \mathbb{Z} (given by the Fredholm index).
- The **polycyclic** (or Cuntz) inverse semigroup is given by the presentation $P_X = \langle X \mid x^*y = \delta_{x,y} \rangle$ (as an inverse semigroup with 0).
- It is a (congruence) simple inverse semigroup and can be identified with the inverse semigroup generated by the generators of the Cuntz algebra.
- Similarly, there is a **graph inverse semigroup** associated to any directed graph which is analogous to the graph C^* -algebra.

More interesting examples

- The **bicyclic** monoid is the inverse semigroup B generated by a unilateral shift and its adjoint.
- Its semilattice of idempotents is (\mathbb{N}, \geq) .
- It is E -unitary with maximal group image is \mathbb{Z} (given by the Fredholm index).
- The **polycyclic** (or Cuntz) inverse semigroup is given by the presentation $P_X = \langle X \mid x^*y = \delta_{x,y} \rangle$ (as an inverse semigroup with 0).
- It is a (congruence) simple inverse semigroup and can be identified with the inverse semigroup generated by the generators of the Cuntz algebra.
- Similarly, there is a **graph inverse semigroup** associated to any directed graph which is analogous to the graph C^* -algebra.

More interesting examples

- The **bicyclic** monoid is the inverse semigroup B generated by a unilateral shift and its adjoint.
- Its semilattice of idempotents is (\mathbb{N}, \geq) .
- It is E -unitary with maximal group image is \mathbb{Z} (given by the Fredholm index).
- The **polycyclic** (or Cuntz) inverse semigroup is given by the presentation $P_X = \langle X \mid x^*y = \delta_{x,y} \rangle$ (as an inverse semigroup with 0).
- It is a (congruence) simple inverse semigroup and can be identified with the inverse semigroup generated by the generators of the Cuntz algebra.
- Similarly, there is a **graph inverse semigroup** associated to any directed graph which is analogous to the graph C^* -algebra.

More interesting examples

- The **bicyclic** monoid is the inverse semigroup B generated by a unilateral shift and its adjoint.
- Its semilattice of idempotents is (\mathbb{N}, \geq) .
- It is E -unitary with maximal group image is \mathbb{Z} (given by the Fredholm index).
- The **polycyclic** (or Cuntz) inverse semigroup is given by the presentation $P_X = \langle X \mid x^*y = \delta_{x,y} \rangle$ (as an inverse semigroup with 0).
- It is a (congruence) simple inverse semigroup and can be identified with the inverse semigroup generated by the generators of the Cuntz algebra.
- Similarly, there is a **graph inverse semigroup** associated to any directed graph which is analogous to the graph C^* -algebra.

The Meakin-Margolis-Munn construction

- Let G be a group with generators X and corresponding Cayley graph Γ .
- $M_A(G)$ consists of all equivalence classes of triples $[g, \Lambda, h]$ where Λ is a finite connected subgraph of Γ and g, h are vertices of Λ .
- Equivalence is up to translation by G .
- The product is given by

$$[g, \Lambda, h][g', \Lambda', h'] = [g, \Lambda \cup h\Lambda', hh'].$$

- $M_A(G)$ is E -unitary with maximal group image G .
- If G is free on A , then $M_A(G)$ is a free inverse monoid on A .
- Free inverse monoids classify graph immersions over a bouquet.

The Meakin-Margolis-Munn construction

- Let G be a group with generators X and corresponding Cayley graph Γ .
- $M_A(G)$ consists of all equivalence classes of triples $[g, \Lambda, h]$ where Λ is a finite connected subgraph of Γ and g, h are vertices of Λ .
- Equivalence is up to translation by G .
- The product is given by

$$[g, \Lambda, h][g', \Lambda', h'] = [g, \Lambda \cup h\Lambda', hh'].$$

- $M_A(G)$ is E -unitary with maximal group image G .
- If G is free on A , then $M_A(G)$ is a free inverse monoid on A .
- Free inverse monoids classify graph immersions over a bouquet.

The Meakin-Margolis-Munn construction

- Let G be a group with generators X and corresponding Cayley graph Γ .
- $M_A(G)$ consists of all equivalence classes of triples $[g, \Lambda, h]$ where Λ is a finite connected subgraph of Γ and g, h are vertices of Λ .
- Equivalence is up to translation by G .
- The product is given by

$$[g, \Lambda, h][g', \Lambda', h'] = [g, \Lambda \cup h\Lambda', hh'].$$

- $M_A(G)$ is E -unitary with maximal group image G .
- If G is free on A , then $M_A(G)$ is a free inverse monoid on A .
- Free inverse monoids classify graph immersions over a bouquet.

The Meakin-Margolis-Munn construction

- Let G be a group with generators X and corresponding Cayley graph Γ .
- $M_A(G)$ consists of all equivalence classes of triples $[g, \Lambda, h]$ where Λ is a finite connected subgraph of Γ and g, h are vertices of Λ .
- Equivalence is up to translation by G .
- The product is given by

$$[g, \Lambda, h][g', \Lambda', h'] = [g, \Lambda \cup h\Lambda', hh'].$$

- $M_A(G)$ is E -unitary with maximal group image G .
- If G is free on A , then $M_A(G)$ is a free inverse monoid on A .
- Free inverse monoids classify graph immersions over a bouquet.

The Meakin-Margolis-Munn construction

- Let G be a group with generators X and corresponding Cayley graph Γ .
- $M_A(G)$ consists of all equivalence classes of triples $[g, \Lambda, h]$ where Λ is a finite connected subgraph of Γ and g, h are vertices of Λ .
- Equivalence is up to translation by G .
- The product is given by

$$[g, \Lambda, h][g', \Lambda', h'] = [g, \Lambda \cup h\Lambda', hh'].$$

- $M_A(G)$ is E -unitary with maximal group image G .
- If G is free on A , then $M_A(G)$ is a free inverse monoid on A .
- Free inverse monoids classify graph immersions over a bouquet.

The Meakin-Margolis-Munn construction

- Let G be a group with generators X and corresponding Cayley graph Γ .
- $M_A(G)$ consists of all equivalence classes of triples $[g, \Lambda, h]$ where Λ is a finite connected subgraph of Γ and g, h are vertices of Λ .
- Equivalence is up to translation by G .
- The product is given by

$$[g, \Lambda, h][g', \Lambda', h'] = [g, \Lambda \cup h\Lambda', hh'].$$

- $M_A(G)$ is E -unitary with maximal group image G .
- If G is free on A , then $M_A(G)$ is a free inverse monoid on A .
- Free inverse monoids classify graph immersions over a bouquet.

The Meakin-Margolis-Munn construction

- Let G be a group with generators X and corresponding Cayley graph Γ .
- $M_A(G)$ consists of all equivalence classes of triples $[g, \Lambda, h]$ where Λ is a finite connected subgraph of Γ and g, h are vertices of Λ .
- Equivalence is up to translation by G .
- The product is given by

$$[g, \Lambda, h][g', \Lambda', h'] = [g, \Lambda \cup h\Lambda', hh'].$$

- $M_A(G)$ is E -unitary with maximal group image G .
- If G is free on A , then $M_A(G)$ is a free inverse monoid on A .
- Free inverse monoids classify graph immersions over a bouquet.

- Let \mathcal{T} be a **tiling** of \mathbb{R}^d .
- Kellendonk's tiling semigroup has elements 0 and equivalence classes $[t, P, u]$ where P is a pattern and t, u are tiles of P .
- Equivalence is up to translation in \mathbb{R}^d .
- The product $[t, P, u][t', P', u']$ is non-zero iff there are $v, v' \in \mathbb{R}^d$ such that $u + v = t' + v'$ and $(P + v) \cup (P' + v')$ is a pattern.
- In this case, the product is $[t + v, (P + v) \cup (P' + v'), u' + v']$.

- Let \mathcal{T} be a **tiling** of \mathbb{R}^d .
- Kellendonk's tiling semigroup has elements 0 and equivalence classes $[t, P, u]$ where P is a pattern and t, u are tiles of P .
- Equivalence is up to translation in \mathbb{R}^d .
- The product $[t, P, u][t', P', u']$ is non-zero iff there are $v, v' \in \mathbb{R}^d$ such that $u + v = t' + v'$ and $(P + v) \cup (P' + v')$ is a pattern.
- In this case, the product is $[t + v, (P + v) \cup (P' + v'), u' + v']$.

Tiling semigroups

- Let \mathcal{T} be a **tiling** of \mathbb{R}^d .
- Kellendonk's tiling semigroup has elements 0 and equivalence classes $[t, P, u]$ where P is a pattern and t, u are tiles of P .
- Equivalence is up to translation in \mathbb{R}^d .
- The product $[t, P, u][t', P', u']$ is non-zero iff there are $v, v' \in \mathbb{R}^d$ such that $u + v = t' + v'$ and $(P + v) \cup (P' + v')$ is a pattern.
- In this case, the product is $[t + v, (P + v) \cup (P' + v'), u' + v']$.

- Let \mathcal{T} be a **tiling** of \mathbb{R}^d .
- Kellendonk's tiling semigroup has elements 0 and equivalence classes $[t, P, u]$ where P is a pattern and t, u are tiles of P .
- Equivalence is up to translation in \mathbb{R}^d .
- The product $[t, P, u][t', P', u']$ is non-zero iff there are $v, v' \in \mathbb{R}^d$ such that $u + v = t' + v'$ and $(P + v) \cup (P' + v')$ is a pattern.
- In this case, the product is $[t + v, (P + v) \cup (P' + v'), u' + v']$.

- Let \mathcal{T} be a **tiling** of \mathbb{R}^d .
- Kellendonk's tiling semigroup has elements 0 and equivalence classes $[t, P, u]$ where P is a pattern and t, u are tiles of P .
- Equivalence is up to translation in \mathbb{R}^d .
- The product $[t, P, u][t', P', u']$ is non-zero iff there are $v, v' \in \mathbb{R}^d$ such that $u + v = t' + v'$ and $(P + v) \cup (P' + v')$ is a pattern.
- In this case, the product is $[t + v, (P + v) \cup (P' + v'), u' + v']$.

Some F -inverse monoids

- Let X be an irreducible algebraic variety and S_X the inverse semigroup of isomorphisms between open subvarieties.
- Then S_X is an F -inverse monoid and the maximal group image is the group of **birational automorphisms** of X .
- When $X = \mathbb{P}^1$, then S_X is the Möbius inverse semigroup and $G(S_X)$ is the group of **linear fractional transformations**.
- Let $\{a, b\}^*$ be a free monoid on $\{a, b\}$ and S the inverse semigroup of isomorphisms between finitely generated essential right ideals of $\{a, b\}^*$.
- Then S is an F -inverse monoid with maximal group image Thompson's simple group V .
- If G is a group and S is the inverse semigroup of virtual automorphisms of G , then S is F -inverse and the maximal group image is the abstract commensurator of G .

Some F -inverse monoids

- Let X be an irreducible algebraic variety and S_X the inverse semigroup of isomorphisms between open subvarieties.
- Then S_X is an F -inverse monoid and the maximal group image is the group of **birational automorphisms** of X .
- When $X = \mathbb{P}^1$, then S_X is the Möbius inverse semigroup and $G(S_X)$ is the group of **linear fractional transformations**.
- Let $\{a, b\}^*$ be a free monoid on $\{a, b\}$ and S the inverse semigroup of isomorphisms between finitely generated essential right ideals of $\{a, b\}^*$.
- Then S is an F -inverse monoid with maximal group image **Thompson's simple group** V .
- If G is a group and S is the inverse semigroup of virtual automorphisms of G , then S is F -inverse and the maximal group image is the abstract commensurator of G .

Some F -inverse monoids

- Let X be an irreducible algebraic variety and S_X the inverse semigroup of isomorphisms between open subvarieties.
- Then S_X is an F -inverse monoid and the maximal group image is the group of **birational automorphisms** of X .
- When $X = \mathbb{P}^1$, then S_X is the Möbius inverse semigroup and $G(S_X)$ is the group of **linear fractional transformations**.
- Let $\{a, b\}^*$ be a free monoid on $\{a, b\}$ and S the inverse semigroup of isomorphisms between finitely generated essential right ideals of $\{a, b\}^*$.
- Then S is an F -inverse monoid with maximal group image **Thompson's simple group** V .
- If G is a group and S is the inverse semigroup of virtual automorphisms of G , then S is F -inverse and the maximal group image is the **abstract commensurator** of G .

Some F -inverse monoids

- Let X be an irreducible algebraic variety and S_X the inverse semigroup of isomorphisms between open subvarieties.
- Then S_X is an F -inverse monoid and the maximal group image is the group of **birational automorphisms** of X .
- When $X = \mathbb{P}^1$, then S_X is the Möbius inverse semigroup and $G(S_X)$ is the group of **linear fractional transformations**.
- Let $\{a, b\}^*$ be a free monoid on $\{a, b\}$ and S the inverse semigroup of isomorphisms between finitely generated essential right ideals of $\{a, b\}^*$.
- Then S is an F -inverse monoid with maximal group image **Thompson's simple group** V .
- If G is a group and S is the inverse semigroup of virtual automorphisms of G , then S is F -inverse and the maximal group image is the **abstract commensurator** of G .

Some F -inverse monoids

- Let X be an irreducible algebraic variety and S_X the inverse semigroup of isomorphisms between open subvarieties.
- Then S_X is an F -inverse monoid and the maximal group image is the group of **birational automorphisms** of X .
- When $X = \mathbb{P}^1$, then S_X is the Möbius inverse semigroup and $G(S_X)$ is the group of **linear fractional transformations**.
- Let $\{a, b\}^*$ be a free monoid on $\{a, b\}$ and S the inverse semigroup of isomorphisms between finitely generated essential right ideals of $\{a, b\}^*$.
- Then S is an F -inverse monoid with maximal group image **Thompson's simple group** V .
- If G is a group and S is the inverse semigroup of virtual automorphisms of G , then S is F -inverse and the maximal group image is the **abstract commensurator** of G .

Some F -inverse monoids

- Let X be an irreducible algebraic variety and S_X the inverse semigroup of isomorphisms between open subvarieties.
- Then S_X is an F -inverse monoid and the maximal group image is the group of **birational automorphisms** of X .
- When $X = \mathbb{P}^1$, then S_X is the Möbius inverse semigroup and $G(S_X)$ is the group of **linear fractional transformations**.
- Let $\{a, b\}^*$ be a free monoid on $\{a, b\}$ and S the inverse semigroup of isomorphisms between finitely generated essential right ideals of $\{a, b\}^*$.
- Then S is an F -inverse monoid with maximal group image **Thompson's simple group** V .
- If G is a group and S is the inverse semigroup of virtual automorphisms of G , then S is F -inverse and the maximal group image is the **abstract commensurator** of G .

- Exel associated to every group G an F -inverse monoid $S(G)$ whose actions correspond to partial actions of G .
- The same semigroup was discovered earlier by Birget and Rhodes in a different context.
- Szendrei had shown $S(G)$ is the universal F -inverse cover of G .
- In some sense, F -inverse monoid covers of G and partial group actions of G are the same subject.

- Exel associated to every group G an F -inverse monoid $S(G)$ whose actions correspond to partial actions of G .
- The same semigroup was discovered earlier by Birget and Rhodes in a different context.
- Szendrei had shown $S(G)$ is the universal F -inverse cover of G .
- In some sense, F -inverse monoid covers of G and partial group actions of G are the same subject.

- Exel associated to every group G an F -inverse monoid $S(G)$ whose actions correspond to partial actions of G .
- The same semigroup was discovered earlier by Birget and Rhodes in a different context.
- Szendrei had shown $S(G)$ is the universal F -inverse cover of G .
- In some sense, F -inverse monoid covers of G and partial group actions of G are the same subject.

- Exel associated to every group G an F -inverse monoid $S(G)$ whose actions correspond to partial actions of G .
- The same semigroup was discovered earlier by Birget and Rhodes in a different context.
- Szendrei had shown $S(G)$ is the universal F -inverse cover of G .
- In some sense, F -inverse monoid covers of G and partial group actions of G are the same subject.

Strongly 0- E -unitary inverse semigroups

- An inverse semigroups with 0 cannot be E -unitary without being a semilattice.
- So one needs an appropriate variation of the concept for this context.
- If S is an inverse semigroup with 0, the **universal group** $U(S)$ is the group with generators $S \setminus \{0\}$ and relations given $s \cdot t = st$ if $st \neq 0$.
- The natural map $\sigma: S \setminus \{0\} \longrightarrow U(S)$ satisfies $\sigma(s)\sigma(t) = \sigma(st)$ whenever $st \neq 0$ and is universal for this property.
- S is called **strongly 0- E -unitary** if $\sigma^{-1}(1) = E(S) \setminus \{0\}$.

Strongly 0- E -unitary inverse semigroups

- An inverse semigroups with 0 cannot be E -unitary without being a semilattice.
- So one needs an appropriate variation of the concept for this context.
- If S is an inverse semigroup with 0, the **universal group** $U(S)$ is the group with generators $S \setminus \{0\}$ and relations given $s \cdot t = st$ if $st \neq 0$.
- The natural map $\sigma: S \setminus \{0\} \longrightarrow U(S)$ satisfies $\sigma(s)\sigma(t) = \sigma(st)$ whenever $st \neq 0$ and is universal for this property.
- S is called **strongly 0- E -unitary** if $\sigma^{-1}(1) = E(S) \setminus \{0\}$.

Strongly 0- E -unitary inverse semigroups

- An inverse semigroups with 0 cannot be E -unitary without being a semilattice.
- So one needs an appropriate variation of the concept for this context.
- If S is an inverse semigroup with 0, the **universal group** $U(S)$ is the group with generators $S \setminus \{0\}$ and relations given $s \cdot t = st$ if $st \neq 0$.
- The natural map $\sigma: S \setminus \{0\} \longrightarrow U(S)$ satisfies $\sigma(s)\sigma(t) = \sigma(st)$ whenever $st \neq 0$ and is universal for this property.
- S is called **strongly 0- E -unitary** if $\sigma^{-1}(1) = E(S) \setminus \{0\}$.

Strongly 0- E -unitary inverse semigroups

- An inverse semigroups with 0 cannot be E -unitary without being a semilattice.
- So one needs an appropriate variation of the concept for this context.
- If S is an inverse semigroup with 0, the **universal group** $U(S)$ is the group with generators $S \setminus \{0\}$ and relations given $s \cdot t = st$ if $st \neq 0$.
- The natural map $\sigma: S \setminus \{0\} \longrightarrow U(S)$ satisfies $\sigma(s)\sigma(t) = \sigma(st)$ whenever $st \neq 0$ and is universal for this property.
- S is called **strongly 0- E -unitary** if $\sigma^{-1}(1) = E(S) \setminus \{0\}$.

Strongly 0- E -unitary inverse semigroups

- An inverse semigroups with 0 cannot be E -unitary without being a semilattice.
- So one needs an appropriate variation of the concept for this context.
- If S is an inverse semigroup with 0, the **universal group** $U(S)$ is the group with generators $S \setminus \{0\}$ and relations given $s \cdot t = st$ if $st \neq 0$.
- The natural map $\sigma: S \setminus \{0\} \longrightarrow U(S)$ satisfies $\sigma(s)\sigma(t) = \sigma(st)$ whenever $st \neq 0$ and is universal for this property.
- S is called **strongly 0- E -unitary** if $\sigma^{-1}(1) = E(S) \setminus \{0\}$.

Strongly 0- E -unitary inverse semigroups II

- The **polycyclic inverse monoid** (or Cuntz semigroup) P_X is strongly 0- E -unitary with universal group the free group on X .
- The map σ takes uv^* to uv^{-1} .
- More generally, **graph inverse semigroups** are strongly 0- E -unitary.
- **Tiling inverse semigroups** are strongly 0- E -unitary (Lawson/Kellendonk).
- It seems most of the inverse semigroups appearing in the C^* -algebra literature are strongly 0- E -unitary.
- A graph immersion over a bouquet of circles can be extended to a regular covering iff the associated inverse semigroup is strongly 0- E -unitary.
- I showed it is undecidable whether a finite inverse semigroup is strongly 0- E -unitary.

Strongly 0- E -unitary inverse semigroups II

- The **polycyclic inverse monoid** (or Cuntz semigroup) P_X is strongly 0- E -unitary with universal group the free group on X .
- The map σ takes uv^* to uv^{-1} .
- More generally, **graph inverse semigroups** are strongly 0- E -unitary.
- **Tiling inverse semigroups** are strongly 0- E -unitary (Lawson/Kellendonk).
- It seems most of the inverse semigroups appearing in the C^* -algebra literature are strongly 0- E -unitary.
- A graph immersion over a bouquet of circles can be extended to a regular covering iff the associated inverse semigroup is strongly 0- E -unitary.
- I showed it is undecidable whether a finite inverse semigroup is strongly 0- E -unitary.

Strongly 0- E -unitary inverse semigroups II

- The **polycyclic inverse monoid** (or Cuntz semigroup) P_X is strongly 0- E -unitary with universal group the free group on X .
- The map σ takes uv^* to uv^{-1} .
- More generally, **graph inverse semigroups** are strongly 0- E -unitary.
- **Tiling inverse semigroups** are strongly 0- E -unitary (Lawson/Kellendonk).
- It seems most of the inverse semigroups appearing in the C^* -algebra literature are strongly 0- E -unitary.
- A graph immersion over a bouquet of circles can be extended to a regular covering iff the associated inverse semigroup is strongly 0- E -unitary.
- I showed it is undecidable whether a finite inverse semigroup is strongly 0- E -unitary.

Strongly 0- E -unitary inverse semigroups II

- The **polycyclic inverse monoid** (or Cuntz semigroup) P_X is strongly 0- E -unitary with universal group the free group on X .
- The map σ takes uv^* to uv^{-1} .
- More generally, **graph inverse semigroups** are strongly 0- E -unitary.
- **Tiling inverse semigroups** are strongly 0- E -unitary (Lawson/Kellendonk).
- It seems most of the inverse semigroups appearing in the C^* -algebra literature are strongly 0- E -unitary.
- A graph immersion over a bouquet of circles can be extended to a regular covering iff the associated inverse semigroup is strongly 0- E -unitary.
- I showed it is undecidable whether a finite inverse semigroup is strongly 0- E -unitary.

Strongly 0- E -unitary inverse semigroups II

- The **polycyclic inverse monoid** (or Cuntz semigroup) P_X is strongly 0- E -unitary with universal group the free group on X .
- The map σ takes uv^* to uv^{-1} .
- More generally, **graph inverse semigroups** are strongly 0- E -unitary.
- **Tiling inverse semigroups** are strongly 0- E -unitary (Lawson/Kellendonk).
- It seems most of the inverse semigroups appearing in the C^* -algebra literature are strongly 0- E -unitary.
- A graph immersion over a bouquet of circles can be extended to a regular covering iff the associated inverse semigroup is strongly 0- E -unitary.
- I showed it is undecidable whether a finite inverse semigroup is strongly 0- E -unitary.

Strongly 0- E -unitary inverse semigroups II

- The **polycyclic inverse monoid** (or Cuntz semigroup) P_X is strongly 0- E -unitary with universal group the free group on X .
- The map σ takes uv^* to uv^{-1} .
- More generally, **graph inverse semigroups** are strongly 0- E -unitary.
- **Tiling inverse semigroups** are strongly 0- E -unitary (Lawson/Kellendonk).
- It seems most of the inverse semigroups appearing in the C^* -algebra literature are strongly 0- E -unitary.
- A graph immersion over a bouquet of circles can be extended to a regular covering iff the associated inverse semigroup is strongly 0- E -unitary.
- I showed it is undecidable whether a finite inverse semigroup is strongly 0- E -unitary.

Strongly 0- E -unitary inverse semigroups II

- The **polycyclic inverse monoid** (or Cuntz semigroup) P_X is strongly 0- E -unitary with universal group the free group on X .
- The map σ takes uv^* to uv^{-1} .
- More generally, **graph inverse semigroups** are strongly 0- E -unitary.
- **Tiling inverse semigroups** are strongly 0- E -unitary (Lawson/Kellendonk).
- It seems most of the inverse semigroups appearing in the C^* -algebra literature are strongly 0- E -unitary.
- A graph immersion over a bouquet of circles can be extended to a regular covering iff the associated inverse semigroup is strongly 0- E -unitary.
- I showed it is undecidable whether a finite inverse semigroup is strongly 0- E -unitary.

Strongly 0- E -unitary inverse semigroups III

- It was observed independently by McAlister and by me and Margolis that S is strongly 0- E -unitary iff there is:
 - an E -unitary inverse semigroup T (with maximal group image $U(S)$)
 - and an ideal I such that $S \cong T/I$.
- This allows a partial action semidirect product type description of S and turns out to be useful for understanding the C^* -algebra of S .

Strongly 0- E -unitary inverse semigroups III

- It was observed independently by McAlister and by me and Margolis that S is strongly 0- E -unitary iff there is:
 - an E -unitary inverse semigroup T (with maximal group image $U(S)$)
 - and an ideal I such that $S \cong T/I$.
- This allows a partial action semidirect product type description of S and turns out to be useful for understanding the C^* -algebra of S .

Strongly 0- E -unitary inverse semigroups III

- It was observed independently by McAlister and by me and Margolis that S is strongly 0- E -unitary iff there is:
 - an E -unitary inverse semigroup T (with maximal group image $U(S)$)
 - and an ideal I such that $S \cong T/I$.
- This allows a partial action semidirect product type description of S and turns out to be useful for understanding the C^* -algebra of S .

Strongly 0- E -unitary inverse semigroups III

- It was observed independently by McAlister and by me and Margolis that S is strongly 0- E -unitary iff there is:
 - an E -unitary inverse semigroup T (with maximal group image $U(S)$)
 - and an ideal I such that $S \cong T/I$.
- This allows a partial action semidirect product type description of S and turns out to be useful for understanding the C^* -algebra of S .

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \rightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \rightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \rightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \rightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \rightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \rightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \rightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \rightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \rightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \rightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \rightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \rightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \rightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \rightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \rightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \rightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \rightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \rightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \longrightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \longrightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \longrightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \rightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \rightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \rightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \rightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \rightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \rightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

The groupoid of germs

- An **action** $S \curvearrowright X$ of an inverse semigroup on a space X is a homomorphism $\theta: S \rightarrow I_X$.
- We always assume that $X = \bigcup_{e \in E(S)} X_e$ where X_e is the domain of e .
- Note $\theta(s): X_{s^*s} \rightarrow X_{ss^*}$.
- Let us call the action **special** if X_e is clopen for each $e \in E(S)$.
- The groupoid of germs $S \ltimes X$ has arrow space $(S \times X)/\sim$ where $(s, x) \sim (t, y)$ iff $x = y$ and $\exists u \leq s, t$ with $x \in X_{u^*u}$.
- The unit space is X .
- $[s, x]: x \rightarrow sx$.
- $[s, ty][t, y] = [st, y]$, $[s, x]^{-1} = [s^*, sx]$.
- $S \ltimes X$ is an étale groupoid, but in general non-Hausdorff.
- When S is a group, this is the usual semidirect product.

Paterson's universal groupoid

- Fix an inverse semigroup S with semilattice of idempotents E .
- Let \widehat{E} be the **character space** of E , that is, the space of non-zero homomorphisms $E \rightarrow \{0, 1\}$ equipped with the topology of pointwise convergence.
- When working in the category of inverse semigroups with 0, we assume characters preserve 0.
- Elements of \widehat{E} can also be identified with **filters** on E .
- Then $S \curvearrowright \widehat{E}$ by putting $\widehat{E}_e = \{\varphi \mid \varphi(e) = 1\}$ and $s\varphi(f) = \varphi(s^*fs)$ for $\varphi \in \widehat{E}_{s^*s}$.
- This is a special action.
- The universal groupoid $\mathcal{G}(S)$ is the germ groupoid $S \ltimes \widehat{E}$.
- Paterson showed that the universal and reduced C^* -algebras of S are isomorphic to the corresponding algebras for $\mathcal{G}(S)$.

Paterson's universal groupoid

- Fix an inverse semigroup S with semilattice of idempotents E .
- Let \widehat{E} be the **character space** of E , that is, the space of non-zero homomorphisms $E \rightarrow \{0, 1\}$ equipped with the topology of pointwise convergence.
- When working in the category of inverse semigroups with 0, we assume characters preserve 0.
- Elements of \widehat{E} can also be identified with **filters** on E .
- Then $S \curvearrowright \widehat{E}$ by putting $\widehat{E}_e = \{\varphi \mid \varphi(e) = 1\}$ and $s\varphi(f) = \varphi(s^*fs)$ for $\varphi \in \widehat{E}_{s^*s}$.
- This is a special action.
- The universal groupoid $\mathcal{G}(S)$ is the germ groupoid $S \ltimes \widehat{E}$.
- Paterson showed that the universal and reduced C^* -algebras of S are isomorphic to the corresponding algebras for $\mathcal{G}(S)$.

Paterson's universal groupoid

- Fix an inverse semigroup S with semilattice of idempotents E .
- Let \hat{E} be the **character space** of E , that is, the space of non-zero homomorphisms $E \rightarrow \{0, 1\}$ equipped with the topology of pointwise convergence.
- When working in the category of inverse semigroups with 0, we assume characters preserve 0.
- Elements of \hat{E} can also be identified with **filters** on E .
- Then $S \curvearrowright \hat{E}$ by putting $\hat{E}_e = \{\varphi \mid \varphi(e) = 1\}$ and $s\varphi(f) = \varphi(s^*fs)$ for $\varphi \in \hat{E}_{s^*s}$.
- This is a special action.
- The universal groupoid $\mathcal{G}(S)$ is the germ groupoid $S \ltimes \hat{E}$.
- Paterson showed that the universal and reduced C^* -algebras of S are isomorphic to the corresponding algebras for $\mathcal{G}(S)$.

Paterson's universal groupoid

- Fix an inverse semigroup S with semilattice of idempotents E .
- Let \hat{E} be the **character space** of E , that is, the space of non-zero homomorphisms $E \rightarrow \{0, 1\}$ equipped with the topology of pointwise convergence.
- When working in the category of inverse semigroups with 0, we assume characters preserve 0.
- Elements of \hat{E} can also be identified with **filters** on E .
- Then $S \curvearrowright \hat{E}$ by putting $\hat{E}_e = \{\varphi \mid \varphi(e) = 1\}$ and $s\varphi(f) = \varphi(s^*fs)$ for $\varphi \in \hat{E}_{s^*s}$.
- This is a special action.
- The **universal groupoid** $\mathcal{G}(S)$ is the germ groupoid $S \times \hat{E}$.
- Paterson showed that the universal and reduced C^* -algebras of S are isomorphic to the corresponding algebras for $\mathcal{G}(S)$.

Paterson's universal groupoid

- Fix an inverse semigroup S with semilattice of idempotents E .
- Let \hat{E} be the **character space** of E , that is, the space of non-zero homomorphisms $E \rightarrow \{0, 1\}$ equipped with the topology of pointwise convergence.
- When working in the category of inverse semigroups with 0, we assume characters preserve 0.
- Elements of \hat{E} can also be identified with **filters** on E .
- Then $S \curvearrowright \hat{E}$ by putting $\hat{E}_e = \{\varphi \mid \varphi(e) = 1\}$ and $s\varphi(f) = \varphi(s^*fs)$ for $\varphi \in \hat{E}_{s^*s}$.
- This is a special action.
- The **universal groupoid** $\mathcal{G}(S)$ is the germ groupoid $S \ltimes \hat{E}$.
- Paterson showed that the universal and reduced C^* -algebras of S are isomorphic to the corresponding algebras for $\mathcal{G}(S)$.

Paterson's universal groupoid

- Fix an inverse semigroup S with semilattice of idempotents E .
- Let \hat{E} be the **character space** of E , that is, the space of non-zero homomorphisms $E \rightarrow \{0, 1\}$ equipped with the topology of pointwise convergence.
- When working in the category of inverse semigroups with 0, we assume characters preserve 0.
- Elements of \hat{E} can also be identified with **filters** on E .
- Then $S \curvearrowright \hat{E}$ by putting $\hat{E}_e = \{\varphi \mid \varphi(e) = 1\}$ and $s\varphi(f) = \varphi(s^*fs)$ for $\varphi \in \hat{E}_{s^*s}$.
- This is a special action.
- The **universal groupoid** $\mathcal{G}(S)$ is the germ groupoid $S \ltimes \hat{E}$.
- Paterson showed that the universal and reduced C^* -algebras of S are isomorphic to the corresponding algebras for $\mathcal{G}(S)$.

Paterson's universal groupoid

- Fix an inverse semigroup S with semilattice of idempotents E .
- Let \hat{E} be the **character space** of E , that is, the space of non-zero homomorphisms $E \rightarrow \{0, 1\}$ equipped with the topology of pointwise convergence.
- When working in the category of inverse semigroups with 0, we assume characters preserve 0.
- Elements of \hat{E} can also be identified with **filters** on E .
- Then $S \curvearrowright \hat{E}$ by putting $\hat{E}_e = \{\varphi \mid \varphi(e) = 1\}$ and $s\varphi(f) = \varphi(s^*fs)$ for $\varphi \in \hat{E}_{s^*s}$.
- This is a special action.
- The **universal groupoid** $\mathcal{G}(S)$ is the germ groupoid $S \ltimes \hat{E}$.
- Paterson showed that the universal and reduced C^* -algebras of S are isomorphic to the corresponding algebras for $\mathcal{G}(S)$.

Paterson's universal groupoid

- Fix an inverse semigroup S with semilattice of idempotents E .
- Let \hat{E} be the **character space** of E , that is, the space of non-zero homomorphisms $E \rightarrow \{0, 1\}$ equipped with the topology of pointwise convergence.
- When working in the category of inverse semigroups with 0, we assume characters preserve 0.
- Elements of \hat{E} can also be identified with **filters** on E .
- Then $S \curvearrowright \hat{E}$ by putting $\hat{E}_e = \{\varphi \mid \varphi(e) = 1\}$ and $s\varphi(f) = \varphi(s^*fs)$ for $\varphi \in \hat{E}_{s^*s}$.
- This is a special action.
- The **universal groupoid** $\mathcal{G}(S)$ is the germ groupoid $S \ltimes \hat{E}$.
- Paterson showed that the universal and reduced C^* -algebras of S are isomorphic to the corresponding algebras for $\mathcal{G}(S)$.

A result of Khoshkam and Skandalis

- Khoshkam and Skandalis proved the following result.

Theorem

Let S be an E -unitary inverse semigroup such that $eSf \cap \sigma^{-1}(g)$ is finitely generated as a downset for all $e, f \in E(S)$ and $g \in G(S)$. Then there is an action $G(S) \curvearrowright X$ such that $\mathcal{G}(S)$ is Morita equivalent to $G(S) \ltimes X$.

- F -inverse semigroups satisfy the Khoshkam-Skandalis condition since $eSf \cap \sigma^{-1}(g)$ has a maximum.
- The proof uses a generalization of the enveloping action of a partial group action.
- An inverse semigroup S is locally E -unitary if eSe is E -unitary for all $e \in E(S)$.
- We have observed that the Khoshkam-Skandalis result also applies to locally E -unitary inverse semigroups.

A result of Khoshkam and Skandalis

- Khoshkam and Skandalis proved the following result.

Theorem

*Let S be an E -unitary inverse semigroup such that $eSf \cap \sigma^{-1}(g)$ is finitely generated as a downset for all $e, f \in E(S)$ and $g \in G(S)$. Then there is an action $G(S) \curvearrowright X$ such that $\mathcal{G}(S)$ is **Morita equivalent** to $G(S) \ltimes X$.*

- F -inverse semigroups satisfy the Khoshkam-Skandalis condition since $eSf \cap \sigma^{-1}(g)$ has a maximum.
- The proof uses a generalization of the enveloping action of a partial group action.
- An inverse semigroup S is locally E -unitary if eSe is E -unitary for all $e \in E(S)$.
- We have observed that the Khoshkam-Skandalis result also applies to locally E -unitary inverse semigroups.

A result of Khoshkam and Skandalis

- Khoshkam and Skandalis proved the following result.

Theorem

*Let S be an E -unitary inverse semigroup such that $eSf \cap \sigma^{-1}(g)$ is finitely generated as a downset for all $e, f \in E(S)$ and $g \in G(S)$. Then there is an action $G(S) \curvearrowright X$ such that $\mathcal{G}(S)$ is **Morita equivalent** to $G(S) \ltimes X$.*

- F -inverse semigroups satisfy the Khoshkam-Skandalis condition since $eSf \cap \sigma^{-1}(g)$ has a maximum.
- The proof uses a generalization of the enveloping action of a partial group action.
- An inverse semigroup S is **locally E -unitary** if eSe is E -unitary for all $e \in E(S)$.
- We have observed that the Khoshkam-Skandalis result also applies to locally E -unitary inverse semigroups.

A result of Khoshkam and Skandalis

- Khoshkam and Skandalis proved the following result.

Theorem

*Let S be an E -unitary inverse semigroup such that $eSf \cap \sigma^{-1}(g)$ is finitely generated as a downset for all $e, f \in E(S)$ and $g \in G(S)$. Then there is an action $G(S) \curvearrowright X$ such that $\mathcal{G}(S)$ is **Morita equivalent** to $G(S) \ltimes X$.*

- F -inverse semigroups satisfy the Khoshkam-Skandalis condition since $eSf \cap \sigma^{-1}(g)$ has a maximum.
- The proof uses a generalization of the enveloping action of a partial group action.
- An inverse semigroup S is **locally E -unitary** if eSe is E -unitary for all $e \in E(S)$.
- We have observed that the Khoshkam-Skandalis result also applies to locally E -unitary inverse semigroups.

A result of Khoshkam and Skandalis

- Khoshkam and Skandalis proved the following result.

Theorem

*Let S be an E -unitary inverse semigroup such that $eSf \cap \sigma^{-1}(g)$ is finitely generated as a downset for all $e, f \in E(S)$ and $g \in G(S)$. Then there is an action $G(S) \curvearrowright X$ such that $\mathcal{G}(S)$ is **Morita equivalent** to $G(S) \ltimes X$.*

- F -inverse semigroups satisfy the Khoshkam-Skandalis condition since $eSf \cap \sigma^{-1}(g)$ has a maximum.
- The proof uses a generalization of the enveloping action of a partial group action.
- An inverse semigroup S is **locally E -unitary** if eSe is E -unitary for all $e \in E(S)$.
- We have observed that the Khoshkam-Skandalis result also applies to locally E -unitary inverse semigroups.

A result of Khoshkam and Skandalis

- Khoshkam and Skandalis proved the following result.

Theorem

*Let S be an E -unitary inverse semigroup such that $eSf \cap \sigma^{-1}(g)$ is finitely generated as a downset for all $e, f \in E(S)$ and $g \in G(S)$. Then there is an action $G(S) \curvearrowright X$ such that $\mathcal{G}(S)$ is **Morita equivalent** to $G(S) \ltimes X$.*

- F -inverse semigroups satisfy the Khoshkam-Skandalis condition since $eSf \cap \sigma^{-1}(g)$ has a maximum.
- The proof uses a generalization of the enveloping action of a partial group action.
- An inverse semigroup S is **locally E -unitary** if eSe is E -unitary for all $e \in E(S)$.
- We have observed that the Khoshkam-Skandalis result also applies to locally E -unitary inverse semigroups.

Khoshkam-Skandalis from a partial action viewpoint

- There is another way to view the Khoshkam-Skandalis result.

Theorem (Milan, BS)

Let S be an E -unitary inverse semigroup. Then there is a partial action of $G(S)$ on $\widehat{E(S)}$ such that $\mathcal{G}(S) \cong G(S) \ltimes \widehat{E(S)}$.

- Here the partial action semidirect product is as in the sense of Abadie.
- The enveloping action is Hausdorff precisely when the Khoshkam-Skandalis condition holds and so Abadie's results also imply the Morita equivalence.
- In the locally E -unitary case there is no partial group action.

Khoshkam-Skandalis from a partial action viewpoint

- There is another way to view the Khoshkam-Skandalis result.

Theorem (Milan, BS)

Let S be an E -unitary inverse semigroup. Then there is a partial action of $G(S)$ on $\widehat{E(S)}$ such that $\mathcal{G}(S) \cong G(S) \ltimes \widehat{E(S)}$.

- Here the partial action semidirect product is as in the sense of Abadie.
- The enveloping action is Hausdorff precisely when the Khoshkam-Skandalis condition holds and so Abadie's results also imply the Morita equivalence.
- In the locally E -unitary case there is no partial group action.

Khoshkam-Skandalis from a partial action viewpoint

- There is another way to view the Khoshkam-Skandalis result.

Theorem (Milan, BS)

Let S be an E -unitary inverse semigroup. Then there is a partial action of $G(S)$ on $\widehat{E(S)}$ such that $\mathcal{G}(S) \cong G(S) \ltimes \widehat{E(S)}$.

- Here the partial action semidirect product is as in the sense of Abadie.
- The enveloping action is Hausdorff precisely when the Khoshkam-Skandalis condition holds and so Abadie's results also imply the Morita equivalence.
- In the locally E -unitary case there is no partial group action.

Khoshkam-Skandalis from a partial action viewpoint

- There is another way to view the Khoshkam-Skandalis result.

Theorem (Milan, BS)

Let S be an E -unitary inverse semigroup. Then there is a partial action of $G(S)$ on $\widehat{E(S)}$ such that $\mathcal{G}(S) \cong G(S) \ltimes \widehat{E(S)}$.

- Here the partial action semidirect product is as in the sense of Abadie.
- The enveloping action is Hausdorff precisely when the Khoshkam-Skandalis condition holds and so Abadie's results also imply the Morita equivalence.
- In the locally E -unitary case there is no partial group action.

Khoshkam-Skandalis from a partial action viewpoint

- There is another way to view the Khoshkam-Skandalis result.

Theorem (Milan, BS)

Let S be an E -unitary inverse semigroup. Then there is a partial action of $G(S)$ on $\widehat{E(S)}$ such that $\mathcal{G}(S) \cong G(S) \ltimes \widehat{E(S)}$.

- Here the partial action semidirect product is as in the sense of Abadie.
- The enveloping action is Hausdorff precisely when the Khoshkam-Skandalis condition holds and so Abadie's results also imply the Morita equivalence.
- In the locally E -unitary case there is no partial group action.

The strongly 0- E -unitary case

- Let S be an inverse semigroup and I an ideal.
- Then $\widehat{E(S/I)}$ can be identified with those characters of $E(S)$ vanishing on I .
- If S is E -unitary, then $\widehat{E(S/I)}$ is invariant under the partial action of $G(S)$.

Theorem (Milan, BS)

If S is strongly 0- E -unitary with universal group $U(S)$, then $\mathcal{G}(S) \cong U(S) \ltimes \widehat{E}(S)$ for an appropriate partial action of $U(S)$.

- A similar result holds for Exel's tight groupoid and explains a number of partial action cross product results in the literature.
- An analogue of the Khoshkam-Skandalis result holds in the strongly 0- E -unitary setting.

The strongly 0- E -unitary case

- Let S be an inverse semigroup and I an ideal.
- Then $\widehat{E(S/I)}$ can be identified with those characters of $E(S)$ vanishing on I .
- If S is E -unitary, then $\widehat{E(S/I)}$ is invariant under the partial action of $G(S)$.

Theorem (Milan, BS)

If S is strongly 0- E -unitary with universal group $U(S)$, then $\mathcal{G}(S) \cong U(S) \ltimes \widehat{E}(S)$ for an appropriate partial action of $U(S)$.

- A similar result holds for Exel's tight groupoid and explains a number of partial action cross product results in the literature.
- An analogue of the Khoshkam-Skandalis result holds in the strongly 0- E -unitary setting.

The strongly 0- E -unitary case

- Let S be an inverse semigroup and I an ideal.
- Then $\widehat{E(S/I)}$ can be identified with those characters of $E(S)$ vanishing on I .
- If S is E -unitary, then $\widehat{E(S/I)}$ is invariant under the partial action of $G(S)$.

Theorem (Milan, BS)

If S is strongly 0- E -unitary with universal group $U(S)$, then $\mathcal{G}(S) \cong U(S) \ltimes \widehat{E}(S)$ for an appropriate partial action of $U(S)$.

- A similar result holds for Exel's tight groupoid and explains a number of partial action cross product results in the literature.
- An analogue of the Khoshkam-Skandalis result holds in the strongly 0- E -unitary setting.

The strongly 0- E -unitary case

- Let S be an inverse semigroup and I an ideal.
- Then $\widehat{E(S/I)}$ can be identified with those characters of $E(S)$ vanishing on I .
- If S is E -unitary, then $\widehat{E(S/I)}$ is invariant under the partial action of $G(S)$.

Theorem (Milan, BS)

If S is strongly 0- E -unitary with universal group $U(S)$, then $\mathcal{G}(S) \cong U(S) \ltimes \widehat{E}(S)$ for an appropriate partial action of $U(S)$.

- A similar result holds for Exel's tight groupoid and explains a number of partial action cross product results in the literature.
- An analogue of the Khoshkam-Skandalis result holds in the strongly 0- E -unitary setting.

The strongly 0- E -unitary case

- Let S be an inverse semigroup and I an ideal.
- Then $\widehat{E(S/I)}$ can be identified with those characters of $E(S)$ vanishing on I .
- If S is E -unitary, then $\widehat{E(S/I)}$ is invariant under the partial action of $G(S)$.

Theorem (Milan, BS)

If S is strongly 0- E -unitary with universal group $U(S)$, then $\mathcal{G}(S) \cong U(S) \ltimes \widehat{E}(S)$ for an appropriate partial action of $U(S)$.

- A similar result holds for Exel's tight groupoid and explains a number of partial action cross product results in the literature.
- An analogue of the Khoshkam-Skandalis result holds in the strongly 0- E -unitary setting.

The strongly 0- E -unitary case

- Let S be an inverse semigroup and I an ideal.
- Then $\widehat{E(S/I)}$ can be identified with those characters of $E(S)$ vanishing on I .
- If S is E -unitary, then $\widehat{E(S/I)}$ is invariant under the partial action of $G(S)$.

Theorem (Milan, BS)

If S is strongly 0- E -unitary with universal group $U(S)$, then $\mathcal{G}(S) \cong U(S) \ltimes \widehat{E}(S)$ for an appropriate partial action of $U(S)$.

- A similar result holds for Exel's tight groupoid and explains a number of partial action cross product results in the literature.
- An analogue of the Khoshkam-Skandalis result holds in the strongly 0- E -unitary setting.

- The Khoshkam-Skandalis result gives a condition that guarantees $C^*(S)$ is strongly Morita equivalent to $C_0(X) \rtimes G(S)$ for some action $G(S) \curvearrowright X$.
- What are conditions on a homomorphism $S \xrightarrow{\varphi} T$ that guarantee that $C^*(S)$ is strongly Morita equivalent to a cross product $C_0(X) \rtimes T$?
- In other words, when is $\mathcal{G}(S)$ Morita equivalent to a groupoid of germs of an action of T ?
- First we need to address the issue of functoriality of the universal groupoid.

- The Khoshkam-Skandalis result gives a condition that guarantees $C^*(S)$ is strongly Morita equivalent to $C_0(X) \rtimes G(S)$ for some action $G(S) \curvearrowright X$.
- What are conditions on a homomorphism $S \xrightarrow{\varphi} T$ that guarantee that $C^*(S)$ is strongly Morita equivalent to a cross product $C_0(X) \rtimes T$?
- In other words, when is $\mathcal{G}(S)$ Morita equivalent to a groupoid of germs of an action of T ?
- First we need to address the issue of functoriality of the universal groupoid.

- The Khoshkam-Skandalis result gives a condition that guarantees $C^*(S)$ is strongly Morita equivalent to $C_0(X) \rtimes G(S)$ for some action $G(S) \curvearrowright X$.
- What are conditions on a homomorphism $S \xrightarrow{\varphi} T$ that guarantee that $C^*(S)$ is strongly Morita equivalent to a cross product $C_0(X) \rtimes T$?
- In other words, when is $\mathcal{G}(S)$ Morita equivalent to a groupoid of germs of an action of T ?
- First we need to address the issue of functoriality of the universal groupoid.

- The Khoshkam-Skandalis result gives a condition that guarantees $C^*(S)$ is strongly Morita equivalent to $C_0(X) \rtimes G(S)$ for some action $G(S) \curvearrowright X$.
- What are conditions on a homomorphism $S \xrightarrow{\varphi} T$ that guarantee that $C^*(S)$ is strongly Morita equivalent to a cross product $C_0(X) \rtimes T$?
- In other words, when is $\mathcal{G}(S)$ Morita equivalent to a groupoid of germs of an action of T ?
- First we need to address the issue of functoriality of the universal groupoid.

- If $E \xrightarrow{\varphi} F$ is a semilattice homomorphism, then there is a natural cts map $\widehat{F} \longrightarrow \widehat{E}$ given by “pulling back.”
- But we want a map the other way — and there is one.
- If \mathcal{F} is a filter on E , then $\varphi(\mathcal{F})$ is a filter base on F and hence yields a filter $\varphi_*(\mathcal{F})$.
- However, φ_* need not be cts.

- If $E \xrightarrow{\varphi} F$ is a semilattice homomorphism, then there is a natural cts map $\hat{F} \longrightarrow \hat{E}$ given by “pulling back.”
- But we want a map the other way — and there is one.
- If \mathcal{F} is a filter on E , then $\varphi(\mathcal{F})$ is a filter base on F and hence yields a filter $\varphi_*(\mathcal{F})$.
- However, φ_* need not be cts.

- If $E \xrightarrow{\varphi} F$ is a semilattice homomorphism, then there is a natural cts map $\widehat{F} \longrightarrow \widehat{E}$ given by “pulling back.”
- But we want a map the other way — and there is one.
- If \mathcal{F} is a filter on E , then $\varphi(\mathcal{F})$ is a filter base on F and hence yields a filter $\varphi_*(\mathcal{F})$.
- However, φ_* need not be cts.

- If $E \xrightarrow{\varphi} F$ is a semilattice homomorphism, then there is a natural cts map $\widehat{F} \longrightarrow \widehat{E}$ given by “pulling back.”
- But we want a map the other way — and there is one.
- If \mathcal{F} is a filter on E , then $\varphi(\mathcal{F})$ is a filter base on F and hence yields a filter $\varphi_*(\mathcal{F})$.
- However, φ_* need not be cts.

Functoriality II

- The **Alexandrov topology** on a poset takes the downsets as the open sets. (It is T_0 .)
- The quasi-compact open sets are the finitely generated downsets.
- A map of spaces is **coherent** if the inverse image of a quasi-compact open is quasi-compact open.
- It is a folklore fact from the theory of locales that φ_* is **cts** iff φ is **locally coherent**.

Theorem (Milan,BS)

The universal groupoid is functorial on the category of inverse semigroups with maps that are locally coherent on the semilattices of idempotents.

- The maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ belongs to this category for instance.

Functoriality II

- The **Alexandrov topology** on a poset takes the downsets as the open sets. (It is T_0 .)
- The quasi-compact open sets are the finitely generated downsets.
- A map of spaces is **coherent** if the inverse image of a quasi-compact open is quasi-compact open.
- It is a folklore fact from the theory of locales that φ_* is **cts** iff φ is **locally coherent**.

Theorem (Milan,BS)

The universal groupoid is functorial on the category of inverse semigroups with maps that are locally coherent on the semilattices of idempotents.

- The maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ belongs to this category for instance.

Functoriality II

- The **Alexandrov topology** on a poset takes the downsets as the open sets. (It is T_0 .)
- The quasi-compact open sets are the finitely generated downsets.
- A map of spaces is **coherent** if the inverse image of a quasi-compact open is quasi-compact open.
- It is a folklore fact from the theory of locales that φ_* is **cts** iff φ is **locally coherent**.

Theorem (Milan,BS)

The universal groupoid is functorial on the category of inverse semigroups with maps that are locally coherent on the semilattices of idempotents.

- The maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ belongs to this category for instance.

Functoriality II

- The **Alexandrov topology** on a poset takes the downsets as the open sets. (It is T_0 .)
- The quasi-compact open sets are the finitely generated downsets.
- A map of spaces is **coherent** if the inverse image of a quasi-compact open is quasi-compact open.
- It is a folklore fact from the theory of locales that φ_* is **cts** iff φ is **locally coherent**.

Theorem (Milan,BS)

The universal groupoid is functorial on the category of inverse semigroups with maps that are locally coherent on the semilattices of idempotents.

- The maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ belongs to this category for instance.

Functoriality II

- The **Alexandrov topology** on a poset takes the downsets as the open sets. (It is T_0 .)
- The quasi-compact open sets are the finitely generated downsets.
- A map of spaces is **coherent** if the inverse image of a quasi-compact open is quasi-compact open.
- It is a folklore fact from the theory of locales that φ_* is **cts** iff φ is **locally coherent**.

Theorem (Milan,BS)

The universal groupoid is functorial on the category of inverse semigroups with maps that are locally coherent on the semilattices of idempotents.

- The maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ belongs to this category for instance.

Functoriality II

- The **Alexandrov topology** on a poset takes the downsets as the open sets. (It is T_0 .)
- The quasi-compact open sets are the finitely generated downsets.
- A map of spaces is **coherent** if the inverse image of a quasi-compact open is quasi-compact open.
- It is a folklore fact from the theory of locales that φ_* is **cts** iff φ is **locally coherent**.

Theorem (Milan,BS)

The universal groupoid is functorial on the category of inverse semigroups with maps that are locally coherent on the semilattices of idempotents.

- The maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ belongs to this category for instance.

Idempotent-pure homomorphisms

- An inverse semigroup homomorphism $S \xrightarrow{\varphi} T$ is **idempotent-pure** if $\varphi^{-1}(E(T)) = E(S)$.
- For example, S is **E -unitary** iff the maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ is idempotent-pure.
- O'Carroll showed that every idempotent-pure morphism is equivalent to a partial action semidirect product projection.
- Let us say $S \xrightarrow{\varphi} T$ is **locally idempotent-pure** if $\varphi|_{eSe}$ is idempotent-pure for each $e \in E(S)$.
- So S is **locally E -unitary** iff $S \xrightarrow{\sigma} G(S)$ is locally idempotent-pure.

Idempotent-pure homomorphisms

- An inverse semigroup homomorphism $S \xrightarrow{\varphi} T$ is **idempotent-pure** if $\varphi^{-1}(E(T)) = E(S)$.
- For example, S is **E -unitary** iff the maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ is idempotent-pure.
- O'Carroll showed that every idempotent-pure morphism is equivalent to a partial action semidirect product projection.
- Let us say $S \xrightarrow{\varphi} T$ is **locally idempotent-pure** if $\varphi|_{eSe}$ is idempotent-pure for each $e \in E(S)$.
- So S is **locally E -unitary** iff $S \xrightarrow{\sigma} G(S)$ is locally idempotent-pure.

Idempotent-pure homomorphisms

- An inverse semigroup homomorphism $S \xrightarrow{\varphi} T$ is **idempotent-pure** if $\varphi^{-1}(E(T)) = E(S)$.
- For example, S is **E -unitary** iff the maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ is idempotent-pure.
- O'Carroll showed that every idempotent-pure morphism is equivalent to a partial action semidirect product projection.
- Let us say $S \xrightarrow{\varphi} T$ is **locally idempotent-pure** if $\varphi|_{eSe}$ is idempotent-pure for each $e \in E(S)$.
- So S is **locally E -unitary** iff $S \xrightarrow{\sigma} G(S)$ is locally idempotent-pure.

Idempotent-pure homomorphisms

- An inverse semigroup homomorphism $S \xrightarrow{\varphi} T$ is **idempotent-pure** if $\varphi^{-1}(E(T)) = E(S)$.
- For example, S is **E -unitary** iff the maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ is idempotent-pure.
- O'Carroll showed that every idempotent-pure morphism is equivalent to a partial action semidirect product projection.
- Let us say $S \xrightarrow{\varphi} T$ is **locally idempotent-pure** if $\varphi|_{eSe}$ is idempotent-pure for each $e \in E(S)$.
- So S is **locally E -unitary** iff $S \xrightarrow{\sigma} G(S)$ is locally idempotent-pure.

Idempotent-pure homomorphisms

- An inverse semigroup homomorphism $S \xrightarrow{\varphi} T$ is **idempotent-pure** if $\varphi^{-1}(E(T)) = E(S)$.
- For example, S is **E -unitary** iff the maximal group image homomorphism $S \xrightarrow{\sigma} G(S)$ is idempotent-pure.
- O'Carroll showed that every idempotent-pure morphism is equivalent to a partial action semidirect product projection.
- Let us say $S \xrightarrow{\varphi} T$ is **locally idempotent-pure** if $\varphi|_{eSe}$ is idempotent-pure for each $e \in E(S)$.
- So S is **locally E -unitary** iff $S \xrightarrow{\sigma} G(S)$ is locally idempotent-pure.

The Khoshkam-Skandalis condition

- Inspired by the result of Khoshkam and Skandalis, we say $S \xrightarrow{\varphi} T$ satisfies the **Khoshkam-Skandalis (KS) condition** if $\varphi|_{eSf}$ is coherent for all $e, f \in E(S)$.
- For the case of $S \xrightarrow{\sigma} G(S)$, this is the original Khoshkam-Skandalis condition.

Theorem (Milan, BS)

*Let $S \xrightarrow{\varphi} T$ be a locally idempotent-pure homomorphism satisfying the Khoshkam-Skandalis condition. Then there is an **action** $T \curvearrowright X$ such that $\mathcal{G}(S)$ is **Morita equivalent** to the groupoid of germs $T \ltimes X$.*

The Khoshkam-Skandalis condition

- Inspired by the result of Khoshkam and Skandalis, we say $S \xrightarrow{\varphi} T$ satisfies the **Khoshkam-Skandalis (KS) condition** if $\varphi|_{eSf}$ is coherent for all $e, f \in E(S)$.
- For the case of $S \xrightarrow{\sigma} G(S)$, this is the original Khoshkam-Skandalis condition.

Theorem (Milan, BS)

*Let $S \xrightarrow{\varphi} T$ be a locally idempotent-pure homomorphism satisfying the Khoshkam-Skandalis condition. Then there is an **action** $T \curvearrowright X$ such that $\mathcal{G}(S)$ is **Morita equivalent** to the groupoid of germs $T \ltimes X$.*

The Khoshkam-Skandalis condition

- Inspired by the result of Khoshkam and Skandalis, we say $S \xrightarrow{\varphi} T$ satisfies the **Khoshkam-Skandalis (KS) condition** if $\varphi|_{eSf}$ is coherent for all $e, f \in E(S)$.
- For the case of $S \xrightarrow{\sigma} G(S)$, this is the original Khoshkam-Skandalis condition.

Theorem (Milan, BS)

*Let $S \xrightarrow{\varphi} T$ be a locally idempotent-pure homomorphism satisfying the Khoshkam-Skandalis condition. Then there is an **action** $T \curvearrowright X$ such that $\mathcal{G}(S)$ is **Morita equivalent** to the groupoid of germs $T \ltimes X$.*

- Khoshkam and Skandalis proved their result along the following lines.
- First they gave a general condition on a functor $\mathcal{G} \longrightarrow G$ from a groupoid to a group (a cocycle) that guarantees \mathcal{G} is Morita equivalent to $G \ltimes X$ for an appropriate action $G \curvearrowright X$.
- The results for inverse semigroups are obtained by considering the cocycle $\mathcal{G}(S) \longrightarrow G(S)$.
- Since an inverse semigroup is not an étale groupoid, we need to be more careful in how to proceed in the general case.

- Khoshkam and Skandalis proved their result along the following lines.
- First they gave a general condition on a functor $\mathcal{G} \rightarrow G$ from a groupoid to a group (a cocycle) that guarantees \mathcal{G} is Morita equivalent to $G \ltimes X$ for an appropriate action $G \curvearrowright X$.
- The results for inverse semigroups are obtained by considering the cocycle $\mathcal{G}(S) \rightarrow G(S)$.
- Since an inverse semigroup is not an étale groupoid, we need to be more careful in how to proceed in the general case.

- Khoshkam and Skandalis proved their result along the following lines.
- First they gave a general condition on a functor $\mathcal{G} \rightarrow G$ from a groupoid to a group (a cocycle) that guarantees \mathcal{G} is Morita equivalent to $G \ltimes X$ for an appropriate action $G \curvearrowright X$.
- The results for inverse semigroups are obtained by considering the cocycle $\mathcal{G}(S) \rightarrow G(S)$.
- Since an inverse semigroup is not an étale groupoid, we need to be more careful in how to proceed in the general case.

- Khoshkam and Skandalis proved their result along the following lines.
- First they gave a general condition on a functor $\mathcal{G} \longrightarrow G$ from a groupoid to a group (a cocycle) that guarantees \mathcal{G} is Morita equivalent to $G \ltimes X$ for an appropriate action $G \curvearrowright X$.
- The results for inverse semigroups are obtained by considering the cocycle $\mathcal{G}(S) \longrightarrow G(S)$.
- Since an inverse semigroup is not an étale groupoid, we need to be more careful in how to proceed in the general case.

Actions of groupoids

- An **action** $\mathcal{G} \curvearrowright X$ of a groupoid \mathcal{G} on a space X consists of a cts map $X \xrightarrow{p} \mathcal{G}^0$ and a map $\theta: \mathcal{G} \times_{\mathcal{G}^0} X \rightarrow X$ such that:
 - $p(x)x = x$;
 - $g(hx) = (gh)x$;
 - $p(gx) = r(g)$.
- The semidirect product $\mathcal{G} \ltimes X$ has arrow space $\mathcal{G} \times_{\mathcal{G}^0} X$.
- The unit space is X .
- One has $x \xrightarrow{(g,x)} gx$.
- $(g, hx)(h, x) = (gh, x)$, $(g, x)^{-1} = (g^{-1}, gx)$.
- If \mathcal{G} is étale, then so is $\mathcal{G} \ltimes X$.

Actions of groupoids

- An **action** $\mathcal{G} \curvearrowright X$ of a groupoid \mathcal{G} on a space X consists of a cts map $X \xrightarrow{p} \mathcal{G}^0$ and a map $\theta: \mathcal{G} \times_{\mathcal{G}^0} X \rightarrow X$ such that:
 - $p(x)x = x$;
 - $g(hx) = (gh)x$;
 - $p(gx) = r(g)$.
- The **semidirect product** $\mathcal{G} \ltimes X$ has arrow space $\mathcal{G} \times_{\mathcal{G}^0} X$.
- The unit space is X .
- One has $x \xrightarrow{(g,x)} gx$.
- $(g, hx)(h, x) = (gh, x)$, $(g, x)^{-1} = (g^{-1}, gx)$.
- If \mathcal{G} is étale, then so is $\mathcal{G} \ltimes X$.

Actions of groupoids

- An **action** $\mathcal{G} \curvearrowright X$ of a groupoid \mathcal{G} on a space X consists of a cts map $X \xrightarrow{p} \mathcal{G}^0$ and a map $\theta: \mathcal{G} \times_{\mathcal{G}^0} X \rightarrow X$ such that:
 - $p(x)x = x$;
 - $g(hx) = (gh)x$;
 - $p(gx) = r(g)$.
- The **semidirect product** $\mathcal{G} \ltimes X$ has arrow space $\mathcal{G} \times_{\mathcal{G}^0} X$.
- The unit space is X .
- One has $x \xrightarrow{(g,x)} gx$.
- $(g, hx)(h, x) = (gh, x)$, $(g, x)^{-1} = (g^{-1}, gx)$.
- If \mathcal{G} is étale, then so is $\mathcal{G} \ltimes X$.

Actions of groupoids

- An **action** $\mathcal{G} \curvearrowright X$ of a groupoid \mathcal{G} on a space X consists of a cts map $X \xrightarrow{p} \mathcal{G}^0$ and a map $\theta: \mathcal{G} \times_{\mathcal{G}^0} X \rightarrow X$ such that:
 - $p(x)x = x$;
 - $g(hx) = (gh)x$;
 - $p(gx) = r(g)$.
- The **semidirect product** $\mathcal{G} \ltimes X$ has arrow space $\mathcal{G} \times_{\mathcal{G}^0} X$.
- The unit space is X .
- One has $x \xrightarrow{(g,x)} gx$.
- $(g, hx)(h, x) = (gh, x)$, $(g, x)^{-1} = (g^{-1}, gx)$.
- If \mathcal{G} is étale, then so is $\mathcal{G} \ltimes X$.

Actions of groupoids

- An **action** $\mathcal{G} \curvearrowright X$ of a groupoid \mathcal{G} on a space X consists of a cts map $X \xrightarrow{p} \mathcal{G}^0$ and a map $\theta: \mathcal{G} \times_{\mathcal{G}^0} X \rightarrow X$ such that:
 - $p(x)x = x$;
 - $g(hx) = (gh)x$;
 - $p(gx) = r(g)$.
- The **semidirect product** $\mathcal{G} \ltimes X$ has arrow space $\mathcal{G} \times_{\mathcal{G}^0} X$.
- The unit space is X .
- One has $x \xrightarrow{(g,x)} gx$.
- $(g, hx)(h, x) = (gh, x)$, $(g, x)^{-1} = (g^{-1}, gx)$.
- If \mathcal{G} is étale, then so is $\mathcal{G} \ltimes X$.

Actions of groupoids

- An **action** $\mathcal{G} \curvearrowright X$ of a groupoid \mathcal{G} on a space X consists of a cts map $X \xrightarrow{p} \mathcal{G}^0$ and a map $\theta: \mathcal{G} \times_{\mathcal{G}^0} X \rightarrow X$ such that:
 - $p(x)x = x$;
 - $g(hx) = (gh)x$;
 - $p(gx) = r(g)$.
- The **semidirect product** $\mathcal{G} \ltimes X$ has arrow space $\mathcal{G} \times_{\mathcal{G}^0} X$.
- The unit space is X .
- One has $x \xrightarrow{(g,x)} gx$.
- $(g, hx)(h, x) = (gh, x)$, $(g, x)^{-1} = (g^{-1}, gx)$.
- If \mathcal{G} is étale, then so is $\mathcal{G} \ltimes X$.

Actions of groupoids

- An **action** $\mathcal{G} \curvearrowright X$ of a groupoid \mathcal{G} on a space X consists of a cts map $X \xrightarrow{p} \mathcal{G}^0$ and a map $\theta: \mathcal{G} \times_{\mathcal{G}^0} X \rightarrow X$ such that:
 - $p(x)x = x$;
 - $g(hx) = (gh)x$;
 - $p(gx) = r(g)$.
- The **semidirect product** $\mathcal{G} \ltimes X$ has arrow space $\mathcal{G} \times_{\mathcal{G}^0} X$.
- The unit space is X .
- One has $x \xrightarrow{(g,x)} gx$.
- $(g, hx)(h, x) = (gh, x)$, $(g, x)^{-1} = (g^{-1}, gx)$.
- If \mathcal{G} is étale, then so is $\mathcal{G} \ltimes X$.

Actions of groupoids

- An **action** $\mathcal{G} \curvearrowright X$ of a groupoid \mathcal{G} on a space X consists of a cts map $X \xrightarrow{p} \mathcal{G}^0$ and a map $\theta: \mathcal{G} \times_{\mathcal{G}^0} X \rightarrow X$ such that:
 - $p(x)x = x$;
 - $g(hx) = (gh)x$;
 - $p(gx) = r(g)$.
- The **semidirect product** $\mathcal{G} \ltimes X$ has arrow space $\mathcal{G} \times_{\mathcal{G}^0} X$.
- The unit space is X .
- One has $x \xrightarrow{(g,x)} gx$.
- $(g, hx)(h, x) = (gh, x)$, $(g, x)^{-1} = (g^{-1}, gx)$.
- If \mathcal{G} is étale, then so is $\mathcal{G} \ltimes X$.

Actions of groupoids

- An **action** $\mathcal{G} \curvearrowright X$ of a groupoid \mathcal{G} on a space X consists of a cts map $X \xrightarrow{p} \mathcal{G}^0$ and a map $\theta: \mathcal{G} \times_{\mathcal{G}^0} X \rightarrow X$ such that:
 - $p(x)x = x$;
 - $g(hx) = (gh)x$;
 - $p(gx) = r(g)$.
- The **semidirect product** $\mathcal{G} \ltimes X$ has arrow space $\mathcal{G} \times_{\mathcal{G}^0} X$.
- The unit space is X .
- One has $x \xrightarrow{(g,x)} gx$.
- $(g, hx)(h, x) = (gh, x)$, $(g, x)^{-1} = (g^{-1}, gx)$.
- If \mathcal{G} is étale, then so is $\mathcal{G} \ltimes X$.

A Morita equivalence theorem

Theorem (Milan, BS)

Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a cts functor with \mathcal{G} locally compact and \mathcal{H} étale. Suppose that:

- the map $\mathcal{G} \rightarrow \mathcal{G}^0 \times_{\mathcal{H}^0} \mathcal{H} \times_{\mathcal{H}^0} \mathcal{G}^0$ given by

$$g \mapsto (d(g), \varphi(g), r(g))$$

is injective and closed;

- the map $\mathcal{H} \times_{\mathcal{H}^0} \mathcal{G} \rightarrow \mathcal{H} \times_{\mathcal{H}^0} \mathcal{G}^0$ given by

$$(h, g) \mapsto (h\varphi(g), d(g))$$

is open.

Then there is an action $\mathcal{H} \curvearrowright X$ such that \mathcal{G} is Morita equivalent to $\mathcal{H} \ltimes X$.

A Morita equivalence theorem

Theorem (Milan, BS)

Let $\varphi: \mathcal{G} \longrightarrow \mathcal{H}$ be a cts functor with \mathcal{G} locally compact and \mathcal{H} étale. Suppose that:

- the map $\mathcal{G} \longrightarrow \mathcal{G}^0 \times_{\mathcal{H}^0} \mathcal{H} \times_{\mathcal{H}^0} \mathcal{G}^0$ given by

$$g \longmapsto (d(g), \varphi(g), r(g))$$

is injective and closed;

- the map $\mathcal{H} \times_{\mathcal{H}^0} \mathcal{G} \longrightarrow \mathcal{H} \times_{\mathcal{H}^0} \mathcal{G}^0$ given by

$$(h, g) \longmapsto (h\varphi(g), d(g))$$

is open.

Then there is an *action* $\mathcal{H} \curvearrowright X$ such that \mathcal{G} is Morita equivalent to $\mathcal{H} \ltimes X$.

A Morita equivalence theorem

Theorem (Milan, BS)

Let $\varphi: \mathcal{G} \rightarrow \mathcal{H}$ be a cts functor with \mathcal{G} locally compact and \mathcal{H} étale. Suppose that:

- the map $\mathcal{G} \rightarrow \mathcal{G}^0 \times_{\mathcal{H}^0} \mathcal{H} \times_{\mathcal{H}^0} \mathcal{G}^0$ given by

$$g \mapsto (d(g), \varphi(g), r(g))$$

is injective and closed;

- the map $\mathcal{H} \times_{\mathcal{H}^0} \mathcal{G} \rightarrow \mathcal{H} \times_{\mathcal{H}^0} \mathcal{G}^0$ given by

$$(h, g) \mapsto (h\varphi(g), d(g))$$

is open.

Then there is an *action* $\mathcal{H} \curvearrowright X$ such that \mathcal{G} is Morita equivalent to $\mathcal{H} \ltimes X$.

A Morita equivalence theorem

Theorem (Milan, BS)

Let $\varphi: \mathcal{G} \longrightarrow \mathcal{H}$ be a cts functor with \mathcal{G} locally compact and \mathcal{H} étale. Suppose that:

- the map $\mathcal{G} \longrightarrow \mathcal{G}^0 \times_{\mathcal{H}^0} \mathcal{H} \times_{\mathcal{H}^0} \mathcal{G}^0$ given by

$$g \longmapsto (d(g), \varphi(g), r(g))$$

is injective and closed;

- the map $\mathcal{H} \times_{\mathcal{H}^0} \mathcal{G} \longrightarrow \mathcal{H} \times_{\mathcal{H}^0} \mathcal{G}^0$ given by

$$(h, g) \longmapsto (h\varphi(g), d(g))$$

is open.

Then there is an *action* $\mathcal{H} \curvearrowright X$ such that \mathcal{G} is *Morita equivalent* to $\mathcal{H} \ltimes X$.

Groupoids of germs as semidirect products

- To apply the previous situation, we need to identify groupoids of germs as semidirect products.

Theorem (Milan, BS)

- 1 *There is a bijection between special actions $T \curvearrowright X$ and actions $\mathcal{G}(T) \curvearrowright X$.*
- 2 *Moreover, the groupoid of germs $T \ltimes X$ is isomorphic to the semidirect product $\mathcal{G}(T) \ltimes X$.*

- For instance, if $\mathcal{G}(T)$ acts on X with $X \xrightarrow{p} \widehat{E(T)}$, then we put $X_e = p^{-1}(\widehat{E}_e)$.
- For $x \in X_{p(t)}$, we define $tx = [t, p(x)]x$.
- Conversely, if $T \curvearrowright X$, then $X \xrightarrow{p} \widehat{E(T)}$ is given by the "neighborhood filter:" $p(x) = \{e \mid x \in X_e\}$.
- One has $[s, \varphi]x = sx$.

Groupoids of germs as semidirect products

- To apply the previous situation, we need to identify groupoids of germs as semidirect products.

Theorem (Milan, BS)

- 1 *There is a bijection between special actions $T \curvearrowright X$ and actions $\mathcal{G}(T) \curvearrowright X$.*
- 2 *Moreover, the groupoid of germs $T \ltimes X$ is isomorphic to the semidirect product $\mathcal{G}(T) \ltimes X$.*

- For instance, if $\mathcal{G}(T)$ acts on X with $X \xrightarrow{p} \widehat{E(T)}$, then we put $X_e = p^{-1}(\widehat{E}_e)$.
- For $x \in X_{t^*t}$, we define $tx = [t, p(x)]x$.
- Conversely, if $T \curvearrowright X$, then $X \xrightarrow{p} \widehat{E(T)}$ is given by the "neighborhood filter:" $p(x) = \{e \mid x \in X_e\}$.
- One has $[s, \varphi]x = sx$.

Groupoids of germs as semidirect products

- To apply the previous situation, we need to identify groupoids of germs as semidirect products.

Theorem (Milan, BS)

- 1 *There is a bijection between special actions $T \curvearrowright X$ and actions $\mathcal{G}(T) \curvearrowright X$.*
- 2 *Moreover, the groupoid of germs $T \ltimes X$ is isomorphic to the semidirect product $\mathcal{G}(T) \ltimes X$.*

- For instance, if $\mathcal{G}(T)$ acts on X with $X \xrightarrow{p} \widehat{E(T)}$, then we put $X_e = p^{-1}(\widehat{E}_e)$.
- For $x \in X_{t^*t}$, we define $tx = [t, p(x)]x$.
- Conversely, if $T \curvearrowright X$, then $X \xrightarrow{p} \widehat{E(T)}$ is given by the "neighborhood filter:" $p(x) = \{e \mid x \in X_e\}$.
- One has $[s, \varphi]x = sx$.

Groupoids of germs as semidirect products

- To apply the previous situation, we need to identify groupoids of germs as semidirect products.

Theorem (Milan, BS)

- 1 *There is a bijection between special actions $T \curvearrowright X$ and actions $\mathcal{G}(T) \curvearrowright X$.*
- 2 *Moreover, the groupoid of germs $T \ltimes X$ is isomorphic to the semidirect product $\mathcal{G}(T) \ltimes X$.*

- For instance, if $\mathcal{G}(T)$ acts on X with $X \xrightarrow{p} \widehat{E(T)}$, then we put $X_e = p^{-1}(\widehat{E}_e)$.
- For $x \in X_{t^*t}$, we define $tx = [t, p(x)]x$.
- Conversely, if $T \curvearrowright X$, then $X \xrightarrow{p} \widehat{E(T)}$ is given by the “neighborhood filter:” $p(x) = \{e \mid x \in X_e\}$.
- One has $[s, \varphi]x = sx$.

Groupoids of germs as semidirect products

- To apply the previous situation, we need to identify groupoids of germs as semidirect products.

Theorem (Milan, BS)

- 1 *There is a bijection between special actions $T \curvearrowright X$ and actions $\mathcal{G}(T) \curvearrowright X$.*
- 2 *Moreover, the groupoid of germs $T \ltimes X$ is isomorphic to the semidirect product $\mathcal{G}(T) \ltimes X$.*

- For instance, if $\mathcal{G}(T)$ acts on X with $X \xrightarrow{p} \widehat{E(T)}$, then we put $X_e = p^{-1}(\widehat{E}_e)$.
- For $x \in X_{t^*t}$, we define $tx = [t, p(x)]x$.
- Conversely, if $T \curvearrowright X$, then $X \xrightarrow{p} \widehat{E(T)}$ is given by the “neighborhood filter:” $p(x) = \{e \mid x \in X_e\}$.
- One has $[s, \varphi]x = sx$.

Groupoids of germs as semidirect products

- To apply the previous situation, we need to identify groupoids of germs as semidirect products.

Theorem (Milan, BS)

- 1 *There is a bijection between special actions $T \curvearrowright X$ and actions $\mathcal{G}(T) \curvearrowright X$.*
- 2 *Moreover, the groupoid of germs $T \ltimes X$ is isomorphic to the semidirect product $\mathcal{G}(T) \ltimes X$.*

- For instance, if $\mathcal{G}(T)$ acts on X with $X \xrightarrow{p} \widehat{E(T)}$, then we put $X_e = p^{-1}(\widehat{E}_e)$.
- For $x \in X_{t^*t}$, we define $tx = [t, p(x)]x$.
- Conversely, if $T \curvearrowright X$, then $X \xrightarrow{p} \widehat{E(T)}$ is given by the “neighborhood filter:” $p(x) = \{e \mid x \in X_e\}$.
- One has $[s, \varphi]x = sx$.

Groupoids of germs as semidirect products

- To apply the previous situation, we need to identify groupoids of germs as semidirect products.

Theorem (Milan, BS)

- 1 *There is a bijection between special actions $T \curvearrowright X$ and actions $\mathcal{G}(T) \curvearrowright X$.*
- 2 *Moreover, the groupoid of germs $T \ltimes X$ is isomorphic to the semidirect product $\mathcal{G}(T) \ltimes X$.*

- For instance, if $\mathcal{G}(T)$ acts on X with $X \xrightarrow{p} \widehat{E(T)}$, then we put $X_e = p^{-1}(\widehat{E}_e)$.
- For $x \in X_{t^*t}$, we define $tx = [t, p(x)]x$.
- Conversely, if $T \curvearrowright X$, then $X \xrightarrow{p} \widehat{E(T)}$ is given by the “neighborhood filter:” $p(x) = \{e \mid x \in X_e\}$.
- One has $[s, \varphi]x = sx$.

The Khoshkam-Skandalis condition: revisited

- To prove our main result, one verifies that if $S \xrightarrow{\varphi} T$ is a **locally idempotent-pure morphism** satisfying the **KS condition**, then the induced map $\mathcal{G}(S) \longrightarrow \mathcal{G}(T)$ satisfies the conditions of our Morita theorem.
- Thus $\mathcal{G}(S)$ is Morita equivalent to $\mathcal{G}(T) \rtimes X \cong T \rtimes X$.
- Therefore, $C^*(S)$ is strongly Morita equivalent to $C_0(X) \rtimes T$.

The Khoshkam-Skandalis condition: revisited

- To prove our main result, one verifies that if $S \xrightarrow{\varphi} T$ is a **locally idempotent-pure morphism** satisfying the **KS condition**, then the induced map $\mathcal{G}(S) \longrightarrow \mathcal{G}(T)$ satisfies the conditions of our Morita theorem.
- Thus $\mathcal{G}(S)$ is Morita equivalent to $\mathcal{G}(T) \ltimes X \cong T \ltimes X$.
- Therefore, $C^*(S)$ is strongly Morita equivalent to $C_0(X) \ltimes T$.

The Khoshkam-Skandalis condition: revisited

- To prove our main result, one verifies that if $S \xrightarrow{\varphi} T$ is a **locally idempotent-pure morphism** satisfying the **KS condition**, then the induced map $\mathcal{G}(S) \longrightarrow \mathcal{G}(T)$ satisfies the conditions of our Morita theorem.
- Thus $\mathcal{G}(S)$ is Morita equivalent to $\mathcal{G}(T) \rtimes X \cong T \rtimes X$.
- Therefore, $C^*(S)$ is strongly Morita equivalent to $C_0(X) \rtimes T$.

Morita equivalence

- Let S be an inverse semigroup.
- Let $S\text{-Set}$ be the category of sets X equipped with an action $S \times X \longrightarrow X$ by **total** functions such that $SX = X$.
- Talwar defined two inverse semigroups S and T to be **Morita equivalent** if $S\text{-Set}$ is equivalent to $T\text{-Set}$.
- For example, if S is an inverse monoid, then S and T are Morita equivalent iff there is an idempotent e with $T = TeT$ and $S \cong eTe$.
- An inverse semigroup is F -inverse iff it is Morita equivalent to a semidirect product of a group and a semilattice.
- Lawson calls S an enlargement of a subsemigroup T if $S = STS$ and $T = TST$.
- In this case S and T are Morita equivalent.
- Two inverse semigroups are Morita equivalent iff they have a common von Neumann regular enlargement (Lawson).

Morita equivalence

- Let S be an inverse semigroup.
- Let $S\text{-Set}$ be the category of sets X equipped with an action $S \times X \longrightarrow X$ by **total** functions such that $SX = X$.
- Talwar defined two inverse semigroups S and T to be **Morita equivalent** if $S\text{-Set}$ is equivalent to $T\text{-Set}$.
- For example, if S is an inverse monoid, then S and T are Morita equivalent iff there is an idempotent e with $T = TeT$ and $S \cong eTe$.
- An inverse semigroup is **F -inverse** iff it is Morita equivalent to a **semidirect product** of a group and a semilattice.
- Lawson calls S an **enlargement** of a subsemigroup T if $S = STS$ and $T = TST$.
- In this case S and T are Morita equivalent.
- Two inverse semigroups are Morita equivalent iff they have a common von Neumann regular enlargement (Lawson).

Morita equivalence

- Let S be an inverse semigroup.
- Let $S\text{-Set}$ be the category of sets X equipped with an action $S \times X \longrightarrow X$ by **total** functions such that $SX = X$.
- Talwar defined two inverse semigroups S and T to be **Morita equivalent** if $S\text{-Set}$ is equivalent to $T\text{-Set}$.
- For example, if S is an inverse monoid, then S and T are Morita equivalent iff there is an idempotent e with $T = TeT$ and $S \cong eTe$.
- An inverse semigroup is **F -inverse** iff it is Morita equivalent to a **semidirect product** of a group and a semilattice.
- Lawson calls S an **enlargement** of a subsemigroup T if $S = STS$ and $T = TST$.
- In this case S and T are Morita equivalent.
- Two inverse semigroups are Morita equivalent iff they have a common von Neumann regular enlargement (Lawson).

Morita equivalence

- Let S be an inverse semigroup.
- Let $S\text{-Set}$ be the category of sets X equipped with an action $S \times X \longrightarrow X$ by **total** functions such that $SX = X$.
- Talwar defined two inverse semigroups S and T to be **Morita equivalent** if $S\text{-Set}$ is equivalent to $T\text{-Set}$.
- For example, if S is an inverse monoid, then S and T are Morita equivalent iff there is an idempotent e with $T = TeT$ and $S \cong eTe$.
- An inverse semigroup is **F -inverse** iff it is Morita equivalent to a **semidirect product** of a group and a semilattice.
- Lawson calls S an **enlargement** of a subsemigroup T if $S = STS$ and $T = TST$.
- In this case S and T are Morita equivalent.
- Two inverse semigroups are Morita equivalent iff they have a common von Neumann regular enlargement (Lawson).

Morita equivalence

- Let S be an inverse semigroup.
- Let $S\text{-Set}$ be the category of sets X equipped with an action $S \times X \longrightarrow X$ by **total** functions such that $SX = X$.
- Talwar defined two inverse semigroups S and T to be **Morita equivalent** if $S\text{-Set}$ is equivalent to $T\text{-Set}$.
- For example, if S is an inverse monoid, then S and T are Morita equivalent iff there is an idempotent e with $T = TeT$ and $S \cong eTe$.
- An inverse semigroup is **F -inverse** iff it is Morita equivalent to a **semidirect product** of a group and a semilattice.
- Lawson calls S an **enlargement** of a subsemigroup T if $S = STS$ and $T = TST$.
- In this case S and T are Morita equivalent.
- Two inverse semigroups are Morita equivalent iff they have a common von Neumann regular enlargement (Lawson).

Morita equivalence

- Let S be an inverse semigroup.
- Let $S\text{-Set}$ be the category of sets X equipped with an action $S \times X \longrightarrow X$ by **total** functions such that $SX = X$.
- Talwar defined two inverse semigroups S and T to be **Morita equivalent** if $S\text{-Set}$ is equivalent to $T\text{-Set}$.
- For example, if S is an inverse monoid, then S and T are Morita equivalent iff there is an idempotent e with $T = TeT$ and $S \cong eTe$.
- An inverse semigroup is **F -inverse** iff it is Morita equivalent to a **semidirect product** of a group and a semilattice.
- Lawson calls S an **enlargement** of a subsemigroup T if $S = STS$ and $T = TST$.
- In this case S and T are Morita equivalent.
- Two inverse semigroups are Morita equivalent iff they have a common von Neumann regular enlargement (Lawson).

Morita equivalence

- Let S be an inverse semigroup.
- Let $S\text{-Set}$ be the category of sets X equipped with an action $S \times X \longrightarrow X$ by **total** functions such that $SX = X$.
- Talwar defined two inverse semigroups S and T to be **Morita equivalent** if $S\text{-Set}$ is equivalent to $T\text{-Set}$.
- For example, if S is an inverse monoid, then S and T are Morita equivalent iff there is an idempotent e with $T = TeT$ and $S \cong eTe$.
- An inverse semigroup is **F -inverse** iff it is Morita equivalent to a **semidirect product** of a group and a semilattice.
- Lawson calls S an **enlargement** of a subsemigroup T if $S = STS$ and $T = TST$.
- In this case S and T are Morita equivalent.
- Two inverse semigroups are Morita equivalent iff they have a common von Neumann regular enlargement (Lawson).

Morita equivalence

- Let S be an inverse semigroup.
- Let $S\text{-Set}$ be the category of sets X equipped with an action $S \times X \longrightarrow X$ by **total** functions such that $SX = X$.
- Talwar defined two inverse semigroups S and T to be **Morita equivalent** if $S\text{-Set}$ is equivalent to $T\text{-Set}$.
- For example, if S is an inverse monoid, then S and T are Morita equivalent iff there is an idempotent e with $T = TeT$ and $S \cong eTe$.
- An inverse semigroup is **F -inverse** iff it is Morita equivalent to a **semidirect product** of a group and a semilattice.
- Lawson calls S an **enlargement** of a subsemigroup T if $S = STS$ and $T = TST$.
- In this case S and T are Morita equivalent.
- Two inverse semigroups are Morita equivalent iff they have a common von Neumann regular enlargement (Lawson).

Strong Morita equivalence

- I defined inverse semigroups S and T to be **strongly Morita equivalent** if there is an **equivalence bimodule** for S and T .
- By definition, this consists of a set X , which is an (S, T) -biset equipped with surjective “inner products”

$$\langle -, - \rangle: X \times X \longrightarrow S, \text{ and } [-, -]: X \times X \longrightarrow T$$

such that the following axioms hold, where $x, y, z \in X$, $s \in S$, and $t \in T$:

- $\langle sx, y \rangle = s \langle x, y \rangle$ $[x, yt] = [x, y]t$
 - $\langle y, x \rangle = \langle x, y \rangle^*$ $[x, y] = [y, x]^*$
 - $\langle x, x \rangle x = x$ $x[x, x] = x$
 - $\langle x, y \rangle z = x[y, z]$.
- For instance, if S is an enlargement of T , then $X = ST$ is an equivalence bimodule.

Strong Morita equivalence

- I defined inverse semigroups S and T to be **strongly Morita equivalent** if there is an **equivalence bimodule** for S and T .
- By definition, this consists of a set X , which is an (S, T) -biset equipped with surjective “inner products”

$$\langle -, - \rangle: X \times X \longrightarrow S, \text{ and } [-, -]: X \times X \longrightarrow T$$

such that the following axioms hold, where $x, y, z \in X$, $s \in S$, and $t \in T$:

- $\langle sx, y \rangle = s \langle x, y \rangle$ $[x, yt] = [x, y]t$
- $\langle y, x \rangle = \langle x, y \rangle^*$ $[x, y] = [y, x]^*$
- $\langle x, x \rangle x = x$ $x[x, x] = x$
- $\langle x, y \rangle z = x[y, z]$.

- For instance, if S is an enlargement of T , then $X = ST$ is an equivalence bimodule.

Strong Morita equivalence

- I defined inverse semigroups S and T to be **strongly Morita equivalent** if there is an **equivalence bimodule** for S and T .
- By definition, this consists of a set X , which is an (S, T) -biset equipped with surjective “inner products”

$$\langle -, - \rangle: X \times X \longrightarrow S, \text{ and } [-, -]: X \times X \longrightarrow T$$

such that the following axioms hold, where $x, y, z \in X$, $s \in S$, and $t \in T$:

- $\langle sx, y \rangle = s \langle x, y \rangle$ $[x, yt] = [x, y]t$
 - $\langle y, x \rangle = \langle x, y \rangle^*$ $[x, y] = [y, x]^*$
 - $\langle x, x \rangle x = x$ $x[x, x] = x$
 - $\langle x, y \rangle z = x[y, z]$.
- For instance, if S is an enlargement of T , then $X = ST$ is an equivalence bimodule.

Strong Morita equivalence and groupoids

Theorem (BS)

If S and T are strongly Morita equivalent, then $\mathcal{G}(S)$ and $\mathcal{G}(T)$ are Morita equivalent.

- Every F -inverse semigroup S has an enlargement $E \rtimes G(S)$.
- Thus $C^*(S)$ is strongly Morita equivalent to $C_0(\widehat{E}) \rtimes G(S)$.
- This yields an alternative proof to that of Khoshkam and Skandalis.
- It is easy to see that strong Morita equivalence implies Morita equivalence.
- I had conjectured the converse is false.

Theorem (Funk, Lawson, BS)

Morita equivalence and strong Morita equivalence are the same.

- The proof uses topos theoretic ideas.

Strong Morita equivalence and groupoids

Theorem (BS)

If S and T are strongly Morita equivalent, then $\mathcal{G}(S)$ and $\mathcal{G}(T)$ are Morita equivalent.

- Every F -inverse semigroup S has an enlargement $E \rtimes G(S)$.
- Thus $C^*(S)$ is strongly Morita equivalent to $C_0(\widehat{E}) \rtimes G(S)$.
- This yields an alternative proof to that of Khoshkam and Skandalis.
- It is easy to see that strong Morita equivalence implies Morita equivalence.
- I had conjectured the converse is false.

Theorem (Funk, Lawson, BS)

Morita equivalence and strong Morita equivalence are the same.

- The proof uses topos theoretic ideas.

Strong Morita equivalence and groupoids

Theorem (BS)

If S and T are strongly Morita equivalent, then $\mathcal{G}(S)$ and $\mathcal{G}(T)$ are Morita equivalent.

- Every F -inverse semigroup S has an enlargement $E \rtimes G(S)$.
- Thus $C^*(S)$ is strongly Morita equivalent to $C_0(\widehat{E}) \rtimes G(S)$.
- This yields an alternative proof to that of Khoshkam and Skandalis.
- It is easy to see that strong Morita equivalence implies Morita equivalence.
- I had conjectured the converse is false.

Theorem (Funk, Lawson, BS)

Morita equivalence and strong Morita equivalence are the same.

- The proof uses topos theoretic ideas.

Strong Morita equivalence and groupoids

Theorem (BS)

If S and T are strongly Morita equivalent, then $\mathcal{G}(S)$ and $\mathcal{G}(T)$ are Morita equivalent.

- Every F -inverse semigroup S has an enlargement $E \rtimes G(S)$.
- Thus $C^*(S)$ is strongly Morita equivalent to $C_0(\widehat{E}) \rtimes G(S)$.
- This yields an alternative proof to that of Khoshkam and Skandalis.
- It is easy to see that strong Morita equivalence implies Morita equivalence.
- I had conjectured the converse is false.

Theorem (Funk, Lawson, BS)

Morita equivalence and strong Morita equivalence are the same.

- The proof uses topos theoretic ideas.

Strong Morita equivalence and groupoids

Theorem (BS)

If S and T are strongly Morita equivalent, then $\mathcal{G}(S)$ and $\mathcal{G}(T)$ are Morita equivalent.

- Every F -inverse semigroup S has an enlargement $E \rtimes G(S)$.
- Thus $C^*(S)$ is strongly Morita equivalent to $C_0(\widehat{E}) \rtimes G(S)$.
- This yields an alternative proof to that of Khoshkam and Skandalis.
- It is easy to see that strong Morita equivalence implies Morita equivalence.
- I had conjectured the converse is false.

Theorem (Funk, Lawson, BS)

Morita equivalence and strong Morita equivalence are the same.

- The proof uses topos theoretic ideas.

Strong Morita equivalence and groupoids

Theorem (BS)

If S and T are strongly Morita equivalent, then $\mathcal{G}(S)$ and $\mathcal{G}(T)$ are Morita equivalent.

- Every F -inverse semigroup S has an enlargement $E \rtimes G(S)$.
- Thus $C^*(S)$ is strongly Morita equivalent to $C_0(\widehat{E}) \rtimes G(S)$.
- This yields an alternative proof to that of Khoshkam and Skandalis.
- It is easy to see that strong Morita equivalence implies Morita equivalence.
- I had conjectured the converse is false.

Theorem (Funk, Lawson, BS)

Morita equivalence and strong Morita equivalence are the same.

- The proof uses topos theoretic ideas.

Strong Morita equivalence and groupoids

Theorem (BS)

If S and T are strongly Morita equivalent, then $\mathcal{G}(S)$ and $\mathcal{G}(T)$ are Morita equivalent.

- Every F -inverse semigroup S has an enlargement $E \rtimes G(S)$.
- Thus $C^*(S)$ is strongly Morita equivalent to $C_0(\widehat{E}) \rtimes G(S)$.
- This yields an alternative proof to that of Khoshkam and Skandalis.
- It is easy to see that strong Morita equivalence implies Morita equivalence.
- I had conjectured the converse is false.

Theorem (Funk, Lawson, BS)

Morita equivalence and strong Morita equivalence are the same.

- The proof uses topos theoretic ideas.

Strong Morita equivalence and groupoids

Theorem (BS)

If S and T are strongly Morita equivalent, then $\mathcal{G}(S)$ and $\mathcal{G}(T)$ are Morita equivalent.

- Every F -inverse semigroup S has an enlargement $E \rtimes G(S)$.
- Thus $C^*(S)$ is strongly Morita equivalent to $C_0(\widehat{E}) \rtimes G(S)$.
- This yields an alternative proof to that of Khoshkam and Skandalis.
- It is easy to see that strong Morita equivalence implies Morita equivalence.
- I had conjectured the converse is false.

Theorem (Funk, Lawson, BS)

Morita equivalence and strong Morita equivalence are the same.

- The proof uses topos theoretic ideas.

Thanks for your attention!

Je vous remercie de votre attention!

Obrigado pela sua atenção!