

# Koszul algebras associated to zonotopes and $CAT(0)$ cube complexes

Stuart Margolis, Bar-Ilan University

Franco Saliola, Université du Québec à Montréal

**Benjamin Steinberg**, City College of New York



December 7, 2013  
Winter CMS Meeting

# Outline

Hyperplane arrangements and cell complexes

Left regular bands

Representation theory

# Goal

- We introduce some Koszul algebras associated to certain regular cell complexes.

# Goal

- We introduce some Koszul algebras associated to certain regular cell complexes.
- One can use algebraic topology to compute minimal projective resolutions of the simple modules.

# Goal

- We introduce some Koszul algebras associated to certain regular cell complexes.
- One can use algebraic topology to compute minimal projective resolutions of the simple modules.
- These algebras are dual to incidence algebras of certain Cohen-Macaulay posets.

# Goal


- We introduce some Koszul algebras associated to certain regular cell complexes.
- One can use algebraic topology to compute minimal projective resolutions of the simple modules.
- These algebras are dual to incidence algebras of certain Cohen-Macaulay posets.
- The original motivation for studying these algebras came from analyzing Markov chains.

# Goal

- We introduce some Koszul algebras associated to certain regular cell complexes.
- One can use algebraic topology to compute minimal projective resolutions of the simple modules.
- These algebras are dual to incidence algebras of certain Cohen-Macaulay posets.
- The original motivation for studying these algebras came from analyzing Markov chains.
- All these algebras are semigroup algebras.

# Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:





## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



the origin

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**rays** emanating from the origin

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**rays** emanating from the origin

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**rays** emanating from the origin

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**rays** emanating from the origin

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**rays** emanating from the origin

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**rays** emanating from the origin

## Faces of a hyperplane arrangement

A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:




**chambers** cut out by the hyperplanes



## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**chambers** cut out by the hyperplanes

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**chambers** cut out by the hyperplanes

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**chambers** cut out by the hyperplanes

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**chambers** cut out by the hyperplanes

## Faces of a hyperplane arrangement


A set of hyperplanes partitions  $\mathbb{R}^n$  into *faces*:



**chambers** cut out by the hyperplanes


## Product of faces (LRB structure)

$xy := \left\{ \begin{array}{l} \text{the face first encountered after a small} \\ \text{movement along a line from } x \text{ toward } y \end{array} \right.$




## Product of faces (LRB structure)

$xy := \left\{ \begin{array}{l} \text{the face first encountered after a small} \\ \text{movement along a line from } x \text{ toward } y \end{array} \right.$



## Product of faces (LRB structure)


$xy := \left\{ \begin{array}{l} \text{the face first encountered after a small} \\ \text{movement along a line from } x \text{ toward } y \end{array} \right.$





## Product of faces (LRB structure)

$xy := \left\{ \begin{array}{l} \text{the face first encountered after a small} \\ \text{movement along a line from } x \text{ toward } y \end{array} \right.$



## A one-dimensional arrangement

- Let's give a more formal definition of hyperplane face semigroups.

## A one-dimensional arrangement

- Let's give a more formal definition of hyperplane face semigroups.
- We begin with the case of a one-dimensional central arrangement.



## A one-dimensional arrangement

- Let's give a more formal definition of hyperplane face semigroups.
- We begin with the case of a one-dimensional central arrangement.



- There are three regions  $+$ ,  $-$ ,  $0$ .

## A one-dimensional arrangement

- Let's give a more formal definition of hyperplane face semigroups.
- We begin with the case of a one-dimensional central arrangement.



- There are three regions  $+$ ,  $-$ ,  $0$ .

$\circ$	$0$	$+$	$-$
$0$	$0$	$+$	$-$
$+$	$+$	$+$	$+$
$-$	$-$	$-$	$-$

## A one-dimensional arrangement

- Let's give a more formal definition of hyperplane face semigroups.
- We begin with the case of a one-dimensional central arrangement.



- There are three regions  $+$ ,  $-$ ,  $0$ .

$\circ$	$0$	$+$	$-$
$0$	$0$	$+$	$-$
$+$	$+$	$+$	$+$
$-$	$-$	$-$	$-$

- One easily checks  $\{+, -, 0\}$  satisfies the identities  $x^2 = x$  and  $xyx = xy$ .

## Hyperplane face semigroups

- Let  $\mathcal{A} = \{H_1, \dots, H_d\}$  be a hyperplane arrangement in a Euclidean space  $V$ .

## Hyperplane face semigroups

- Let  $\mathcal{A} = \{H_1, \dots, H_d\}$  be a hyperplane arrangement in a Euclidean space  $V$ .
- Assume  $H_i$  is given by  $f_i = c_i$  with  $f_i \in V^*$ ,  $c_i \in \mathbb{R}$ .



## Hyperplane face semigroups

- Let  $\mathcal{A} = \{H_1, \dots, H_d\}$  be a hyperplane arrangement in a Euclidean space  $V$ .
- Assume  $H_i$  is given by  $f_i = c_i$  with  $f_i \in V^*$ ,  $c_i \in \mathbb{R}$ .
- Define  $\sigma: V \rightarrow \{+, -, 0\}^d$  by  $\sigma(v) = (\sigma_i)_{i=1}^d$  where

$$\sigma_i = \operatorname{sgn}(f_i(v) - c_i).$$

## Hyperplane face semigroups

- Let  $\mathcal{A} = \{H_1, \dots, H_d\}$  be a hyperplane arrangement in a Euclidean space  $V$ .
- Assume  $H_i$  is given by  $f_i = c_i$  with  $f_i \in V^*$ ,  $c_i \in \mathbb{R}$ .
- Define  $\sigma: V \rightarrow \{+, -, 0\}^d$  by  $\sigma(v) = (\sigma_i)_{i=1}^d$  where

$$\sigma_i = \operatorname{sgn}(f_i(v) - c_i).$$

- $\sigma(v) = \sigma(w)$  iff they are in the same face.

## Hyperplane face semigroups

- Let  $\mathcal{A} = \{H_1, \dots, H_d\}$  be a hyperplane arrangement in a Euclidean space  $V$ .
- Assume  $H_i$  is given by  $f_i = c_i$  with  $f_i \in V^*$ ,  $c_i \in \mathbb{R}$ .
- Define  $\sigma: V \rightarrow \{+, -, 0\}^d$  by  $\sigma(v) = (\sigma_i)_{i=1}^d$  where

$$\sigma_i = \operatorname{sgn}(f_i(v) - c_i).$$

- $\sigma(v) = \sigma(w)$  iff they are in the same face.
- $\mathcal{F}(\mathcal{A}) = \sigma(V)$  is a subsemigroup called the **face semigroup of  $\mathcal{A}$** .

## Hyperplane face semigroups

- Let  $\mathcal{A} = \{H_1, \dots, H_d\}$  be a hyperplane arrangement in a Euclidean space  $V$ .
- Assume  $H_i$  is given by  $f_i = c_i$  with  $f_i \in V^*$ ,  $c_i \in \mathbb{R}$ .
- Define  $\sigma: V \rightarrow \{+, -, 0\}^d$  by  $\sigma(v) = (\sigma_i)_{i=1}^d$  where

$$\sigma_i = \operatorname{sgn}(f_i(v) - c_i).$$

- $\sigma(v) = \sigma(w)$  iff they are in the same face.
- $\mathcal{F}(\mathcal{A}) = \sigma(V)$  is a subsemigroup called the **face semigroup of  $\mathcal{A}$** .
- It is a submonoid iff  $\mathcal{A}$  is **central**, i.e.,  $\bigcap H_i \neq \emptyset$ .

## Hyperplane face semigroups

- Let  $\mathcal{A} = \{H_1, \dots, H_d\}$  be a hyperplane arrangement in a Euclidean space  $V$ .
- Assume  $H_i$  is given by  $f_i = c_i$  with  $f_i \in V^*$ ,  $c_i \in \mathbb{R}$ .
- Define  $\sigma: V \rightarrow \{+, -, 0\}^d$  by  $\sigma(v) = (\sigma_i)_{i=1}^d$  where

$$\sigma_i = \operatorname{sgn}(f_i(v) - c_i).$$

- $\sigma(v) = \sigma(w)$  iff they are in the same face.
- $\mathcal{F}(\mathcal{A}) = \sigma(V)$  is a subsemigroup called the **face semigroup of  $\mathcal{A}$** .
- It is a submonoid iff  $\mathcal{A}$  is **central**, i.e.,  $\bigcap H_i \neq \emptyset$ .
- $\sigma(v) \circ \sigma(w) = \sigma(z)$  where  $z = (1 - t)v + tw$  with  $t > 0$  sufficiently small.


## Hyperplane face semigroups

- Let  $\mathcal{A} = \{H_1, \dots, H_d\}$  be a hyperplane arrangement in a Euclidean space  $V$ .
- Assume  $H_i$  is given by  $f_i = c_i$  with  $f_i \in V^*$ ,  $c_i \in \mathbb{R}$ .
- Define  $\sigma: V \rightarrow \{+, -, 0\}^d$  by  $\sigma(v) = (\sigma_i)_{i=1}^d$  where

$$\sigma_i = \operatorname{sgn}(f_i(v) - c_i).$$


- $\sigma(v) = \sigma(w)$  iff they are in the same face.
- $\mathcal{F}(\mathcal{A}) = \sigma(V)$  is a subsemigroup called the **face semigroup of  $\mathcal{A}$** .
- It is a submonoid iff  $\mathcal{A}$  is **central**, i.e.,  $\bigcap H_i \neq \emptyset$ .
- $\sigma(v) \circ \sigma(w) = \sigma(z)$  where  $z = (1 - t)v + tw$  with  $t > 0$  sufficiently small.
- This whole setup generalizes to **oriented matroids**.

## Examples



**Figure :** The sign sequences of the faces of the hyperplane arrangement in  $\mathbb{R}^2$  consisting of three distinct lines.

## Examples



**Figure :** The sign sequences of the faces of the hyperplane arrangement in  $\mathbb{R}^2$  consisting of three distinct lines.

### Example (The coordinate arrangement)

The face monoid of  $\{x_i = 0 \mid i = 1, \dots, n\}$  in  $\mathbb{R}^n$  is  $\{0, +, -\}^n$ . The chambers are the orthants.



## Complex hyperplane arrangements


- This framework was generalized to complex hyperplane arrangements by Björner and Ziegler.

## Complex hyperplane arrangements

- This framework was generalized to complex hyperplane arrangements by Björner and Ziegler.
- We need only consider the one-dimensional complex arrangement consisting of the origin in  $\mathbb{C}$ .


## Complex hyperplane arrangements

- This framework was generalized to complex hyperplane arrangements by Björner and Ziegler.
- We need only consider the one-dimensional complex arrangement consisting of the origin in  $\mathbb{C}$ .



## Complex hyperplane arrangements

- This framework was generalized to complex hyperplane arrangements by Björner and Ziegler.
- We need only consider the one-dimensional complex arrangement consisting of the origin in  $\mathbb{C}$ .



- $\{0, +, -\}$  is a submonoid and  $\pm i \circ x = \pm i$ ,  $y \circ \pm i = \pm i$  if  $y \in \{0, +, -\}$ .

## Zonotopes: the dual picture

- Let  $\mathcal{A}$  be a central hyperplane arrangement in  $V$ .

## Zonotopes: the dual picture

- Let  $\mathcal{A}$  be a central hyperplane arrangement in  $V$ .
- Assume  $\mathcal{A}$  is given by forms  $f_i = 0$  for  $i = 1, \dots, d$ .

## Zonotopes: the dual picture

- Let  $\mathcal{A}$  be a central hyperplane arrangement in  $V$ .
- Assume  $\mathcal{A}$  is given by forms  $f_i = 0$  for  $i = 1, \dots, d$ .
- The **zonotope** dual to  $\mathcal{A}$  is the Minkowski sum

$$\mathcal{Z}(\mathcal{A}) = [-f_1, f_1] + \cdots + [-f_d, f_d] = \left\{ \sum_{i=1}^d t_i f_i \mid -1 \leq t_i \leq 1 \right\}.$$

## Zonotopes: the dual picture

- Let  $\mathcal{A}$  be a central hyperplane arrangement in  $V$ .
- Assume  $\mathcal{A}$  is given by forms  $f_i = 0$  for  $i = 1, \dots, d$ .
- The **zonotope** dual to  $\mathcal{A}$  is the Minkowski sum

$$\mathcal{Z}(\mathcal{A}) = [-f_1, f_1] + \dots + [-f_d, f_d] = \left\{ \sum_{i=1}^d t_i f_i \mid -1 \leq t_i \leq 1 \right\}.$$

- The zonotope of the coordinate arrangement  $\{x_i = 0\}$  in  $\mathbb{R}^n$  is the **hypercube**  $[-1, 1]^n$ .



## Zonotopes: the dual picture

- Let  $\mathcal{A}$  be a central hyperplane arrangement in  $V$ .
- Assume  $\mathcal{A}$  is given by forms  $f_i = 0$  for  $i = 1, \dots, d$ .
- The **zonotope** dual to  $\mathcal{A}$  is the Minkowski sum

$$\mathcal{Z}(\mathcal{A}) = [-f_1, f_1] + \dots + [-f_d, f_d] = \left\{ \sum_{i=1}^d t_i f_i \mid -1 \leq t_i \leq 1 \right\}.$$

- The zonotope of the coordinate arrangement  $\{x_i = 0\}$  in  $\mathbb{R}^n$  is the **hypercube**  $[-1, 1]^n$ .
- The zonotope of the braid arrangement  $\{x_i = x_j\}$  in  $\mathbb{R}^n$  is the **permutohedron**.

## Zonotopes: the dual picture

- Let  $\mathcal{A}$  be a central hyperplane arrangement in  $V$ .
- Assume  $\mathcal{A}$  is given by forms  $f_i = 0$  for  $i = 1, \dots, d$ .
- The **zonotope** dual to  $\mathcal{A}$  is the Minkowski sum

$$\mathcal{Z}(\mathcal{A}) = [-f_1, f_1] + \dots + [-f_d, f_d] = \left\{ \sum_{i=1}^d t_i f_i \mid -1 \leq t_i \leq 1 \right\}.$$

- The zonotope of the coordinate arrangement  $\{x_i = 0\}$  in  $\mathbb{R}^n$  is the **hypercube**  $[-1, 1]^n$ .
- The zonotope of the braid arrangement  $\{x_i = x_j\}$  in  $\mathbb{R}^n$  is the **permutohedron**.
- The faces of  $\mathcal{Z}(\mathcal{A})$  are in order-reversing bijection with the faces of  $\mathcal{A}$ .

## Zonotopes: the dual picture

- Let  $\mathcal{A}$  be a central hyperplane arrangement in  $V$ .
- Assume  $\mathcal{A}$  is given by forms  $f_i = 0$  for  $i = 1, \dots, d$ .
- The **zonotope** dual to  $\mathcal{A}$  is the Minkowski sum

$$\mathcal{Z}(\mathcal{A}) = [-f_1, f_1] + \cdots + [-f_d, f_d] = \left\{ \sum_{i=1}^d t_i f_i \mid -1 \leq t_i \leq 1 \right\}.$$

- The zonotope of the coordinate arrangement  $\{x_i = 0\}$  in  $\mathbb{R}^n$  is the **hypercube**  $[-1, 1]^n$ .
- The zonotope of the braid arrangement  $\{x_i = x_j\}$  in  $\mathbb{R}^n$  is the **permutohedron**.
- The faces of  $\mathcal{Z}(\mathcal{A})$  are in order-reversing bijection with the faces of  $\mathcal{A}$ .
- Similarly, complex arrangements, affine arrangements and oriented matroids have dual regular cell complexes.

# Permutohedron




Figure : Permutohedron for  $S_4$  by Mike Zabrocki

# Zonohedra




Figure : 3-dimensional zontopes (zonohedra)

# An oriented matroid




Figure : An oriented matroid corresponding to a zonotopal tiling

# Advantages of the dual approach

- Working with the zonotope has a number of advantages.

## Advantages of the dual approach

- Working with the zonotope has a number of advantages.
- The face semigroup acts by cellular maps on the zonotope.



## Advantages of the dual approach

- Working with the zonotope has a number of advantages.
- The face semigroup acts by cellular maps on the zonotope.
- The augmented cellular chain complex of  $\mathcal{Z}(\mathcal{A})$  becomes a minimal projective resolution of the trivial  $\mathbb{k}\mathcal{F}(\mathcal{A})$ -module.

## Advantages of the dual approach

- Working with the zonotope has a number of advantages.
- The face semigroup acts by cellular maps on the zonotope.
- The augmented cellular chain complex of  $\mathcal{Z}(\mathcal{A})$  becomes a minimal projective resolution of the trivial  $\mathbb{k}\mathcal{F}(\mathcal{A})$ -module.
- Projective resolutions of the other simple modules arise by contracting the hyperplane arrangement and using zonotopes.

## Advantages of the dual approach

- Working with the zonotope has a number of advantages.
- The face semigroup acts by cellular maps on the zonotope.
- The augmented cellular chain complex of  $\mathcal{Z}(\mathcal{A})$  becomes a minimal projective resolution of the trivial  $\mathbb{k}\mathcal{F}(\mathcal{A})$ -module.
- Projective resolutions of the other simple modules arise by contracting the hyperplane arrangement and using zonotopes.
- Every subcomplex of the zonotope is a subsemigroup of  $\mathcal{F}(\mathcal{A})$ .

## CAT(0) cube complexes

- A **cube complex** is a regular cell complex obtained by gluing together cubes along their faces.

## CAT(0) cube complexes

- A **cube complex** is a regular cell complex obtained by gluing together cubes along their faces.
- A simply connected cube complex is **CAT(0)** if the link of each vertex is a flag complex (Gromov).

## CAT(0) cube complexes

- A **cube complex** is a regular cell complex obtained by gluing together cubes along their faces.
- A simply connected cube complex is **CAT(0)** if the link of each vertex is a flag complex (Gromov).
- CAT(0) cube complexes have played a major role in geometric group theory.

## CAT(0) cube complexes

- A **cube complex** is a regular cell complex obtained by gluing together cubes along their faces.
- A simply connected cube complex is **CAT(0)** if the link of each vertex is a flag complex (Gromov).
- CAT(0) cube complexes have played a major role in geometric group theory.
- They were used by Agol and Wise to solve Thurston's virtual fibering and virtual Haken conjectures.

## CAT(0) cube complexes

- A **cube complex** is a regular cell complex obtained by gluing together cubes along their faces.
- A simply connected cube complex is **CAT(0)** if the link of each vertex is a flag complex (Gromov).
- CAT(0) cube complexes have played a major role in geometric group theory.
- They were used by Agol and Wise to solve Thurston's virtual fibering and virtual Haken conjectures.
- They also arise in combinatorics and tropical geometry.



## CAT(0) cube complexes

- A **cube complex** is a regular cell complex obtained by gluing together cubes along their faces.
- A simply connected cube complex is **CAT(0)** if the link of each vertex is a flag complex (Gromov).
- CAT(0) cube complexes have played a major role in geometric group theory.
- They were used by Agol and Wise to solve Thurston's virtual fibering and virtual Haken conjectures.
- They also arise in combinatorics and tropical geometry.
- Billera, Holmes and Vogtmann introduced a space of phylogenetic trees, which is a CAT(0) cube complex.

## CAT(0) cube complexes

- A **cube complex** is a regular cell complex obtained by gluing together cubes along their faces.
- A simply connected cube complex is **CAT(0)** if the link of each vertex is a flag complex (Gromov).
- CAT(0) cube complexes have played a major role in geometric group theory.
- They were used by Agol and Wise to solve Thurston's virtual fibering and virtual Haken conjectures.
- They also arise in combinatorics and tropical geometry.
- Billera, Holmes and Vogtmann introduced a space of phylogenetic trees, which is a CAT(0) cube complex.
- Speyer showed that the space of phylogenetic trees is the tropical analog of the Grassmannian.

# A CAT(0) square complex




Figure : A CAT(0) square complex

## CAT(0) cube complexes (ctd.)

- Finite CAT(0) cube complexes are subcomplexes of hypercubes.

## CAT(0) cube complexes (ctd.)

- Finite CAT(0) cube complexes are subcomplexes of hypercubes.
- Hence they are subsemigroups of the face semigroup of the coordinate hyperplane arrangement.

## CAT(0) cube complexes (ctd.)

- Finite CAT(0) cube complexes are subcomplexes of hypercubes.
- Hence they are subsemigroups of the face semigroup of the coordinate hyperplane arrangement.
- The product of a cube  $C$  with a vertex  $v$  is the vertex of  $C$  closest to  $v$ .

## CAT(0) cube complexes (ctd.)

- Finite CAT(0) cube complexes are subcomplexes of hypercubes.
- Hence they are subsemigroups of the face semigroup of the coordinate hyperplane arrangement.
- The product of a cube  $C$  with a vertex  $v$  is the vertex of  $C$  closest to  $v$ .
- This is just as in the case of a zonotope.

## CAT(0) cube complexes (ctd.)

- Finite CAT(0) cube complexes are subcomplexes of hypercubes.
- Hence they are subsemigroups of the face semigroup of the coordinate hyperplane arrangement.
- The product of a cube  $C$  with a vertex  $v$  is the vertex of  $C$  closest to  $v$ .
- This is just as in the case of a zonotope.
- The face semigroup of a CAT(0) cube complex acts on the cube complex.



## CAT(0) cube complexes (ctd.)

- Finite CAT(0) cube complexes are subcomplexes of hypercubes.
- Hence they are subsemigroups of the face semigroup of the coordinate hyperplane arrangement.
- The product of a cube  $C$  with a vertex  $v$  is the vertex of  $C$  closest to  $v$ .
- This is just as in the case of a zonotope.
- The face semigroup of a CAT(0) cube complex acts on the cube complex.
- Again minimal projective resolutions can be constructed from the augmented cellular chain complex.

## Left regular bands (LRBs)

- Real and complex hyperplane face semigroups and their subsemigroups are examples of left regular bands.

## Left regular bands (LRBs)

- Real and complex hyperplane face semigroups and their subsemigroups are examples of left regular bands.

### Definition (LRB)

A **left regular band** is a semigroup  $B$  satisfying the identities:

- $x^2 = x$  *( $B$  is a “band”)*
- $xyx = xy$  *(“left regularity”)*

## Left regular bands (LRBs)

- Real and complex hyperplane face semigroups and their subsemigroups are examples of left regular bands.

### Definition (LRB)

A **left regular band** is a semigroup  $B$  satisfying the identities:

- $x^2 = x$  *( $B$  is a “band”)*
  - $xyx = xy$  *(“left regularity”)*
- Informally: identities say ignore ‘repetitions’.

## Left regular bands (LRBs)

- Real and complex hyperplane face semigroups and their subsemigroups are examples of left regular bands.

### Definition (LRB)

A **left regular band** is a semigroup  $B$  satisfying the identities:

- $x^2 = x$  *( $B$  is a “band”)*
  - $xyx = xy$  *(“left regularity”)*
- 
- Informally: identities say ignore ‘repetitions’.
  - Commutative LRBs are posets with meets.

## Left regular bands (LRBs)

- Real and complex hyperplane face semigroups and their subsemigroups are examples of left regular bands.

### Definition (LRB)

A **left regular band** is a semigroup  $B$  satisfying the identities:

- $x^2 = x$  *( $B$  is a “band”)*
  - $xyx = xy$  *(“left regularity”)*
- 
- Informally: identities say ignore ‘repetitions’.
  - Commutative LRBs are posets with meets.
  - All LRBs are assumed to be finite.

## LRBs as posets

- An LRB  $B$  is a poset with

$$a \leq b \iff aB \subseteq bB \iff ba = a$$

## LRBs as posets

- An LRB  $B$  is a poset with

$$a \leq b \iff aB \subseteq bB \iff ba = a$$

- For hyperplane face semigroups, oriented matroids and CAT(0) cube complexes, this corresponds to inclusion of faces in the dual cell complex.



## LRBs as posets

- An LRB  $B$  is a poset with

$$a \leq b \iff aB \subseteq bB \iff ba = a$$

- For hyperplane face semigroups, oriented matroids and CAT(0) cube complexes, this corresponds to inclusion of faces in the dual cell complex.
- The **support semilattice** is  $\Lambda(B) = \{Bb \mid b \in B\}$  ordered by inclusion.

## LRBs as posets

- An LRB  $B$  is a poset with

$$a \leq b \iff aB \subseteq bB \iff ba = a$$

- For hyperplane face semigroups, oriented matroids and CAT(0) cube complexes, this corresponds to inclusion of faces in the dual cell complex.
- The **support semilattice** is  $\Lambda(B) = \{Bb \mid b \in B\}$  ordered by inclusion.
- It is the **intersection semilattice** in the case of a hyperplane arrangement.

## LRBs as posets

- An LRB  $B$  is a poset with

$$a \leq b \iff aB \subseteq bB \iff ba = a$$

- For hyperplane face semigroups, oriented matroids and CAT(0) cube complexes, this corresponds to inclusion of faces in the dual cell complex.
- The **support semilattice** is  $\Lambda(B) = \{Bb \mid b \in B\}$  ordered by inclusion.
- It is the **intersection semilattice** in the case of a hyperplane arrangement.
- $Ba \cap Bb = Bab$  and so  $\sigma: B \rightarrow \Lambda(B)$  given by  $\sigma(b) = Bb$  is a homomorphism.

## LRBs as posets

- An LRB  $B$  is a poset with

$$a \leq b \iff aB \subseteq bB \iff ba = a$$

- For hyperplane face semigroups, oriented matroids and CAT(0) cube complexes, this corresponds to inclusion of faces in the dual cell complex.
- The **support semilattice** is  $\Lambda(B) = \{Bb \mid b \in B\}$  ordered by inclusion.
- It is the **intersection semilattice** in the case of a hyperplane arrangement.
- $Ba \cap Bb = Bab$  and so  $\sigma: B \rightarrow \Lambda(B)$  given by  $\sigma(b) = Bb$  is a homomorphism.
- $\mathbb{k}B \rightarrow \mathbb{k}\Lambda(B) \cong \mathbb{k}^{\Lambda(B)}$  is the semisimple quotient.

## LRBs as posets

- An LRB  $B$  is a poset with

$$a \leq b \iff aB \subseteq bB \iff ba = a$$

- For hyperplane face semigroups, oriented matroids and CAT(0) cube complexes, this corresponds to inclusion of faces in the dual cell complex.
- The **support semilattice** is  $\Lambda(B) = \{Bb \mid b \in B\}$  ordered by inclusion.
- It is the **intersection semilattice** in the case of a hyperplane arrangement.
- $Ba \cap Bb = Bab$  and so  $\sigma: B \rightarrow \Lambda(B)$  given by  $\sigma(b) = Bb$  is a homomorphism.
- $\mathbb{k}B \rightarrow \mathbb{k}\Lambda(B) \cong \mathbb{k}^{\Lambda(B)}$  is the semisimple quotient.
- So  $\mathbb{k}B$  is split basic over any field.

# The order complex and contractions

- Associated to a poset  $P$  is its **order complex**  $\Delta(P)$ .

## The order complex and contractions

- Associated to a poset  $P$  is its **order complex**  $\Delta(P)$ .
- The vertex set of  $\Delta(P)$  is  $P$ .

## The order complex and contractions

- Associated to a poset  $P$  is its **order complex**  $\Delta(P)$ .
- The vertex set of  $\Delta(P)$  is  $P$ .
- A subset of  $P$  is a simplex if it is a chain.




Figure : Order complex of the power set of  $\{1, 2, 3\}$



## The order complex and contractions

- Associated to a poset  $P$  is its **order complex**  $\Delta(P)$ .
- The vertex set of  $\Delta(P)$  is  $P$ .
- A subset of  $P$  is a simplex if it is a chain.
- If  $B$  is an LRB and  $Ba \in \Lambda(B)$ ,

$$B_{\geq Ba} = \{b \in B \mid Bb \supseteq Ba\}$$

is a subsemigroup called the **contraction** of  $B$  to  $a$ .

## The order complex and contractions

- Associated to a poset  $P$  is its **order complex**  $\Delta(P)$ .
- The vertex set of  $\Delta(P)$  is  $P$ .
- A subset of  $P$  is a simplex if it is a chain.
- If  $B$  is an LRB and  $Ba \in \Lambda(B)$ ,

$$B_{\geq Ba} = \{b \in B \mid Bb \supseteq Ba\}$$

is a subsemigroup called the **contraction** of  $B$  to  $a$ .

- $\mathbb{k}B$  is unital iff  $\Delta(B_{\geq Ba})$  is connected for all  $Ba \in \Lambda(B)$ .

## The order complex and contractions

- Associated to a poset  $P$  is its **order complex**  $\Delta(P)$ .
- The vertex set of  $\Delta(P)$  is  $P$ .
- A subset of  $P$  is a simplex if it is a chain.
- If  $B$  is an LRB and  $Ba \in \Lambda(B)$ ,

$$B_{\geq Ba} = \{b \in B \mid Bb \supseteq Ba\}$$

is a subsemigroup called the **contraction** of  $B$  to  $a$ .

- $\mathbb{k}B$  is unital iff  $\Delta(B_{\geq Ba})$  is connected for all  $Ba \in \Lambda(B)$ .
- There is an epi  $\mathbb{k}B \rightarrow \mathbb{k}B_{\geq Ba}$  sending  $B \setminus B_{\geq Ba}$  to 0.

## The order complex and contractions

- Associated to a poset  $P$  is its **order complex**  $\Delta(P)$ .
- The vertex set of  $\Delta(P)$  is  $P$ .
- A subset of  $P$  is a simplex if it is a chain.
- If  $B$  is an LRB and  $Ba \in \Lambda(B)$ ,

$$B_{\geq Ba} = \{b \in B \mid Bb \supseteq Ba\}$$

is a subsemigroup called the **contraction** of  $B$  to  $a$ .

- $\mathbb{k}B$  is unital iff  $\Delta(B_{\geq Ba})$  is connected for all  $Ba \in \Lambda(B)$ .
- There is an epi  $\mathbb{k}B \rightarrow \mathbb{k}B_{\geq Ba}$  sending  $B \setminus B_{\geq Ba}$  to 0.
- $\mathbb{k}B_{\geq Ba}$  is a projective  $\mathbb{k}B$ -module.

## The order complex and contractions

- Associated to a poset  $P$  is its **order complex**  $\Delta(P)$ .
- The vertex set of  $\Delta(P)$  is  $P$ .
- A subset of  $P$  is a simplex if it is a chain.
- If  $B$  is an LRB and  $Ba \in \Lambda(B)$ ,

$$B_{\geq Ba} = \{b \in B \mid Bb \supseteq Ba\}$$

is a subsemigroup called the **contraction** of  $B$  to  $a$ .

- $\mathbb{k}B$  is unital iff  $\Delta(B_{\geq Ba})$  is connected for all  $Ba \in \Lambda(B)$ .
- There is an epi  $\mathbb{k}B \rightarrow \mathbb{k}B_{\geq Ba}$  sending  $B \setminus B_{\geq Ba}$  to 0.
- $\mathbb{k}B_{\geq Ba}$  is a projective  $\mathbb{k}B$ -module.
- The trivial  $\mathbb{k}B_{\geq Ba}$ -module is a simple  $\mathbb{k}B$ -module

## The order complex and contractions

- Associated to a poset  $P$  is its **order complex**  $\Delta(P)$ .
- The vertex set of  $\Delta(P)$  is  $P$ .
- A subset of  $P$  is a simplex if it is a chain.
- If  $B$  is an LRB and  $Ba \in \Lambda(B)$ ,

$$B_{\geq Ba} = \{b \in B \mid Bb \supseteq Ba\}$$

is a subsemigroup called the **contraction** of  $B$  to  $a$ .

- $\mathbb{k}B$  is unital iff  $\Delta(B_{\geq Ba})$  is connected for all  $Ba \in \Lambda(B)$ .
- There is an epi  $\mathbb{k}B \rightarrow \mathbb{k}B_{\geq Ba}$  sending  $B \setminus B_{\geq Ba}$  to 0.
- $\mathbb{k}B_{\geq Ba}$  is a projective  $\mathbb{k}B$ -module.
- The trivial  $\mathbb{k}B_{\geq Ba}$ -module is a simple  $\mathbb{k}B$ -module
- All simples arise like this.

## Regular CW complexes and face posets

- A **regular CW complex** is a Hausdorff space  $X$  that decomposes as a union of finitely many cells homeomorphic to closed balls such that the boundary of each cell is a union of cells of lower dimension.

## Regular CW complexes and face posets

- A **regular CW complex** is a Hausdorff space  $X$  that decomposes as a union of finitely many cells homeomorphic to closed balls such that the boundary of each cell is a union of cells of lower dimension.
- The **face poset**  $\mathcal{P}(X)$  is the set of cells ordered by inclusion.



## Regular CW complexes and face posets

- A **regular CW complex** is a Hausdorff space  $X$  that decomposes as a union of finitely many cells homeomorphic to closed balls such that the boundary of each cell is a union of cells of lower dimension.
- The **face poset**  $\mathcal{P}(X)$  is the set of cells ordered by inclusion.
- $\Delta(\mathcal{P}(X))$  is the **barycentric subdivision** of  $X$ .

## Regular CW complexes and face posets

- A **regular CW complex** is a Hausdorff space  $X$  that decomposes as a union of finitely many cells homeomorphic to closed balls such that the boundary of each cell is a union of cells of lower dimension.
- The **face poset**  $\mathcal{P}(X)$  is the set of cells ordered by inclusion.
- $\Delta(\mathcal{P}(X))$  is the **barycentric subdivision** of  $X$ .
- So  $X \cong \Delta(\mathcal{P}(X))$ .

## Regular CW complexes and face posets

- A **regular CW complex** is a Hausdorff space  $X$  that decomposes as a union of finitely many cells homeomorphic to closed balls such that the boundary of each cell is a union of cells of lower dimension.
- The **face poset**  $\mathcal{P}(X)$  is the set of cells ordered by inclusion.
- $\Delta(\mathcal{P}(X))$  is the **barycentric subdivision** of  $X$ .
- So  $X \cong \Delta(\mathcal{P}(X))$ .
- Moreover, the cell structure can be functorially recovered the order structure on  $\mathcal{P}(X)$ .

## Regular CW complexes and face posets

- A **regular CW complex** is a Hausdorff space  $X$  that decomposes as a union of finitely many cells homeomorphic to closed balls such that the boundary of each cell is a union of cells of lower dimension.
- The **face poset**  $\mathcal{P}(X)$  is the set of cells ordered by inclusion.
- $\Delta(\mathcal{P}(X))$  is the **barycentric subdivision** of  $X$ .
- So  $X \cong \Delta(\mathcal{P}(X))$ .
- Moreover, the cell structure can be functorially recovered the order structure on  $\mathcal{P}(X)$ .
- There are axioms characterizing face posets of regular CW complexes (cf. Björner).

## CW left regular bands

- We say  $B$  is a **CW LRB** if each contraction  $B_{\geq Ba}$  is the face poset of a connected regular CW complex  $X_{Ba}$ .

## CW left regular bands

- We say  $B$  is a **CW LRB** if each contraction  $B_{\geq Ba}$  is the face poset of a connected regular CW complex  $X_{Ba}$ .
- It turns out that  $X_{Ba}$  is always acyclic.

## CW left regular bands

- We say  $B$  is a **CW LRB** if each contraction  $B_{\geq Ba}$  is the face poset of a connected regular CW complex  $X_{Ba}$ .
- It turns out that  $X_{Ba}$  is always acyclic.
- $\mathbb{k}B$  is a unital algebra over any field.

## CW left regular bands

- We say  $B$  is a **CW LRB** if each contraction  $B_{\geq Ba}$  is the face poset of a connected regular CW complex  $X_{Ba}$ .
- It turns out that  $X_{Ba}$  is always acyclic.
- $\mathbb{k}B$  is a unital algebra over any field.
- Face semigroups of real and complex hyperplane arrangements, oriented matroids and CAT(0) cube complexes all are examples of CW LRBs.



## CW left regular bands

- We say  $B$  is a **CW LRB** if each contraction  $B_{\geq Ba}$  is the face poset of a connected regular CW complex  $X_{Ba}$ .
- It turns out that  $X_{Ba}$  is always acyclic.
- $\mathbb{k}B$  is a unital algebra over any field.
- Face semigroups of real and complex hyperplane arrangements, oriented matroids and CAT(0) cube complexes all are examples of CW LRBs.
- The support semilattice  $\Lambda(B)$  is always Cohen-Macaulay.

## CW left regular bands

- We say  $B$  is a **CW LRB** if each contraction  $B_{\geq Ba}$  is the face poset of a connected regular CW complex  $X_{Ba}$ .
- It turns out that  $X_{Ba}$  is always acyclic.
- $\mathbb{k}B$  is a unital algebra over any field.
- Face semigroups of real and complex hyperplane arrangements, oriented matroids and CAT(0) cube complexes all are examples of CW LRBs.
- The support semilattice  $\Lambda(B)$  is always Cohen-Macaulay.
- A poset  $P$  is **Cohen-Macaulay** if  $\mathbb{k}[P]/(pq \mid p \not\leq q, q \not\leq p)$  is Cohen-Macaulay.

## CW left regular bands

- We say  $B$  is a **CW LRB** if each contraction  $B_{\geq Ba}$  is the face poset of a connected regular CW complex  $X_{Ba}$ .
- It turns out that  $X_{Ba}$  is always acyclic.
- $\mathbb{k}B$  is a unital algebra over any field.
- Face semigroups of real and complex hyperplane arrangements, oriented matroids and  $\text{CAT}(0)$  cube complexes all are examples of CW LRBs.
- The support semilattice  $\Lambda(B)$  is always Cohen-Macaulay.
- A poset  $P$  is **Cohen-Macaulay** if  $\mathbb{k}[P]/(pq \mid p \not\leq q, q \not\leq p)$  is Cohen-Macaulay.
- This depends only on the homeomorphism type of  $\Delta(P)$ .

## A quiver presentation

### Theorem (Margolis,Saliola,BS)

*Let  $B$  be a CW LRB. Let  $Q$  be the Hasse diagram of  $\Lambda(B) = \{Ba \mid a \in B\}$ . Then  $\mathbb{k}B \cong \mathbb{k}Q/(r)$  where  $r$  is the sum of all paths of length 2.*

## A quiver presentation

### Theorem (Margolis,Saliola,BS)

*Let  $B$  be a CW LRB. Let  $Q$  be the Hasse diagram of  $\Lambda(B) = \{Ba \mid a \in B\}$ . Then  $\mathbb{k}B \cong \mathbb{k}Q/(r)$  where  $r$  is the sum of all paths of length 2.*

- Thus  $\mathbb{k}B$  is graded (by path length) and quadratic.

## A quiver presentation

### Theorem (Margolis,Saliola,BS)

*Let  $B$  be a CW LRB. Let  $Q$  be the Hasse diagram of  $\Lambda(B) = \{Ba \mid a \in B\}$ . Then  $\mathbb{k}B \cong \mathbb{k}Q/(r)$  where  $r$  is the sum of all paths of length 2.*

- Thus  $\mathbb{k}B$  is graded (by path length) and quadratic.
- Notice that  $\mathbb{k}B$  depends only on  $\Lambda(B)$  up to isomorphism.

## A quiver presentation

### Theorem (Margolis,Saliola,BS)

*Let  $B$  be a CW LRB. Let  $Q$  be the Hasse diagram of  $\Lambda(B) = \{Ba \mid a \in B\}$ . Then  $\mathbb{k}B \cong \mathbb{k}Q/(r)$  where  $r$  is the sum of all paths of length 2.*

- Thus  $\mathbb{k}B$  is graded (by path length) and quadratic.
- Notice that  $\mathbb{k}B$  depends only on  $\Lambda(B)$  up to isomorphism.
- This was proved for the face monoid of a central real hyperplane arrangement by Saliola.

## A quiver presentation

### Theorem (Margolis,Saliola,BS)

*Let  $B$  be a CW LRB. Let  $Q$  be the Hasse diagram of  $\Lambda(B) = \{Ba \mid a \in B\}$ . Then  $\mathbb{k}B \cong \mathbb{k}Q/(r)$  where  $r$  is the sum of all paths of length 2.*

- Thus  $\mathbb{k}B$  is graded (by path length) and quadratic.
- Notice that  $\mathbb{k}B$  depends only on  $\Lambda(B)$  up to isomorphism.
- This was proved for the face monoid of a central real hyperplane arrangement by Saliola.
- Our approach was inspired by his, but is more conceptual, which in turn led to simplifications.



# A Koszul algebra

## Theorem (Margolis,Saliola,BS)

*Let  $B$  be a CW LRB. Then  $\mathbb{k}B$  is a **Koszul algebra** with respect to the grading of the previous theorem. Moreover,  $\text{Ext}(\mathbb{k}B)$  is isomorphic to the **incidence algebra** of  $\Lambda(B)$ .*

# A Koszul algebra

## Theorem (Margolis,Saliola,BS)

*Let  $B$  be a CW LRB. Then  $\mathbb{k}B$  is a **Koszul algebra** with respect to the grading of the previous theorem. Moreover,  $\text{Ext}(\mathbb{k}B)$  is isomorphic to the **incidence algebra** of  $\Lambda(B)$ .*

- Again, this was proved by Saliola in the case of the face monoid of a central real hyperplane arrangement.

# A Koszul algebra

## Theorem (Margolis,Saliola,BS)

*Let  $B$  be a CW LRB. Then  $\mathbb{k}B$  is a **Koszul algebra** with respect to the grading of the previous theorem. Moreover,  $\text{Ext}(\mathbb{k}B)$  is isomorphic to the **incidence algebra** of  $\Lambda(B)$ .*

- Again, this was proved by Saliola in the case of the face monoid of a central real hyperplane arrangement.
- Polo and Woodcock independently proved a poset is Cohen-Macaulay iff its incidence algebra is Koszul.

# A Koszul algebra

## Theorem (Margolis,Saliola,BS)

*Let  $B$  be a CW LRB. Then  $\mathbb{k}B$  is a **Koszul algebra** with respect to the grading of the previous theorem. Moreover,  $\text{Ext}(\mathbb{k}B)$  is isomorphic to the **incidence algebra** of  $\Lambda(B)$ .*

- Again, this was proved by Saliola in the case of the face monoid of a central real hyperplane arrangement.
- Polo and Woodcock independently proved a poset is Cohen-Macaulay iff its incidence algebra is Koszul.
- This is how we obtained that  $\Lambda(B)$  is Cohen-Macaulay.

# Minimal projective resolutions

- Let  $B$  be a CW LRB.

## Minimal projective resolutions

- Let  $B$  be a CW LRB.
- By passing to contractions it suffices to compute a minimal projective resolution for the trivial module.

## Minimal projective resolutions

- Let  $B$  be a CW LRB.
- By passing to contractions it suffices to compute a minimal projective resolution for the trivial module.
- Let  $B$  be the face poset of the regular CW complex  $X$ .

## Minimal projective resolutions

- Let  $B$  be a CW LRB.
- By passing to contractions it suffices to compute a minimal projective resolution for the trivial module.
- Let  $B$  be the face poset of the regular CW complex  $X$ .
- $B$  acts on  $X$  by cellular maps.



## Minimal projective resolutions

- Let  $B$  be a CW LRB.
- By passing to contractions it suffices to compute a minimal projective resolution for the trivial module.
- Let  $B$  be the face poset of the regular CW complex  $X$ .
- $B$  acts on  $X$  by cellular maps.
- Since  $X$  is acyclic, the augmented cellular chain complex of  $X$  is a  $\mathbb{k}B$ -resolution of  $\mathbb{k}$ .

## Minimal projective resolutions

- Let  $B$  be a CW LRB.
- By passing to contractions it suffices to compute a minimal projective resolution for the trivial module.
- Let  $B$  be the face poset of the regular CW complex  $X$ .
- $B$  acts on  $X$  by cellular maps.
- Since  $X$  is acyclic, the augmented cellular chain complex of  $X$  is a  $\mathbb{k}B$ -resolution of  $\mathbb{k}$ .
- We have shown it is the minimal projective resolution.

## Minimal projective resolutions

- Let  $B$  be a CW LRB.
- By passing to contractions it suffices to compute a minimal projective resolution for the trivial module.
- Let  $B$  be the face poset of the regular CW complex  $X$ .
- $B$  acts on  $X$  by cellular maps.
- Since  $X$  is acyclic, the augmented cellular chain complex of  $X$  is a  $\mathbb{k}B$ -resolution of  $\mathbb{k}$ .
- We have shown it is the minimal projective resolution.
- It is straightforward to show it is a linear resolution.

## Minimal projective resolutions

- Let  $B$  be a CW LRB.
- By passing to contractions it suffices to compute a minimal projective resolution for the trivial module.
- Let  $B$  be the face poset of the regular CW complex  $X$ .
- $B$  acts on  $X$  by cellular maps.
- Since  $X$  is acyclic, the augmented cellular chain complex of  $X$  is a  $\mathbb{k}B$ -resolution of  $\mathbb{k}$ .
- We have shown it is the minimal projective resolution.
- It is straightforward to show it is a linear resolution.
- This was obtained in an ad hoc way by Brown, Diaconis and Saliola for real central hyperplane arrangements.

## Minimal projective resolutions

- Let  $B$  be a CW LRB.
- By passing to contractions it suffices to compute a minimal projective resolution for the trivial module.
- Let  $B$  be the face poset of the regular CW complex  $X$ .
- $B$  acts on  $X$  by cellular maps.
- Since  $X$  is acyclic, the augmented cellular chain complex of  $X$  is a  $\mathbb{k}B$ -resolution of  $\mathbb{k}$ .
- We have shown it is the minimal projective resolution.
- It is straightforward to show it is a linear resolution.
- This was obtained in an ad hoc way by Brown, Diaconis and Saliola for real central hyperplane arrangements.
- It follows that the global dimension of  $\mathbb{k}B$  is  $\dim X$ .

## An example

- Let  $L_n = \{0, \pm 1, \dots, \pm n\}$  with the product given by

$$xy = \begin{cases} y, & \text{if } |x| < |y| \\ x, & \text{if } |x| \geq |y|. \end{cases}$$

## An example

- Let  $L_n = \{0, \pm 1, \dots, \pm n\}$  with the product given by

$$xy = \begin{cases} y, & \text{if } |x| < |y| \\ x, & \text{if } |x| \geq |y|. \end{cases}$$

- $L_n$  is a CW LRB which is the face poset of a regular cell decomposition of the closed  $n$ -ball.





Figure : The Hasse diagram of  $L_3$  and cell decomposition of the 3-ball.

## An example

- Let  $L_n = \{0, \pm 1, \dots, \pm n\}$  with the product given by

$$xy = \begin{cases} y, & \text{if } |x| < |y| \\ x, & \text{if } |x| \geq |y|. \end{cases}$$

- $L_n$  is a CW LRB which is the face poset of a regular cell decomposition of the closed  $n$ -ball.
- $\Lambda(L_n) \cong \{0 < 1 < \dots < n\}$  via  $L_n x \mapsto n - |x|$ .



## An example

- Let  $L_n = \{0, \pm 1, \dots, \pm n\}$  with the product given by

$$xy = \begin{cases} y, & \text{if } |x| < |y| \\ x, & \text{if } |x| \geq |y|. \end{cases}$$

- $L_n$  is a CW LRB which is the face poset of a regular cell decomposition of the closed  $n$ -ball.
- $\Lambda(L_n) \cong \{0 < 1 < \dots < n\}$  via  $L_n x \mapsto n - |x|$ .
- The Hasse diagram of  $L_n$  is the  $A_{n+1}$ -quiver.

$$0 \longrightarrow 1 \longrightarrow 2 \longrightarrow 3$$

## An example

- Let  $L_n = \{0, \pm 1, \dots, \pm n\}$  with the product given by

$$xy = \begin{cases} y, & \text{if } |x| < |y| \\ x, & \text{if } |x| \geq |y|. \end{cases}$$

- $L_n$  is a CW LRB which is the face poset of a regular cell decomposition of the closed  $n$ -ball.
- $\Lambda(L_n) \cong \{0 < 1 < \dots < n\}$  via  $L_n x \mapsto n - |x|$ .
- The Hasse diagram of  $L_n$  is the  $A_{n+1}$ -quiver.

$$0 \longrightarrow 1 \longrightarrow 2 \longrightarrow 3$$

- $\mathbb{k}L_n \cong \mathbb{k}A_{n+1}/\text{Rad}^2(\mathbb{k}A_{n+1})$ .

## An example

- Let  $L_n = \{0, \pm 1, \dots, \pm n\}$  with the product given by

$$xy = \begin{cases} y, & \text{if } |x| < |y| \\ x, & \text{if } |x| \geq |y|. \end{cases}$$

- $L_n$  is a CW LRB which is the face poset of a regular cell decomposition of the closed  $n$ -ball.
- $\Lambda(L_n) \cong \{0 < 1 < \dots < n\}$  via  $L_n x \mapsto n - |x|$ .
- The Hasse diagram of  $L_n$  is the  $A_{n+1}$ -quiver.

$$0 \longrightarrow 1 \longrightarrow 2 \longrightarrow 3$$

- $\mathbb{k}L_n \cong \mathbb{k}A_{n+1}/\text{Rad}^2(\mathbb{k}A_{n+1})$ .
- It is Koszul with Ext-algebra isomorphic  $\mathbb{k}A_{n+1}$ , which is the incidence algebra of  $\{0 < 1 < \dots < n\}$ .

# The end

# Thank you for your attention!