

The Karoubi envelope and the classification of Markov-Dyck shifts

Alfredo Costa, Universidade de Coimbra

Benjamin Steinberg, City College of New York



November 21, 2013

Outline

- 1 The Karoubi Envelope
- 2 Markov-Dyck shifts: a success story
- 3 Conjugacy Invariance

Goals

- Our goal is to introduce a syntactic invariant of flow equivalence of shifts.

Goals

- Our goal is to introduce a syntactic invariant of flow equivalence of shifts.
- The invariant is a small category constructed from the syntactic semigroup of the shift.

Goals

- Our goal is to introduce a syntactic invariant of flow equivalence of shifts.
- The invariant is a small category constructed from the syntactic semigroup of the shift.
- The behavior of this invariant is somewhat orthogonal to classical invariants.

Goals

- Our goal is to introduce a syntactic invariant of flow equivalence of shifts.
- The invariant is a small category constructed from the syntactic semigroup of the shift.
- The behavior of this invariant is somewhat orthogonal to classical invariants.
- We will show how it classifies Markov-Dyck shifts up to flow equivalence under mild hypotheses on the graphs.

Goals

- Our goal is to introduce a syntactic invariant of flow equivalence of shifts.
- The invariant is a small category constructed from the syntactic semigroup of the shift.
- The behavior of this invariant is somewhat orthogonal to classical invariants.
- We will show how it classifies Markov-Dyck shifts up to flow equivalence under mild hypotheses on the graphs.
- I will sketch a simple proof of our main result using results of Nasu.

The Karoubi envelope of a semigroup

- Let S be a semigroup.

The Karoubi envelope of a semigroup

- Let S be a semigroup.
- The Karoubi envelope $\mathbb{K}(S)$ is the following small category.

The Karoubi envelope of a semigroup

- Let S be a semigroup.
- The Karoubi envelope $\mathbb{K}(S)$ is the following small category.
- $\text{Obj}(\mathbb{K}(S)) = E(S)$ (the idempotents of S).

The Karoubi envelope of a semigroup

- Let S be a semigroup.
- The Karoubi envelope $\mathbb{K}(S)$ is the following small category.
- $\text{Obj}(\mathbb{K}(S)) = E(S)$ (the idempotents of S).
- An arrow $e \rightarrow f$ is a triple (f, s, e) with $s \in fSe$.
That is, $fs = s = se$.

The Karoubi envelope of a semigroup

- Let S be a semigroup.
- The Karoubi envelope $\mathbb{K}(S)$ is the following small category.
- $\text{Obj}(\mathbb{K}(S)) = E(S)$ (the idempotents of S).
- An arrow $e \rightarrow f$ is a triple (f, s, e) with $s \in fSe$.
That is, $fs = s = se$.
- $(f, s, e)(e, s', e') = (f, ss', e')$

The Karoubi envelope of a semigroup

- Let S be a semigroup.
- The Karoubi envelope $\mathbb{K}(S)$ is the following small category.
- $\text{Obj}(\mathbb{K}(S)) = E(S)$ (the idempotents of S).
- An arrow $e \rightarrow f$ is a triple (f, s, e) with $s \in fSe$.
That is, $fs = s = se$.
- $(f, s, e)(e, s', e') = (f, ss', e')$
- The identity at e is (e, e, e) .

The Karoubi envelope of a semigroup

- Let S be a semigroup.
- The Karoubi envelope $\mathbb{K}(S)$ is the following small category.
- $\text{Obj}(\mathbb{K}(S)) = E(S)$ (the idempotents of S).
- An arrow $e \rightarrow f$ is a triple (f, s, e) with $s \in fSe$.
That is, $fs = s = se$.
- $(f, s, e)(e, s', e') = (f, ss', e')$
- The identity at e is (e, e, e) .
- The endomorphism monoid at e is eSe .

The Karoubi envelope of a semigroup

- Let S be a semigroup.
- The Karoubi envelope $\mathbb{K}(S)$ is the following small category.
- $\text{Obj}(\mathbb{K}(S)) = E(S)$ (the idempotents of S).
- An arrow $e \rightarrow f$ is a triple (f, s, e) with $s \in fSe$.
That is, $fs = s = se$.
- $(f, s, e)(e, s', e') = (f, ss', e')$
- The identity at e is (e, e, e) .
- The endomorphism monoid at e is eSe .
- $e \cong f$ iff $e = ab$ and $f = ba$ for some $a, b \in S$ with $aba = a, bab = b$. (Called \mathcal{D} -equivalence.)

The Karoubi envelope of a semigroup

- Let S be a semigroup.
- The Karoubi envelope $\mathbb{K}(S)$ is the following small category.
- $\text{Obj}(\mathbb{K}(S)) = E(S)$ (the idempotents of S).
- An arrow $e \rightarrow f$ is a triple (f, s, e) with $s \in fSe$.
That is, $fs = s = se$.
- $(f, s, e)(e, s', e') = (f, ss', e')$
- The identity at e is (e, e, e) .
- The endomorphism monoid at e is eSe .
- $e \cong f$ iff $e = ab$ and $f = ba$ for some $a, b \in S$ with $aba = a, bab = b$. (Called \mathcal{D} -equivalence.)
- This is analogous to von Neumann-Murray equivalence of projections.

A more conceptual definition

- A right S -set is a set X with an action $X \times S \rightarrow X$.

A more conceptual definition

- A right S -set is a set X with an action $X \times S \rightarrow X$.
- $\mathbb{K}(S)$ is equivalent to the category of S -sets of the form eS , with $e \in E(S)$, and S -equivariant maps.

A more conceptual definition

- A right S -set is a set X with an action $X \times S \rightarrow X$.
- $\mathbb{K}(S)$ is equivalent to the category of S -sets of the form eS , with $e \in E(S)$, and S -equivariant maps.
- You should think of eS as a projective indecomposable.

A more conceptual definition

- A right S -set is a set X with an action $X \times S \rightarrow X$.
- $\mathbb{K}(S)$ is equivalent to the category of S -sets of the form eS , with $e \in E(S)$, and S -equivariant maps.
- You should think of eS as a projective indecomposable.
- Hence if $\mathbb{K}(S)$ is equivalent to $\mathbb{K}(T)$, we should think of S and T as Morita equivalent in some sense.

Semigroups with local units

- A semigroup S has **local units** if $E(S)SE(S) = S$.

Semigroups with local units

- A semigroup S has **local units** if $E(S)SE(S) = S$.
- Equivalently, for all $s \in S$, there are $e, f \in E(S)$ with $fs = s = se$.

Semigroups with local units

- A semigroup S has **local units** if $E(S)SE(S) = S$.
- Equivalently, for all $s \in S$, there are $e, f \in E(S)$ with $fs = s = se$.
- Every monoid has local units.

Semigroups with local units

- A semigroup S has **local units** if $E(S)SE(S) = S$.
- Equivalently, for all $s \in S$, there are $e, f \in E(S)$ with $fs = s = se$.
- Every monoid has local units.
- If S is any semigroup, then $LU(S) = E(S)SE(S)$ is the largest subsemigroup of S with local units.

Semigroups with local units

- A semigroup S has **local units** if $E(S)SE(S) = S$.
- Equivalently, for all $s \in S$, there are $e, f \in E(S)$ with $fs = s = se$.
- Every monoid has local units.
- If S is any semigroup, then $LU(S) = E(S)SE(S)$ is the largest subsemigroup of S with local units.
- From the construction, $\mathbb{K}(S) = \mathbb{K}(LU(S))$.

Morita equivalence of semigroups with local units

- Let S be a semigroup with local units.

Morita equivalence of semigroups with local units

- Let S be a semigroup with local units.
- An S -set X is **closed** if the natural map $X \otimes_S S \rightarrow X$ given by $x \otimes s \mapsto xs$ is a bijection.

Morita equivalence of semigroups with local units

- Let S be a semigroup with local units.
- An S -set X is **closed** if the natural map $X \otimes_S S \rightarrow X$ given by $x \otimes s \mapsto xs$ is a bijection.
- For a monoid S , all unital S -sets are closed.

Morita equivalence of semigroups with local units

- Let S be a semigroup with local units.
- An S -set X is **closed** if the natural map $X \otimes_S S \rightarrow X$ given by $x \otimes s \mapsto xs$ is a bijection.
- For a monoid S , all unital S -sets are closed.
- Talwar defined semigroups S, T with local units to be **Morita equivalent** if the category of closed S -sets is equivalent to the category of closed T -sets.

Morita equivalence of semigroups with local units

- Let S be a semigroup with local units.
- An S -set X is **closed** if the natural map $X \otimes_S S \rightarrow X$ given by $x \otimes s \mapsto xs$ is a bijection.
- For a monoid S , all unital S -sets are closed.
- Talwar defined semigroups S, T with local units to be **Morita equivalent** if the category of closed S -sets is equivalent to the category of closed T -sets.
- Lawson first explicitly proved S is Morita equivalent to T iff $\mathbb{K}(S)$ is equivalent to $\mathbb{K}(T)$.

Morita equivalence of semigroups with local units

- Let S be a semigroup with local units.
- An S -set X is **closed** if the natural map $X \otimes_S S \rightarrow X$ given by $x \otimes s \mapsto xs$ is a bijection.
- For a monoid S , all unital S -sets are closed.
- Talwar defined semigroups S, T with local units to be **Morita equivalent** if the category of closed S -sets is equivalent to the category of closed T -sets.
- Lawson first explicitly proved S is Morita equivalent to T iff $\mathbb{K}(S)$ is equivalent to $\mathbb{K}(T)$.
- If S, T are any semigroups, we say S and T are **Morita equivalent up to local units** if $\mathbb{K}(S)$ is equivalent to $\mathbb{K}(T)$.

Examples

- A semigroup $S = LU(S)$ is Morita equivalent to a monoid M iff there is $e \in E(S)$ with $S = SeS$ and $eSe \cong M$.

Examples

- A semigroup $S = LU(S)$ is Morita equivalent to a monoid M iff there is $e \in E(S)$ with $S = SeS$ and $eSe \cong M$.
- In particular, S is Morita equivalent to a group G iff S is completely simple with maximal subgroup G .

Examples

- A semigroup $S = LU(S)$ is Morita equivalent to a monoid M iff there is $e \in E(S)$ with $S = SeS$ and $eSe \cong M$.
- In particular, S is Morita equivalent to a group G iff S is completely simple with maximal subgroup G .
- Morita equivalent semigroups with local units have a number of structural properties in common.

Examples

- A semigroup $S = LU(S)$ is Morita equivalent to a monoid M iff there is $e \in E(S)$ with $S = SeS$ and $eSe \cong M$.
- In particular, S is Morita equivalent to a group G iff S is completely simple with maximal subgroup G .
- Morita equivalent semigroups with local units have a number of structural properties in common.
- They have isomorphic lattices of ideals.

Examples

- A semigroup $S = LU(S)$ is Morita equivalent to a monoid M iff there is $e \in E(S)$ with $S = SeS$ and $eSe \cong M$.
- In particular, S is Morita equivalent to a group G iff S is completely simple with maximal subgroup G .
- Morita equivalent semigroups with local units have a number of structural properties in common.
- They have isomorphic lattices of ideals.
- Their sets of \mathcal{D} -classes of idempotents are in bijection, and moreover the bijection preserves maximal subgroups.

Examples

- A semigroup $S = LU(S)$ is Morita equivalent to a monoid M iff there is $e \in E(S)$ with $S = SeS$ and $eSe \cong M$.
- In particular, S is Morita equivalent to a group G iff S is completely simple with maximal subgroup G .
- Morita equivalent semigroups with local units have a number of structural properties in common.
- They have isomorphic lattices of ideals.
- Their sets of \mathcal{D} -classes of idempotents are in bijection, and moreover the bijection preserves maximal subgroups.
- Our syntactic invariants of shifts mostly come from Morita invariant properties of semigroups with local units.

Shifts

- Let $\mathcal{X} \subseteq A^{\mathbb{Z}}$ be a subshift.

Shifts

- Let $\mathcal{X} \subseteq A^{\mathbb{Z}}$ be a subshift.
- $L(\mathcal{X})$ denotes the language of blocks (or factors) of \mathcal{X} .

Shifts

- Let $\mathcal{X} \subseteq A^{\mathbb{Z}}$ be a subshift.
- $L(\mathcal{X})$ denotes the language of blocks (or factors) of \mathcal{X} .
- It is a nonempty, factorial, prolongable language and all such languages come from shifts.

Shifts

- Let $\mathcal{X} \subseteq A^{\mathbb{Z}}$ be a subshift.
- $L(\mathcal{X})$ denotes the language of blocks (or factors) of \mathcal{X} .
- It is a nonempty, factorial, prolongable language and all such languages come from shifts.
- Many syntactic invariants of \mathcal{X} have been constructed from $L(\mathcal{X})$ via its syntactic semigroup.

The syntactic semigroup of a shift

- A is a finite alphabet.

The syntactic semigroup of a shift

- A is a finite alphabet.
- A^+ (resp. A^*) is the free semigroup (monoid) on A .

The syntactic semigroup of a shift

- A is a finite alphabet.
- A^+ (resp. A^*) is the free semigroup (monoid) on A .
- The **syntactic congruence** for $L \subseteq A^+$ is defined by
 - $x \equiv_L y$ if, for all $u, v \in A^*$, we have $uxv \in L \iff uyv \in L$.

The syntactic semigroup of a shift

- A is a finite alphabet.
- A^+ (resp. A^*) is the free semigroup (monoid) on A .
- The **syntactic congruence** for $L \subseteq A^+$ is defined by
 - $x \equiv_L y$ if, for all $u, v \in A^*$, we have $uxv \in L \iff uyv \in L$.
- The quotient semigroup $S(L) = A^+ / \equiv_L$ is called the **syntactic semigroup** of L .

The syntactic semigroup of a shift

- A is a finite alphabet.
- A^+ (resp. A^*) is the free semigroup (monoid) on A .
- The **syntactic congruence** for $L \subseteq A^+$ is defined by
 - $x \equiv_L y$ if, for all $u, v \in A^*$, we have $uxv \in L \iff uyv \in L$.
- The quotient semigroup $S(L) = A^+ / \equiv_L$ is called the **syntactic semigroup** of L .
- For a subshift $\mathcal{X} \subseteq A^{\mathbb{Z}}$, we put $S(\mathcal{X}) = S(L(\mathcal{X}))$ and call it the **syntactic semigroup** of \mathcal{X} .

The syntactic semigroup of a shift

- A is a finite alphabet.
- A^+ (resp. A^*) is the free semigroup (monoid) on A .
- The **syntactic congruence** for $L \subseteq A^+$ is defined by
 - $x \equiv_L y$ if, for all $u, v \in A^*$, we have $uxv \in L \iff uyv \in L$.
- The quotient semigroup $S(L) = A^+ / \equiv_L$ is called the **syntactic semigroup** of L .
- For a subshift $\mathcal{X} \subseteq A^{\mathbb{Z}}$, we put $S(\mathcal{X}) = S(L(\mathcal{X}))$ and call it the **syntactic semigroup of \mathcal{X}** .
- If $\mathcal{X} \subsetneq A^{\mathbb{Z}}$, then $S(\mathcal{X})$ contains a zero element (the class of a non-factor).

The syntactic semigroup of a shift

- A is a finite alphabet.
- A^+ (resp. A^*) is the free semigroup (monoid) on A .
- The **syntactic congruence** for $L \subseteq A^+$ is defined by
 - $x \equiv_L y$ if, for all $u, v \in A^*$, we have $uxv \in L \iff uyv \in L$.
- The quotient semigroup $S(L) = A^+ / \equiv_L$ is called the **syntactic semigroup** of L .
- For a subshift $\mathcal{X} \subseteq A^{\mathbb{Z}}$, we put $S(\mathcal{X}) = S(L(\mathcal{X}))$ and call it the **syntactic semigroup of \mathcal{X}** .
- If $\mathcal{X} \subsetneq A^{\mathbb{Z}}$, then $S(\mathcal{X})$ contains a zero element (the class of a non-factor).
- **Convention:** We put $S(A^{\mathbb{Z}}) = S(A^+) \cup \{0\}$.

Some basic properties of $S(\mathcal{X})$

- \mathcal{X} is **sofic** iff $S(\mathcal{X})$ is finite.

Some basic properties of $S(\mathcal{X})$

- \mathcal{X} is **sofic** iff $S(\mathcal{X})$ is finite.
- A sofic shift \mathcal{X} is **irreducible** iff $S(\mathcal{X})$ has a unique minimal non-zero ideal.

Some basic properties of $S(\mathcal{X})$

- \mathcal{X} is **sofic** iff $S(\mathcal{X})$ is finite.
- A sofic shift \mathcal{X} is **irreducible** iff $S(\mathcal{X})$ has a unique minimal non-zero ideal.
- An irreducible sofic shift \mathcal{X} is of **finite** type iff $eS(\mathcal{X})e$ is commutative for all $e \in E(S(\mathcal{X}))$.

Some basic properties of $S(\mathcal{X})$

- \mathcal{X} is **sofic** iff $S(\mathcal{X})$ is finite.
- A sofic shift \mathcal{X} is **irreducible** iff $S(\mathcal{X})$ has a unique minimal non-zero ideal.
- An irreducible sofic shift \mathcal{X} is of **finite** type iff $eS(\mathcal{X})e$ is commutative for all $e \in E(S(\mathcal{X}))$.
- An irreducible sofic shift is of **almost finite type** iff $eS(\mathcal{X})e$ has commuting idempotents for all $e \in E(S(\mathcal{X}))$.

Some basic properties of $S(\mathcal{X})$

- \mathcal{X} is **sofic** iff $S(\mathcal{X})$ is finite.
- A sofic shift \mathcal{X} is **irreducible** iff $S(\mathcal{X})$ has a unique minimal non-zero ideal.
- An irreducible sofic shift \mathcal{X} is of **finite** type iff $eS(\mathcal{X})e$ is commutative for all $e \in E(S(\mathcal{X}))$.
- An irreducible sofic shift is of **almost finite type** iff $eS(\mathcal{X})e$ has commuting idempotents for all $e \in E(S(\mathcal{X}))$.
- These last three properties are all Morita invariants up to local units.

The Karoubi envelope of a shift

- If \mathcal{X} is a shift, we put $\mathbb{K}(\mathcal{X}) = \mathbb{K}(S(\mathcal{X}))$ and call it the Karoubi envelope of \mathcal{X} .

The Karoubi envelope of a shift

- If \mathcal{X} is a shift, we put $\mathbb{K}(\mathcal{X}) = \mathbb{K}(S(\mathcal{X}))$ and call it the Karoubi envelope of \mathcal{X} .

Theorem (Costa/BS)

If \mathcal{X} and \mathcal{Y} are flow equivalent shifts, then $\mathbb{K}(\mathcal{X})$ is naturally equivalent to $\mathbb{K}(\mathcal{Y})$.

The Karoubi envelope of a shift

- If \mathcal{X} is a shift, we put $\mathbb{K}(\mathcal{X}) = \mathbb{K}(S(\mathcal{X}))$ and call it the Karoubi envelope of \mathcal{X} .

Theorem (Costa/BS)

If \mathcal{X} and \mathcal{Y} are flow equivalent shifts, then $\mathbb{K}(\mathcal{X})$ is naturally equivalent to $\mathbb{K}(\mathcal{Y})$.

- In other words, the Morita equivalence class of $S(\mathcal{X})$ (up to local units) is a flow equivalence invariant of \mathcal{X} .

The Karoubi envelope of a shift

- If \mathcal{X} is a shift, we put $\mathbb{K}(\mathcal{X}) = \mathbb{K}(S(\mathcal{X}))$ and call it the Karoubi envelope of \mathcal{X} .

Theorem (Costa/BS)

If \mathcal{X} and \mathcal{Y} are flow equivalent shifts, then $\mathbb{K}(\mathcal{X})$ is naturally equivalent to $\mathbb{K}(\mathcal{Y})$.

- In other words, the Morita equivalence class of $S(\mathcal{X})$ (up to local units) is a flow equivalence invariant of \mathcal{X} .
- Alfredo will explain in his talk how a number of syntactic invariants of shifts in the literature can be deduced from this result.

The good, the bad and the ugly

The good news:

The Karoubi envelope can tell if an irreducible sofic shift is:

The good, the bad and the ugly

The good news:

The Karoubi envelope can tell if an irreducible sofic shift is:

- finite type

The good, the bad and the ugly

The good news:

The Karoubi envelope can tell if an irreducible sofic shift is:

- finite type
- almost finite type

The good, the bad and the ugly

The good news:

The Karoubi envelope can tell if an irreducible sofic shift is:

- finite type
- almost finite type

The bad news:

The good, the bad and the ugly

The good news:

The Karoubi envelope can tell if an irreducible sofic shift is:

- finite type
- almost finite type

The bad news:

- All finite type shifts have equivalent Karoubi envelopes.

The good, the bad and the ugly

The good news:

The Karoubi envelope can tell if an irreducible sofic shift is:

- finite type
- almost finite type

The bad news:

- All finite type shifts have equivalent Karoubi envelopes.
- All minimal shifts have equivalent Karoubi envelopes.

Graph inverse semigroups

- Let $G = (V, E)$ be a finite (di)graph.

Graph inverse semigroups

- Let $G = (V, E)$ be a finite (di)graph.
- Assume G has no sinks and no vertex of in-degree 1.

Graph inverse semigroups

- Let $G = (V, E)$ be a finite (di)graph.
- Assume G has no sinks and no vertex of in-degree 1.
- G^* denotes the set of all (directed) paths in G .

Graph inverse semigroups

- Let $G = (V, E)$ be a finite (di)graph.
- Assume G has no sinks and no vertex of in-degree 1.
- G^* denotes the set of all (directed) paths in G .
- There is an empty path at each vertex; path composition is right-to-left.

Graph inverse semigroups

- Let $G = (V, E)$ be a finite (di)graph.
- Assume G has no sinks and no vertex of in-degree 1.
- G^* denotes the set of all (directed) paths in G .
- There is an empty path at each vertex; path composition is right-to-left.
- Let $\mathfrak{h} = \ell_2(G^*)$.

Graph inverse semigroups

- Let $G = (V, E)$ be a finite (di)graph.
- Assume G has no sinks and no vertex of in-degree 1.
- G^* denotes the set of all (directed) paths in G .
- There is an empty path at each vertex; path composition is right-to-left.
- Let $\mathfrak{h} = \ell_2(G^*)$.
- For each path p , define a partial isometry s_p on \mathfrak{h} by

$$s_p(q) = \begin{cases} pq, & \text{if } pq \text{ is a path} \\ 0, & \text{else.} \end{cases}$$

Graph inverse semigroups

- Let $G = (V, E)$ be a finite (di)graph.
- Assume G has no sinks and no vertex of in-degree 1.
- G^* denotes the set of all (directed) paths in G .
- There is an empty path at each vertex; path composition is right-to-left.
- Let $\mathfrak{h} = \ell_2(G^*)$.
- For each path p , define a partial isometry s_p on \mathfrak{h} by

$$s_p(q) = \begin{cases} pq, & \text{if } pq \text{ is a path} \\ 0, & \text{else.} \end{cases}$$

- $P_G = \langle \{s_e, s_e^* \mid e \in E\} \rangle$ is called the **graph inverse semigroup** associated to G .

Graph inverse semigroups

- Let $G = (V, E)$ be a finite (di)graph.
- Assume G has no sinks and no vertex of in-degree 1.
- G^* denotes the set of all (directed) paths in G .
- There is an empty path at each vertex; path composition is right-to-left.
- Let $\mathfrak{h} = \ell_2(G^*)$.
- For each path p , define a partial isometry s_p on \mathfrak{h} by

$$s_p(q) = \begin{cases} pq, & \text{if } pq \text{ is a path} \\ 0, & \text{else.} \end{cases}$$

- $P_G = \langle \{s_e, s_e^* \mid e \in E\} \rangle$ is called the **graph inverse semigroup** associated to G .
- Each nonzero element of P_G can be uniquely written $s_p s_q^*$.

Inverse semigroups

- S is an **inverse semigroup** if, for all $s \in S$, there exists *unique* $s^* \in S$ such that $ss^*s = s$ and $s^*ss^* = s^*$.

Inverse semigroups

- S is an **inverse semigroup** if, for all $s \in S$, there exists *unique* $s^* \in S$ such that $ss^*s = s$ and $s^*ss^* = s^*$.
- $ss^*, s^*s \in E(S)$ and so S has local units.

Inverse semigroups

- S is an **inverse semigroup** if, for all $s \in S$, there exists *unique* $s^* \in S$ such that $ss^*s = s$ and $s^*ss^* = s^*$.
- $ss^*, s^*s \in E(S)$ and so S has local units.
- Inverse semigroups are precisely $*$ -semigroups of partial isometries of Hilbert spaces.

Inverse semigroups

- S is an **inverse semigroup** if, for all $s \in S$, there exists *unique* $s^* \in S$ such that $ss^*s = s$ and $s^*ss^* = s^*$.
- $ss^*, s^*s \in E(S)$ and so S has local units.
- Inverse semigroups are precisely $*$ -semigroups of partial isometries of Hilbert spaces.
- Graph inverse semigroups were considered first in the 70s by Ash and Hall for graphs without multiple edges.

Inverse semigroups

- S is an **inverse semigroup** if, for all $s \in S$, there exists *unique* $s^* \in S$ such that $ss^*s = s$ and $s^*ss^* = s^*$.
- $ss^*, s^*s \in E(S)$ and so S has local units.
- Inverse semigroups are precisely $*$ -semigroups of partial isometries of Hilbert spaces.
- Graph inverse semigroups were considered first in the 70s by Ash and Hall for graphs without multiple edges.
- The general case was studied by Paterson in connection with graph C^* -algebras.

Inverse semigroups

- S is an **inverse semigroup** if, for all $s \in S$, there exists *unique* $s^* \in S$ such that $ss^*s = s$ and $s^*ss^* = s^*$.
- $ss^*, s^*s \in E(S)$ and so S has local units.
- Inverse semigroups are precisely $*$ -semigroups of partial isometries of Hilbert spaces.
- Graph inverse semigroups were considered first in the 70s by Ash and Hall for graphs without multiple edges.
- The general case was studied by Paterson in connection with graph C^* -algebras.
- P_G is an inverse semigroup.

Inverse semigroups

- S is an **inverse semigroup** if, for all $s \in S$, there exists *unique* $s^* \in S$ such that $ss^*s = s$ and $s^*ss^* = s^*$.
- $ss^*, s^*s \in E(S)$ and so S has local units.
- Inverse semigroups are precisely $*$ -semigroups of partial isometries of Hilbert spaces.
- Graph inverse semigroups were considered first in the 70s by Ash and Hall for graphs without multiple edges.
- The general case was studied by Paterson in connection with graph C^* -algebras.
- P_G is an inverse semigroup.

Theorem (BS)

Morita equivalent inverse semigroups have Morita equivalent universal, reduced and tight C^ -algebras.*

Markov-Dyck shifts

- Let $\Sigma_G = E \cup E^*$.

Markov-Dyck shifts

- Let $\Sigma_G = E \cup E^*$.
- Define $\pi: \Sigma_G^+ \rightarrow P_G$ by $\pi(e) = s_e$, $\pi(e^*) = s_e^*$.

Markov-Dyck shifts

- Let $\Sigma_G = E \cup E^*$.
- Define $\pi: \Sigma_G^+ \rightarrow P_G$ by $\pi(e) = s_e$, $\pi(e^*) = s_e^*$.
- Then $L_G = \pi^{-1}(P_G \setminus \{0\})$ is a factorial, prolongable language.

Markov-Dyck shifts

- Let $\Sigma_G = E \cup E^*$.
- Define $\pi: \Sigma_G^+ \rightarrow P_G$ by $\pi(e) = s_e$, $\pi(e^*) = s_e^*$.
- Then $L_G = \pi^{-1}(P_G \setminus \{0\})$ is a factorial, prolongable language.
- Let $D_G \subseteq (\Sigma_G)^{\mathbb{Z}}$ be the corresponding subshift.

Markov-Dyck shifts

- Let $\Sigma_G = E \cup E^*$.
- Define $\pi: \Sigma_G^+ \rightarrow P_G$ by $\pi(e) = s_e$, $\pi(e^*) = s_e^*$.
- Then $L_G = \pi^{-1}(P_G \setminus \{0\})$ is a factorial, prolongable language.
- Let $D_G \subseteq (\Sigma_G)^{\mathbb{Z}}$ be the corresponding subshift.
- It is called the **Markov-Dyck shift** associated to G .

Markov-Dyck shifts

- Let $\Sigma_G = E \cup E^*$.
- Define $\pi: \Sigma_G^+ \rightarrow P_G$ by $\pi(e) = s_e$, $\pi(e^*) = s_e^*$.
- Then $L_G = \pi^{-1}(P_G \setminus \{0\})$ is a factorial, prolongable language.
- Let $D_G \subseteq (\Sigma_G)^{\mathbb{Z}}$ be the corresponding subshift.
- It is called the **Markov-Dyck shift** associated to G .
- Markov-Dyck shifts have been studied by Krieger and his collaborators.

Markov-Dyck shifts

- Let $\Sigma_G = E \cup E^*$.
- Define $\pi: \Sigma_G^+ \rightarrow P_G$ by $\pi(e) = s_e$, $\pi(e^*) = s_e^*$.
- Then $L_G = \pi^{-1}(P_G \setminus \{0\})$ is a factorial, prolongable language.
- Let $D_G \subseteq (\Sigma_G)^{\mathbb{Z}}$ be the corresponding subshift.
- It is called the **Markov-Dyck shift** associated to G .
- Markov-Dyck shifts have been studied by Krieger and his collaborators.
- They have been classified by Krieger up to conjugacy when G is strongly connected.

Markov-Dyck shifts

- Let $\Sigma_G = E \cup E^*$.
- Define $\pi: \Sigma_G^+ \rightarrow P_G$ by $\pi(e) = s_e$, $\pi(e^*) = s_e^*$.
- Then $L_G = \pi^{-1}(P_G \setminus \{0\})$ is a factorial, prolongable language.
- Let $D_G \subseteq (\Sigma_G)^{\mathbb{Z}}$ be the corresponding subshift.
- It is called the **Markov-Dyck shift** associated to G .
- Markov-Dyck shifts have been studied by Krieger and his collaborators.
- They have been classified by Krieger up to conjugacy when G is strongly connected.
- Dyck shifts were classified by Matsumoto up to flow equivalence.

Flow equivalence classification

Theorem (Costa/BS)

Let G, H be finite graphs such that $\deg_{\text{out}}(v) \geq 1$ and $\deg_{\text{in}}(v) \neq 1$ for all vertices v . Then the following are equivalent.

Flow equivalence classification

Theorem (Costa/BS)

Let G, H be finite graphs such that $\deg_{\text{out}}(v) \geq 1$ and $\deg_{\text{in}}(v) \neq 1$ for all vertices v . Then the following are equivalent.

- 1 D_G and D_H are conjugate.

Flow equivalence classification

Theorem (Costa/BS)

Let G, H be finite graphs such that $\deg_{\text{out}}(v) \geq 1$ and $\deg_{\text{in}}(v) \neq 1$ for all vertices v . Then the following are equivalent.

- ❶ D_G and D_H are conjugate.
- ❷ D_G and D_H are flow equivalent.

Flow equivalence classification

Theorem (Costa/BS)

Let G, H be finite graphs such that $\deg_{\text{out}}(v) \geq 1$ and $\deg_{\text{in}}(v) \neq 1$ for all vertices v . Then the following are equivalent.

- ❶ D_G and D_H are conjugate.
- ❷ D_G and D_H are flow equivalent.
- ❸ G and H are isomorphic.

Flow equivalence classification

Theorem (Costa/BS)

Let G, H be finite graphs such that $\deg_{\text{out}}(v) \geq 1$ and $\deg_{\text{in}}(v) \neq 1$ for all vertices v . Then the following are equivalent.

- ❶ D_G and D_H are conjugate.
- ❷ D_G and D_H are flow equivalent.
- ❸ G and H are isomorphic.

- An analogous result holds for Markov-Motzkin shifts.

Flow equivalence classification

Theorem (Costa/BS)

Let G, H be finite graphs such that $\deg_{\text{out}}(v) \geq 1$ and $\deg_{\text{in}}(v) \neq 1$ for all vertices v . Then the following are equivalent.

- ❶ D_G and D_H are conjugate.
- ❷ D_G and D_H are flow equivalent.
- ❸ G and H are isomorphic.

- An analogous result holds for Markov-Motzkin shifts.
- In this case, sinks are allowed.

G is determined by $\mathbb{K}(P_G)$

- Under our assumptions, we have P_G is the syntactic semigroup of D_G . (And the same for H .)

G is determined by $\mathbb{K}(P_G)$

- Under our assumptions, we have P_G is the syntactic semigroup of D_G . (And the same for H .)

Theorem (BS)

P_G is Morita equivalent to P_H if and only if $G = H$.

G is determined by $\mathbb{K}(P_G)$

- Under our assumptions, we have P_G is the syntactic semigroup of D_G . (And the same for H .)

Theorem (BS)

P_G is Morita equivalent to P_H if and only if $G = H$.

- If S is a semigroup with 0, then the object 0 of $\mathbb{K}(S)$ is the unique object which is both initial and terminal.

G is determined by $\mathbb{K}(P_G)$

- Under our assumptions, we have P_G is the syntactic semigroup of D_G . (And the same for H .)

Theorem (BS)

P_G is Morita equivalent to P_H if and only if $G = H$.

- If S is a semigroup with 0, then the object 0 of $\mathbb{K}(S)$ is the unique object which is both initial and terminal.
- Let $\mathbb{K}_0(S)$ be the full subcategory of $\mathbb{K}(S)$ on the nonzero elements of $E(S)$. (Call it the **proper part** of $\mathbb{K}(S)$.)

G is determined by $\mathbb{K}(P_G)$

- Under our assumptions, we have P_G is the syntactic semigroup of D_G . (And the same for H .)

Theorem (BS)

P_G is Morita equivalent to P_H if and only if $G = H$.

- If S is a semigroup with 0, then the object 0 of $\mathbb{K}(S)$ is the unique object which is both initial and terminal.
- Let $\mathbb{K}_0(S)$ be the full subcategory of $\mathbb{K}(S)$ on the nonzero elements of $E(S)$. (Call it the **proper part** of $\mathbb{K}(S)$.)
- It follows that if S and T have zeroes, then $\mathbb{K}(S)$ is equivalent to $\mathbb{K}(T)$ iff their proper parts are equivalent.

Split epimorphisms

- A morphism $f: c \rightarrow d$ of a category admitting a section s (so $fs = 1_d$) is called a **split epimorphism**.

Split epimorphisms

- A morphism $f: c \rightarrow d$ of a category admitting a section s (so $fs = 1_d$) is called a **split epimorphism**.
- Clearly, split epimorphisms are preserved by all functors.

Split epimorphisms

- A morphism $f: c \rightarrow d$ of a category admitting a section s (so $fs = 1_d$) is called a **split epimorphism**.
- Clearly, split epimorphisms are preserved by all functors.
- If S is a semigroup with 0, let $\mathbb{L}(S)$ be the subcategory of $\mathbb{K}_0(S)$ whose arrows are the split epimorphisms.

Split epimorphisms

- A morphism $f: c \rightarrow d$ of a category admitting a section s (so $fs = 1_d$) is called a **split epimorphism**.
- Clearly, split epimorphisms are preserved by all functors.
- If S is a semigroup with 0, let $\mathbb{L}(S)$ be the subcategory of $\mathbb{K}_0(S)$ whose arrows are the split epimorphisms.
- If $\mathbb{K}_0(S)$ is equivalent to $\mathbb{K}_0(T)$, then $\mathbb{L}(S)$ is equivalent to $\mathbb{L}(T)$.

Split epimorphisms

- A morphism $f: c \rightarrow d$ of a category admitting a section s (so $fs = 1_d$) is called a **split epimorphism**.
- Clearly, split epimorphisms are preserved by all functors.
- If S is a semigroup with 0, let $\mathbb{L}(S)$ be the subcategory of $\mathbb{K}_0(S)$ whose arrows are the split epimorphisms.
- If $\mathbb{K}_0(S)$ is equivalent to $\mathbb{K}_0(T)$, then $\mathbb{L}(S)$ is equivalent to $\mathbb{L}(T)$.
- One easily checks that $\mathbb{L}(P_G)$ is equivalent to the free category G^* on G .

Split epimorphisms

- A morphism $f: c \rightarrow d$ of a category admitting a section s (so $fs = 1_d$) is called a **split epimorphism**.
- Clearly, split epimorphisms are preserved by all functors.
- If S is a semigroup with 0, let $\mathbb{L}(S)$ be the subcategory of $\mathbb{K}_0(S)$ whose arrows are the split epimorphisms.
- If $\mathbb{K}_0(S)$ is equivalent to $\mathbb{K}_0(T)$, then $\mathbb{L}(S)$ is equivalent to $\mathbb{L}(T)$.
- One easily checks that $\mathbb{L}(P_G)$ is equivalent to the free category G^* on G .
- Two free categories are equivalent iff the graphs are isomorphic.

Split epimorphisms

- A morphism $f: c \rightarrow d$ of a category admitting a section s (so $fs = 1_d$) is called a **split epimorphism**.
- Clearly, split epimorphisms are preserved by all functors.
- If S is a semigroup with 0, let $\mathbb{L}(S)$ be the subcategory of $\mathbb{K}_0(S)$ whose arrows are the split epimorphisms.
- If $\mathbb{K}_0(S)$ is equivalent to $\mathbb{K}_0(T)$, then $\mathbb{L}(S)$ is equivalent to $\mathbb{L}(T)$.
- One easily checks that $\mathbb{L}(P_G)$ is equivalent to the free category G^* on G .
- Two free categories are equivalent iff the graphs are isomorphic.
- So $\mathbb{K}(P_G)$ equivalent to $\mathbb{K}(P_H)$ implies $G \cong H$.

Shifts and automata

- An **automaton** \mathcal{A} over the alphabet A is a graph with edges labeled by elements of A .

Shifts and automata

- An **automaton** \mathcal{A} over the alphabet A is a graph with edges labeled by elements of A .
- $\mathcal{X}_{\mathcal{A}}$ denotes the shift of all labels of bi-infinite paths in \mathcal{A} .


Shifts and automata

- An **automaton** \mathcal{A} over the alphabet A is a graph with edges labeled by elements of A .
- $\mathcal{X}_{\mathcal{A}}$ denotes the shift of all labels of bi-infinite paths in \mathcal{A} .
- $\mathcal{X}_{\mathcal{A}}$ is sofic if \mathcal{A} is finite.

Shifts and automata

- An **automaton** \mathcal{A} over the alphabet A is a graph with edges labeled by elements of A .
- $\mathcal{X}_{\mathcal{A}}$ denotes the shift of all labels of bi-infinite paths in \mathcal{A} .
- $\mathcal{X}_{\mathcal{A}}$ is sofic if \mathcal{A} is finite.


Example (Even shift)



Shifts and automata

- An **automaton** \mathcal{A} over the alphabet A is a graph with edges labeled by elements of A .
- $\mathcal{X}_{\mathcal{A}}$ denotes the shift of all labels of bi-infinite paths in \mathcal{A} .
- $\mathcal{X}_{\mathcal{A}}$ is sofic if \mathcal{A} is finite.

Example (Even shift)




- \mathcal{A} is **deterministic** if there is *at most one* edge leaving any vertex with a given label $a \in A$.

Shifts and automata

- An **automaton** \mathcal{A} over the alphabet A is a graph with edges labeled by elements of A .
- $\mathcal{X}_{\mathcal{A}}$ denotes the shift of all labels of bi-infinite paths in \mathcal{A} .
- $\mathcal{X}_{\mathcal{A}}$ is sofic if \mathcal{A} is finite.

Example (Even shift)




- \mathcal{A} is **deterministic** if there is *at most one* edge leaving any vertex with a given label $a \in A$.
- The Krieger automaton for a shift is deterministic.

Shifts and automata

- An **automaton** \mathcal{A} over the alphabet A is a graph with edges labeled by elements of A .
- $\mathcal{X}_{\mathcal{A}}$ denotes the shift of all labels of bi-infinite paths in \mathcal{A} .
- $\mathcal{X}_{\mathcal{A}}$ is sofic if \mathcal{A} is finite.

Example (Even shift)



- \mathcal{A} is **deterministic** if there is *at most one* edge leaving any vertex with a given label $a \in A$.
- The Krieger automaton for a shift is deterministic.
- The Fischer cover for an irreducible sofic shift is deterministic.

Transition semigroups

- \mathcal{A} is a deterministic automaton over A with state set Q .

Transition semigroups

- \mathcal{A} is a deterministic automaton over A with state set Q .
- For each $a \in A$, define a partial map $f_a: Q \rightarrow Q$ by:

Transition semigroups

- \mathcal{A} is a deterministic automaton over A with state set Q .
- For each $a \in A$, define a partial map $f_a: Q \rightarrow Q$ by:
 - $qf_a = \emptyset$ if there is no edge from q labelled a ;


Transition semigroups

- \mathcal{A} is a deterministic automaton over A with state set Q .
- For each $a \in A$, define a partial map $f_a: Q \rightarrow Q$ by:
 - $qf_a = \emptyset$ if there is no edge from q labelled a ;
 - $qf_a = q'$ if there is an edge $q \xrightarrow{a} q'$.

Transition semigroups

- \mathcal{A} is a deterministic automaton over A with state set Q .
- For each $a \in A$, define a partial map $f_a: Q \rightarrow Q$ by:
 - $qf_a = \emptyset$ if there is no edge from q labelled a ;
 - $qf_a = q'$ if there is an edge $q \xrightarrow{a} q'$.


Example (Even shift)



Transition semigroups

- \mathcal{A} is a deterministic automaton over A with state set Q .
- For each $a \in A$, define a partial map $f_a: Q \rightarrow Q$ by:
 - $qf_a = \emptyset$ if there is no edge from q labelled a ;
 - $qf_a = q'$ if there is an edge $q \xrightarrow{a} q'$.

Example (Even shift)




$$f_a = \begin{pmatrix} p & q \\ q & p \end{pmatrix} \text{ and } f_b = \begin{pmatrix} p & q \\ p & \emptyset \end{pmatrix}$$

Transition semigroups

- \mathcal{A} is a deterministic automaton over A with state set Q .
- For each $a \in A$, define a partial map $f_a: Q \rightarrow Q$ by:
 - $qf_a = \emptyset$ if there is no edge from q labelled a ;
 - $qf_a = q'$ if there is an edge $q \xrightarrow{a} q'$.

Example (Even shift)



$$f_a = \begin{pmatrix} p & q \\ q & p \end{pmatrix} \text{ and } f_b = \begin{pmatrix} p & q \\ p & \emptyset \end{pmatrix}$$

- The semigroup $S(\mathcal{A}) = \langle \{f_a \mid a \in A\} \rangle$ is called the **transition semigroup** of \mathcal{A} .

The syntactic semigroup as a transition semigroup

- If \mathcal{A} is the Krieger automaton of a shift \mathcal{X} , then $S(\mathcal{A}) \cong S(\mathcal{X})$.

The syntactic semigroup as a transition semigroup

- If \mathcal{A} is the Krieger automaton of a shift \mathcal{X} , then $S(\mathcal{A}) \cong S(\mathcal{X})$.
- The analogous statement holds for the Fischer cover of an irreducible sofic shift.

The syntactic semigroup as a transition semigroup

- If \mathcal{A} is the Krieger automaton of a shift \mathcal{X} , then $S(\mathcal{A}) \cong S(\mathcal{X})$.
- The analogous statement holds for the Fischer cover of an irreducible sofic shift.
- The action of $S(\mathcal{X})$ on the states of its Krieger (or Fischer) cover gives a functor $F: \mathbb{K}(\mathcal{X}) \rightarrow \mathbf{Set}$.

The syntactic semigroup as a transition semigroup

- If \mathcal{A} is the Krieger automaton of a shift \mathcal{X} , then $S(\mathcal{A}) \cong S(\mathcal{X})$.
- The analogous statement holds for the Fischer cover of an irreducible sofic shift.
- The action of $S(\mathcal{X})$ on the states of its Krieger (or Fischer) cover gives a functor $F: \mathbb{K}(\mathcal{X}) \rightarrow \mathbf{Set}$.
- If \mathcal{X} is flow equivalent to \mathcal{Y} , then this functor is 'remembered' by the equivalence of $\mathbb{K}(\mathcal{X})$ with $\mathbb{K}(\mathcal{Y})$ (see Alfredo's talk).

Bipartite automata

- An automaton \mathcal{A} over A is **bipartite** if there are partitions $Q = Q_1 \uplus Q_2$ of the states and $A = A_1 \uplus A_2$ of the alphabet such that:

Bipartite automata

- An automaton \mathcal{A} over A is **bipartite** if there are partitions $Q = Q_1 \uplus Q_2$ of the states and $A = A_1 \uplus A_2$ of the alphabet such that:
 - all edges labeled by A_1 go from Q_1 to Q_2

Bipartite automata

- An automaton \mathcal{A} over A is **bipartite** if there are partitions $Q = Q_1 \uplus Q_2$ of the states and $A = A_1 \uplus A_2$ of the alphabet such that:
 - all edges labeled by A_1 go from Q_1 to Q_2
 - all edges labeled by A_2 go from Q_2 to Q_1 .

Bipartite automata

- An automaton \mathcal{A} over A is **bipartite** if there are partitions $Q = Q_1 \uplus Q_2$ of the states and $A = A_1 \uplus A_2$ of the alphabet such that:
 - all edges labeled by A_1 go from Q_1 to Q_2
 - all edges labeled by A_2 go from Q_2 to Q_1 .
- Let \mathcal{A}_1 be the automaton over $A_1 A_2$ with state set Q_1 obtained by turning each path of length 2 from Q_1 to itself into an edge.

Bipartite automata

- An automaton \mathcal{A} over A is **bipartite** if there are partitions $Q = Q_1 \uplus Q_2$ of the states and $A = A_1 \uplus A_2$ of the alphabet such that:
 - all edges labeled by A_1 go from Q_1 to Q_2
 - all edges labeled by A_2 go from Q_2 to Q_1 .
- Let \mathcal{A}_1 be the automaton over A_1A_2 with state set Q_1 obtained by turning each path of length 2 from Q_1 to itself into an edge.
- Define \mathcal{A}_2 with alphabet A_2A_1 and states Q_2 analogously.

Bipartite automata

- An automaton \mathcal{A} over A is **bipartite** if there are partitions $Q = Q_1 \uplus Q_2$ of the states and $A = A_1 \uplus A_2$ of the alphabet such that:
 - all edges labeled by A_1 go from Q_1 to Q_2
 - all edges labeled by A_2 go from Q_2 to Q_1 .
- Let \mathcal{A}_1 be the automaton over A_1A_2 with state set Q_1 obtained by turning each path of length 2 from Q_1 to itself into an edge.
- Define \mathcal{A}_2 with alphabet A_2A_1 and states Q_2 analogously.
- We call $\mathcal{A}_1, \mathcal{A}_2$ the **components** of \mathcal{A} .

Nasu's theorem

- Write $\mathcal{A} \sim \mathcal{B}$ if there is a bipartite automaton with components \mathcal{A}, \mathcal{B} .

Nasu's theorem

- Write $\mathcal{A} \sim \mathcal{B}$ if there is a bipartite automaton with components \mathcal{A}, \mathcal{B} .
- Let \simeq be the equivalence relation on automata generated by \sim .

Nasu's theorem

- Write $\mathcal{A} \sim \mathcal{B}$ if there is a bipartite automaton with components \mathcal{A}, \mathcal{B} .
- Let \simeq be the equivalence relation on automata generated by \sim .

Theorem (Nasu 1986)

Let $\mathcal{X}_1, \mathcal{X}_2$ be sofic shifts with Krieger automata $\mathcal{A}_1, \mathcal{A}_2$, respectively. Then the following are equivalent.

Nasu's theorem

- Write $\mathcal{A} \sim \mathcal{B}$ if there is a bipartite automaton with components \mathcal{A}, \mathcal{B} .
- Let \simeq be the equivalence relation on automata generated by \sim .

Theorem (Nasu 1986)

Let $\mathcal{X}_1, \mathcal{X}_2$ be sofic shifts with Krieger automata $\mathcal{A}_1, \mathcal{A}_2$, respectively. Then the following are equivalent.

1. \mathcal{X}_1 and \mathcal{X}_2 are conjugate.

Nasu's theorem

- Write $\mathcal{A} \sim \mathcal{B}$ if there is a bipartite automaton with components \mathcal{A}, \mathcal{B} .
- Let \simeq be the equivalence relation on automata generated by \sim .

Theorem (Nasu 1986)

Let $\mathcal{X}_1, \mathcal{X}_2$ be sofic shifts with Krieger automata $\mathcal{A}_1, \mathcal{A}_2$, respectively. Then the following are equivalent.

- 1 \mathcal{X}_1 and \mathcal{X}_2 are conjugate.
- 2 $\mathcal{A}_1 \simeq \mathcal{A}_2$ (via finite bipartite automata).

Nasu's theorem

- Write $\mathcal{A} \sim \mathcal{B}$ if there is a bipartite automaton with components \mathcal{A}, \mathcal{B} .
- Let \simeq be the equivalence relation on automata generated by \sim .

Theorem (Nasu 1986)

Let $\mathcal{X}_1, \mathcal{X}_2$ be sofic shifts with Krieger automata $\mathcal{A}_1, \mathcal{A}_2$, respectively. Then the following are equivalent.

- 1 \mathcal{X}_1 and \mathcal{X}_2 are conjugate.
 - 2 $\mathcal{A}_1 \simeq \mathcal{A}_2$ (via finite bipartite automata).
- Looking at Nasu's paper it seems $1 \implies 2$ (without the finiteness condition) does not require the sofic hypothesis.

Nasu's theorem

- Write $\mathcal{A} \sim \mathcal{B}$ if there is a bipartite automaton with components \mathcal{A}, \mathcal{B} .
- Let \simeq be the equivalence relation on automata generated by \sim .

Theorem (Nasu 1986)

Let $\mathcal{X}_1, \mathcal{X}_2$ be sofic shifts with Krieger automata $\mathcal{A}_1, \mathcal{A}_2$, respectively. Then the following are equivalent.

- 1 \mathcal{X}_1 and \mathcal{X}_2 are conjugate.
 - 2 $\mathcal{A}_1 \simeq \mathcal{A}_2$ (via finite bipartite automata).
- Looking at Nasu's paper it seems $1 \implies 2$ (without the finiteness condition) does not require the sofic hypothesis.
 - If so, the following is a simpler proof of our main result.

Conjugacy invariance of the Karoubi envelope

- In light of Nasu's result, proving conjugacy invariance of the Karoubi envelope reduces to the following result.

Conjugacy invariance of the Karoubi envelope

- In light of Nasu's result, proving conjugacy invariance of the Karoubi envelope reduces to the following result.

Theorem (Costa/BS)

Let \mathcal{A} be a bipartite automaton with components $\mathcal{A}_1, \mathcal{A}_2$. Let S, S_1, S_2 be the transition semigroups of $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$, respectively.

Conjugacy invariance of the Karoubi envelope

- In light of Nasu's result, proving conjugacy invariance of the Karoubi envelope reduces to the following result.

Theorem (Costa/BS)

Let \mathcal{A} be a bipartite automaton with components $\mathcal{A}_1, \mathcal{A}_2$. Let S, S_1, S_2 be the transition semigroups of $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$, respectively. Then S, S_1, S_2 are all Morita equivalent up to local units.

Conjugacy invariance of the Karoubi envelope

- In light of Nasu's result, proving conjugacy invariance of the Karoubi envelope reduces to the following result.

Theorem (Costa/BS)

Let \mathcal{A} be a bipartite automaton with components $\mathcal{A}_1, \mathcal{A}_2$. Let S, S_1, S_2 be the transition semigroups of $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$, respectively. Then S, S_1, S_2 are all Morita equivalent up to local units. Moreover, the equivalence is “action-preserving.”

Enlargements

- Let $S \leq T$ be semigroups with local units.

Enlargements

- Let $S \leq T$ be semigroups with local units.
- T is an **enlargement** of S if there is a set of idempotents $E \subseteq T$ such that:

Enlargements

- Let $S \leq T$ be semigroups with local units.
- T is an **enlargement** of S if there is a set of idempotents $E \subseteq T$ such that:
 - $TET = T$;

Enlargements

- Let $S \leq T$ be semigroups with local units.
- T is an **enlargement** of S if there is a set of idempotents $E \subseteq T$ such that:
 - $TET = T$;
 - $ETE = S$.

Enlargements

- Let $S \leq T$ be semigroups with local units.
- T is an **enlargement** of S if there is a set of idempotents $E \subseteq T$ such that:
 - $TET = T$;
 - $ETE = S$.
- Equivalently, $TST = T$ and $STS = S$.

Enlargements

- Let $S \leq T$ be semigroups with local units.
- T is an **enlargement** of S if there is a set of idempotents $E \subseteq T$ such that:
 - $TET = T$;
 - $ETE = S$.
- Equivalently, $TST = T$ and $STS = S$.
- In this setup, the bisets ST and TS give the Morita context.

Enlargements

- Let $S \leq T$ be semigroups with local units.
- T is an **enlargement** of S if there is a set of idempotents $E \subseteq T$ such that:
 - $TET = T$;
 - $ETE = S$.
- Equivalently, $TST = T$ and $STS = S$.
- In this setup, the bisets ST and TS give the Morita context.

Theorem (Lawson)

Two semigroups with local units are Morita equivalent iff they have a common enlargement.

Completion of the proof sketch

- Let \mathcal{A} be a bipartite automata with components $\mathcal{A}_1, \mathcal{A}_2$.

Completion of the proof sketch

- Let \mathcal{A} be a bipartite automata with components $\mathcal{A}_1, \mathcal{A}_2$.
- Let S, S_1, S_2 be the transition semigroups of $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$, respectively.

Completion of the proof sketch

- Let \mathcal{A} be a bipartite automata with components $\mathcal{A}_1, \mathcal{A}_2$.
- Let S, S_1, S_2 be the transition semigroups of $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$, respectively.
- One checks that $LU(S)$ is a common enlargement of $LU(S_1), LU(S_2)$.

Completion of the proof sketch

- Let \mathcal{A} be a bipartite automata with components $\mathcal{A}_1, \mathcal{A}_2$.
- Let S, S_1, S_2 be the transition semigroups of $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$, respectively.
- One checks that $LU(S)$ is a common enlargement of $LU(S_1), LU(S_2)$.
- This follows easily from the following observations of Béal, Berstel, Eilers and Perrin (stated in the finite case).

Completion of the proof sketch

- Let \mathcal{A} be a bipartite automata with components $\mathcal{A}_1, \mathcal{A}_2$.
- Let S, S_1, S_2 be the transition semigroups of $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$, respectively.
- One checks that $LU(S)$ is a common enlargement of $LU(S_1), LU(S_2)$.
- This follows easily from the following observations of Béal, Berstel, Eilers and Perrin (stated in the finite case).
- $E(S) \setminus \{0\} = (E(S_1) \setminus \{0\}) \uplus (E(S_2) \setminus \{0\})$;

Completion of the proof sketch

- Let \mathcal{A} be a bipartite automata with components $\mathcal{A}_1, \mathcal{A}_2$.
- Let S, S_1, S_2 be the transition semigroups of $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$, respectively.
- One checks that $LU(S)$ is a common enlargement of $LU(S_1), LU(S_2)$.
- This follows easily from the following observations of Béal, Berstel, Eilers and Perrin (stated in the finite case).
- $E(S) \setminus \{0\} = (E(S_1) \setminus \{0\}) \uplus (E(S_2) \setminus \{0\})$;
- Each idempotent of S_1 is \mathcal{D} -equivalent to an idempotent of S_2 , and vice versa.

Completion of the proof sketch

- Let \mathcal{A} be a bipartite automata with components $\mathcal{A}_1, \mathcal{A}_2$.
- Let S, S_1, S_2 be the transition semigroups of $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$, respectively.
- One checks that $LU(S)$ is a common enlargement of $LU(S_1), LU(S_2)$.
- This follows easily from the following observations of Béal, Berstel, Eilers and Perrin (stated in the finite case).
- $E(S) \setminus \{0\} = (E(S_1) \setminus \{0\}) \uplus (E(S_2) \setminus \{0\})$;
- Each idempotent of S_1 is \mathcal{D} -equivalent to an idempotent of S_2 , and vice versa.
- The equivalence takes the $LU(S_1)$ -set $Q_1 \otimes LU(S_1)$ to the $LU(S_2)$ -set $Q_2 \otimes LU(S_2)$.

The end

Thank you for your attention!