

# Matrix mortality and the Pin-Černý Conjecture

Jorge Almeida<sup>1</sup> Benjamin Steinberg<sup>2</sup>

<sup>1</sup>University of Porto, <sup>2</sup>Carleton University

[bsteinbg@math.carleton.ca](mailto:bsteinbg@math.carleton.ca)

<http://www.mathstat.carleton.ca/~bsteinbg>

DLT July 3, 2009

# Synchronizing automata

- By an **automaton**  $\mathcal{A} = (Q, \Sigma)$ , we understand a complete deterministic automaton with state set  $Q$ , input alphabet  $\Sigma$  and no initial or final states.
- If  $w \in \Sigma^*$ , then the **rank** of  $w$  is  $\text{rk}(w) = |Qw|$ .
- If  $\text{rk}(w) = 1$ , then  $w$  is called a **reset word**.
- Define  $\text{rk}(\mathcal{A}) = \min\{\text{rk}(w) \mid w \in \Sigma^*\}$  (Pin).
- $\mathcal{A}$  is synchronizing if  $\text{rk}(\mathcal{A}) = 1$ , i.e., it admits a reset word.

## Conjecture (Černý-Pin)

*An automaton  $\mathcal{A}$  of rank  $r$  admits a word  $w$  of length at most  $(n - r)^2$  with  $\text{rk}(w) = r$ .*

- The case  $r = 1$  is due to Černý; the more general conjecture is a variation on an earlier conjecture of Pin.

# Synchronizing automata

- By an **automaton**  $\mathcal{A} = (Q, \Sigma)$ , we understand a complete deterministic automaton with state set  $Q$ , input alphabet  $\Sigma$  and no initial or final states.
- If  $w \in \Sigma^*$ , then the **rank** of  $w$  is  $\text{rk}(w) = |Qw|$ .
- If  $\text{rk}(w) = 1$ , then  $w$  is called a **reset word**.
- Define  $\text{rk}(\mathcal{A}) = \min\{\text{rk}(w) \mid w \in \Sigma^*\}$  (Pin).
- $\mathcal{A}$  is **synchronizing** if  $\text{rk}(\mathcal{A}) = 1$ , i.e., it admits a reset word.

## Conjecture (Černý-Pin)

*An automaton  $\mathcal{A}$  of rank  $r$  admits a word  $w$  of length at most  $(n - r)^2$  with  $\text{rk}(w) = r$ .*

- The case  $r = 1$  is due to Černý; the more general conjecture is a variation on an earlier conjecture of Pin.

# Synchronizing automata

- By an **automaton**  $\mathcal{A} = (Q, \Sigma)$ , we understand a complete deterministic automaton with state set  $Q$ , input alphabet  $\Sigma$  and no initial or final states.
- If  $w \in \Sigma^*$ , then the **rank** of  $w$  is  $\text{rk}(w) = |Qw|$ .
- If  $\text{rk}(w) = 1$ , then  $w$  is called a **reset word**.
- Define  $\text{rk}(\mathcal{A}) = \min\{\text{rk}(w) \mid w \in \Sigma^*\}$  (Pin).
- $\mathcal{A}$  is **synchronizing** if  $\text{rk}(\mathcal{A}) = 1$ , i.e., it admits a reset word.

## Conjecture (Černý-Pin)

*An automaton  $\mathcal{A}$  of rank  $r$  admits a word  $w$  of length at most  $(n - r)^2$  with  $\text{rk}(w) = r$ .*

- The case  $r = 1$  is due to Černý; the more general conjecture is a variation on an earlier conjecture of Pin.

# Synchronizing automata

- By an **automaton**  $\mathcal{A} = (Q, \Sigma)$ , we understand a complete deterministic automaton with state set  $Q$ , input alphabet  $\Sigma$  and no initial or final states.
- If  $w \in \Sigma^*$ , then the **rank** of  $w$  is  $\text{rk}(w) = |Qw|$ .
- If  $\text{rk}(w) = 1$ , then  $w$  is called a **reset word**.
- Define  $\text{rk}(\mathcal{A}) = \min\{\text{rk}(w) \mid w \in \Sigma^*\}$  (Pin).
- $\mathcal{A}$  is **synchronizing** if  $\text{rk}(\mathcal{A}) = 1$ , i.e., it admits a reset word.

## Conjecture (Černý-Pin)

*An automaton  $\mathcal{A}$  of rank  $r$  admits a word  $w$  of length at most  $(n - r)^2$  with  $\text{rk}(w) = r$ .*

- The case  $r = 1$  is due to Černý; the more general conjecture is a variation on an earlier conjecture of Pin.

# Synchronizing automata

- By an **automaton**  $\mathcal{A} = (Q, \Sigma)$ , we understand a complete deterministic automaton with state set  $Q$ , input alphabet  $\Sigma$  and no initial or final states.
- If  $w \in \Sigma^*$ , then the **rank** of  $w$  is  $\text{rk}(w) = |Qw|$ .
- If  $\text{rk}(w) = 1$ , then  $w$  is called a **reset word**.
- Define  $\text{rk}(\mathcal{A}) = \min\{\text{rk}(w) \mid w \in \Sigma^*\}$  (Pin).
- $\mathcal{A}$  is **synchronizing** if  $\text{rk}(\mathcal{A}) = 1$ , i.e., it admits a reset word.

## Conjecture (Černý-Pin)

*An automaton  $\mathcal{A}$  of rank  $r$  admits a word  $w$  of length at most  $(n - r)^2$  with  $\text{rk}(w) = r$ .*

- The case  $r = 1$  is due to Černý; the more general conjecture is a variation on an earlier conjecture of Pin.

# Synchronizing automata

- By an **automaton**  $\mathcal{A} = (Q, \Sigma)$ , we understand a complete deterministic automaton with state set  $Q$ , input alphabet  $\Sigma$  and no initial or final states.
- If  $w \in \Sigma^*$ , then the **rank** of  $w$  is  $\text{rk}(w) = |Qw|$ .
- If  $\text{rk}(w) = 1$ , then  $w$  is called a **reset word**.
- Define  $\text{rk}(\mathcal{A}) = \min\{\text{rk}(w) \mid w \in \Sigma^*\}$  (Pin).
- $\mathcal{A}$  is **synchronizing** if  $\text{rk}(\mathcal{A}) = 1$ , i.e., it admits a reset word.

## Conjecture (Černý-Pin)

*An automaton  $\mathcal{A}$  of rank  $r$  admits a word  $w$  of length at most  $(n - r)^2$  with  $\text{rk}(w) = r$ .*

- The case  $r = 1$  is due to Černý; the more general conjecture is a variation on an earlier conjecture of Pin.

# Synchronizing automata

- By an **automaton**  $\mathcal{A} = (Q, \Sigma)$ , we understand a complete deterministic automaton with state set  $Q$ , input alphabet  $\Sigma$  and no initial or final states.
- If  $w \in \Sigma^*$ , then the **rank** of  $w$  is  $\text{rk}(w) = |Qw|$ .
- If  $\text{rk}(w) = 1$ , then  $w$  is called a **reset word**.
- Define  $\text{rk}(\mathcal{A}) = \min\{\text{rk}(w) \mid w \in \Sigma^*\}$  (Pin).
- $\mathcal{A}$  is **synchronizing** if  $\text{rk}(\mathcal{A}) = 1$ , i.e., it admits a reset word.

## Conjecture (Černý-Pin)

*An automaton  $\mathcal{A}$  of rank  $r$  admits a word  $w$  of length at most  $(n - r)^2$  with  $\text{rk}(w) = r$ .*

- The case  $r = 1$  is due to Černý; the more general conjecture is a variation on an earlier conjecture of Pin.




# Černý's examples

- Černý showed that the shortest length reset word for the  $n$ -state synchronizing automaton with transitions

$$a = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 2 & 3 & 4 & \cdots & 1 \end{pmatrix}, \quad b = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 2 & 2 & 3 & \cdots & n \end{pmatrix}$$

is  $(n - 1)^2$ .

- The Černý automaton for  $n = 4$ :



- The word  $b(a^3b)^2$  resets to state 2.


# Černý's examples

- Černý showed that the shortest length reset word for the  $n$ -state synchronizing automaton with transitions

$$a = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 2 & 3 & 4 & \cdots & 1 \end{pmatrix}, \quad b = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 2 & 2 & 3 & \cdots & n \end{pmatrix}$$

is  $(n - 1)^2$ .

- The Černý automaton for  $n = 4$ :



- The word  $b(a^3b)^2$  resets to state 2.


# Černý's examples

- Černý showed that the shortest length reset word for the  $n$ -state synchronizing automaton with transitions

$$a = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 2 & 3 & 4 & \cdots & 1 \end{pmatrix}, \quad b = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 2 & 2 & 3 & \cdots & n \end{pmatrix}$$

is  $(n - 1)^2$ .

- The Černý automaton for  $n = 4$ :



- The word  $b(a^3b)^2$  resets to state 2.

- It is straightforward to obtain a cubic upper bound of  $\frac{n^3-n}{3}$  on reset words for synchronizing automata.
- The best known upper bound for the synchronizing case is  $\frac{n^3-n}{6}$ , which was proved by Pin modulo an extremal set theory result of Frankl.
- Improving a bound by a factor of 2 can be hard work!
- The lower bound of  $(n-r)^2$  for rank  $r$  is due to Pin.
- Pin also has an analogous cubic upper bound for rank  $r$ .
- Probabilistically speaking, all automata are synchronizing with reset word of length at most  $2n$ .
- The remainder of the Černý literature consists of a vast array of special, but interesting, cases.
- There is no time to survey the whole literature here.

- It is straightforward to obtain a cubic upper bound of  $\frac{n^3-n}{3}$  on reset words for synchronizing automata.
- The best known upper bound for the synchronizing case is  $\frac{n^3-n}{6}$ , which was proved by Pin modulo an extremal set theory result of Frankl.
- Improving a bound by a factor of 2 can be hard work!
- The lower bound of  $(n-r)^2$  for rank  $r$  is due to Pin.
- Pin also has an analogous cubic upper bound for rank  $r$ .
- Probabilistically speaking, all automata are synchronizing with reset word of length at most  $2n$ .
- The remainder of the Černý literature consists of a vast array of special, but interesting, cases.
- There is no time to survey the whole literature here.

- It is straightforward to obtain a cubic upper bound of  $\frac{n^3-n}{3}$  on reset words for synchronizing automata.
- The best known upper bound for the synchronizing case is  $\frac{n^3-n}{6}$ , which was proved by Pin modulo an extremal set theory result of Frankl.
- Improving a bound by a factor of 2 can be hard work!
- The lower bound of  $(n-r)^2$  for rank  $r$  is due to Pin.
- Pin also has an analogous cubic upper bound for rank  $r$ .
- Probabilistically speaking, all automata are synchronizing with reset word of length at most  $2n$ .
- The remainder of the Černý literature consists of a vast array of special, but interesting, cases.
- There is no time to survey the whole literature here.

# Pin's bounds

- It is straightforward to obtain a cubic upper bound of  $\frac{n^3-n}{3}$  on reset words for synchronizing automata.
- The best known upper bound for the synchronizing case is  $\frac{n^3-n}{6}$ , which was proved by Pin modulo an extremal set theory result of Frankl.
- Improving a bound by a factor of 2 can be hard work!
- The lower bound of  $(n-r)^2$  for rank  $r$  is due to Pin.
- Pin also has an analogous cubic upper bound for rank  $r$ .
- Probabilistically speaking, all automata are synchronizing with reset word of length at most  $2n$ .
- The remainder of the Černý literature consists of a vast array of special, but interesting, cases.
- There is no time to survey the whole literature here.

# Pin's bounds

- It is straightforward to obtain a cubic upper bound of  $\frac{n^3-n}{3}$  on reset words for synchronizing automata.
- The best known upper bound for the synchronizing case is  $\frac{n^3-n}{6}$ , which was proved by Pin modulo an extremal set theory result of Frankl.
- Improving a bound by a factor of 2 can be hard work!
- The lower bound of  $(n-r)^2$  for rank  $r$  is due to Pin.
- Pin also has an analogous cubic upper bound for rank  $r$ .
- Probabilistically speaking, all automata are synchronizing with reset word of length at most  $2n$ .
- The remainder of the Černý literature consists of a vast array of special, but interesting, cases.
- There is no time to survey the whole literature here.



# Pin's bounds

- It is straightforward to obtain a cubic upper bound of  $\frac{n^3-n}{3}$  on reset words for synchronizing automata.
- The best known upper bound for the synchronizing case is  $\frac{n^3-n}{6}$ , which was proved by Pin modulo an extremal set theory result of Frankl.
- Improving a bound by a factor of 2 can be hard work!
- The lower bound of  $(n-r)^2$  for rank  $r$  is due to Pin.
- Pin also has an analogous cubic upper bound for rank  $r$ .
- Probabilistically speaking, all automata are synchronizing with reset word of length at most  $2n$ .
- The remainder of the Černý literature consists of a vast array of special, but interesting, cases.
- There is no time to survey the whole literature here.

# Pin's bounds

- It is straightforward to obtain a cubic upper bound of  $\frac{n^3-n}{3}$  on reset words for synchronizing automata.
- The best known upper bound for the synchronizing case is  $\frac{n^3-n}{6}$ , which was proved by Pin modulo an extremal set theory result of Frankl.
- Improving a bound by a factor of 2 can be hard work!
- The lower bound of  $(n-r)^2$  for rank  $r$  is due to Pin.
- Pin also has an analogous cubic upper bound for rank  $r$ .
- Probabilistically speaking, all automata are synchronizing with reset word of length at most  $2n$ .
- The remainder of the Černý literature consists of a vast array of special, but interesting, cases.
- There is no time to survey the whole literature here.

# Pin's bounds

- It is straightforward to obtain a cubic upper bound of  $\frac{n^3-n}{3}$  on reset words for synchronizing automata.
- The best known upper bound for the synchronizing case is  $\frac{n^3-n}{6}$ , which was proved by Pin modulo an extremal set theory result of Frankl.
- Improving a bound by a factor of 2 can be hard work!
- The lower bound of  $(n-r)^2$  for rank  $r$  is due to Pin.
- Pin also has an analogous cubic upper bound for rank  $r$ .
- Probabilistically speaking, all automata are synchronizing with reset word of length at most  $2n$ .
- The remainder of the Černý literature consists of a vast array of special, but interesting, cases.
- There is no time to survey the whole literature here.

# Some known results

- The special cases treated so far tend to be of two sorts:
  - 1 Combinatorial restrictions are imposed on the automata;
  - 2 Algebraic restrictions are imposed on the transition monoid.
- A key example of the first sort is the result of Dubuc that the Černý conjecture holds for **circular automata**: automata where one of the input letters cyclically permutes the state set.
- Kari proved that if the underlying digraph of the automaton is Eulerian, then the Černý conjecture holds.
- An important algebraic result is that of Trahtman establishing the Černý conjecture for automata with aperiodic transition monoid with an upper bound of  $n(n-1)/2$ .
- Rystsov showed that if the transition monoid is commutative, then the Černý conjecture holds with an upper bound of  $n-1$  (which is sharp).

# Some known results

- The special cases treated so far tend to be of two sorts:
  - ① **Combinatorial** restrictions are imposed on the automata;
  - ② **Algebraic** restrictions are imposed on the transition monoid.
- A key example of the first sort is the result of Dubuc that the Černý conjecture holds for **circular automata**: automata where one of the input letters cyclically permutes the state set.
- Kari proved that if the underlying digraph of the automaton is **Eulerian**, then the Černý conjecture holds.
- An important algebraic result is that of Trahtman establishing the Černý conjecture for automata with aperiodic transition monoid with an upper bound of  $n(n-1)/2$ .
- Rystsov showed that if the transition monoid is commutative, then the Černý conjecture holds with an upper bound of  $n-1$  (which is sharp).

# Some known results

- The special cases treated so far tend to be of two sorts:
  - ❶ **Combinatorial** restrictions are imposed on the automata;
  - ❷ **Algebraic** restrictions are imposed on the transition monoid.
- A key example of the first sort is the result of Dubuc that the Černý conjecture holds for **circular automata**: automata where one of the input letters cyclically permutes the state set.
- Kari proved that if the underlying digraph of the automaton is **Eulerian**, then the Černý conjecture holds.
- An important algebraic result is that of Trahtman establishing the Černý conjecture for automata with **aperiodic** transition monoid with an upper bound of  $n(n-1)/2$ .
- Rystsov showed that if the transition monoid is commutative, then the Černý conjecture holds with an upper bound of  $n-1$  (which is sharp).

# Some known results

- The special cases treated so far tend to be of two sorts:
  - 1 Combinatorial restrictions are imposed on the automata;
  - 2 Algebraic restrictions are imposed on the transition monoid.
- A key example of the first sort is the result of Dubuc that the Černý conjecture holds for **circular automata**: automata where one of the input letters cyclically permutes the state set.
- Kari proved that if the underlying digraph of the automaton is **Eulerian**, then the Černý conjecture holds.
- An important algebraic result is that of Trahtman establishing the Černý conjecture for automata with **aperiodic** transition monoid with an upper bound of  $n(n-1)/2$ .
- Rystsov showed that if the transition monoid is **commutative**, then the Černý conjecture holds with an upper bound of  $n-1$  (which is sharp).

# Some known results

- The special cases treated so far tend to be of two sorts:
  - ① **Combinatorial** restrictions are imposed on the automata;
  - ② **Algebraic** restrictions are imposed on the transition monoid.
- A key example of the first sort is the result of Dubuc that the Černý conjecture holds for **circular automata**: automata where one of the input letters cyclically permutes the state set.
- Kari proved that if the underlying digraph of the automaton is **Eulerian**, then the Černý conjecture holds.
- An important algebraic result is that of Trahtman establishing the Černý conjecture for automata with **aperiodic** transition monoid with an upper bound of  $n(n-1)/2$ .
- Rystsov showed that if the transition monoid is **commutative**, then the Černý conjecture holds with an upper bound of  $n-1$  (which is sharp).



# Some known results

- The special cases treated so far tend to be of two sorts:
  - 1 Combinatorial restrictions are imposed on the automata;
  - 2 Algebraic restrictions are imposed on the transition monoid.
- A key example of the first sort is the result of Dubuc that the Černý conjecture holds for **circular automata**: automata where one of the input letters cyclically permutes the state set.
- Kari proved that if the underlying digraph of the automaton is **Eulerian**, then the Černý conjecture holds.
- An important algebraic result is that of Trahtman establishing the Černý conjecture for automata with **aperiodic** transition monoid with an upper bound of  $n(n-1)/2$ .
- Rystsov showed that if the transition monoid is **commutative**, then the Černý conjecture holds with an upper bound of  $n-1$  (which is sharp).

# Some known results

- The special cases treated so far tend to be of two sorts:
  - ① **Combinatorial** restrictions are imposed on the automata;
  - ② **Algebraic** restrictions are imposed on the transition monoid.
- A key example of the first sort is the result of Dubuc that the Černý conjecture holds for **circular automata**: automata where one of the input letters cyclically permutes the state set.
- Kari proved that if the underlying digraph of the automaton is **Eulerian**, then the Černý conjecture holds.
- An important algebraic result is that of Trahtman establishing the Černý conjecture for automata with **aperiodic** transition monoid with an upper bound of  $n(n-1)/2$ .
- Rystsov showed that if the transition monoid is **commutative**, then the Černý conjecture holds with an upper bound of  $n-1$  (which is sharp).

# Representation theoretic approaches

- **Representation theory** is the study of algebraic objects using linear algebra.
- Many papers on Černý's conjecture make use in some form or the other of representation theory without using the full strength of the subject.
- For instance, Dubuc's paper on circular automata implicitly relies on properties of representations of cyclic groups.
- An approach using rational power series, pioneered by Béal, also relies on representation theory as representation theory lies in the foundations of weighted automata theory.
- Rystsov has a number of papers that make use of matrix representations to attack cases of the Černý conjecture.
- My goal is to explore representation theoretic approaches to the Černý conjecture.

# Representation theoretic approaches

- **Representation theory** is the study of algebraic objects using linear algebra.
- Many papers on Černý's conjecture make use in some form or the other of representation theory without using the full strength of the subject.
- For instance, Dubuc's paper on circular automata implicitly relies on properties of representations of cyclic groups.
- An approach using rational power series, pioneered by Béal, also relies on representation theory as representation theory lies in the foundations of weighted automata theory.
- Rystsov has a number of papers that make use of matrix representations to attack cases of the Černý conjecture.
- My goal is to explore representation theoretic approaches to the Černý conjecture.

# Representation theoretic approaches

- **Representation theory** is the study of algebraic objects using linear algebra.
- Many papers on Černý's conjecture make use in some form or the other of representation theory without using the full strength of the subject.
- For instance, Dubuc's paper on circular automata implicitly relies on properties of representations of cyclic groups.
- An approach using rational power series, pioneered by Béal, also relies on representation theory as representation theory lies in the foundations of weighted automata theory.
- Rystsov has a number of papers that make use of matrix representations to attack cases of the Černý conjecture.
- My goal is to explore representation theoretic approaches to the Černý conjecture.

# Representation theoretic approaches

- Representation theory is the study of algebraic objects using linear algebra.
- Many papers on Černý's conjecture make use in some form or the other of representation theory without using the full strength of the subject.
- For instance, Dubuc's paper on circular automata implicitly relies on properties of representations of cyclic groups.
- An approach using rational power series, pioneered by Béal, also relies on representation theory as representation theory lies in the foundations of weighted automata theory.
- Rystsov has a number of papers that make use of matrix representations to attack cases of the Černý conjecture.
- My goal is to explore representation theoretic approaches to the Černý conjecture.

# Representation theoretic approaches

- Representation theory is the study of algebraic objects using linear algebra.
- Many papers on Černý's conjecture make use in some form or the other of representation theory without using the full strength of the subject.
- For instance, Dubuc's paper on circular automata implicitly relies on properties of representations of cyclic groups.
- An approach using rational power series, pioneered by Béal, also relies on representation theory as representation theory lies in the foundations of weighted automata theory.
- Rystsov has a number of papers that make use of matrix representations to attack cases of the Černý conjecture.
- My goal is to explore representation theoretic approaches to the Černý conjecture.

# Representation theoretic approaches

- **Representation theory** is the study of algebraic objects using linear algebra.
- Many papers on Černý's conjecture make use in some form or the other of representation theory without using the full strength of the subject.
- For instance, Dubuc's paper on circular automata implicitly relies on properties of representations of cyclic groups.
- An approach using rational power series, pioneered by Béal, also relies on representation theory as representation theory lies in the foundations of weighted automata theory.
- Rystsov has a number of papers that make use of matrix representations to attack cases of the Černý conjecture.
- My goal is to explore representation theoretic approaches to the Černý conjecture.



## Theorem (Rystsov)

*Suppose that it is true that, given a set  $\Sigma \subseteq M_n(K)$  of  $n \times n$  matrices over a field  $K$  such that*

- 1  $|\langle \Sigma \rangle| < \infty$
- 2  $0 \in \langle \Sigma \rangle$ ,

*there is a word  $w \in \Sigma^*$  of length at most  $n^2$  representing the zero matrix. Then the Černý-Pin conjecture is true.*

- Unfortunately, Rystsov's conjecture is false.
- Paterson showed that it is undecidable whether the monoid generated by a finite subset of  $M_3(\mathbb{Z})$  contains 0 (The Matrix Mortality Problem).
- If Rystsov's conjecture were true, then by considering reduction modulo primes this problem would be decidable.

## Theorem (Rystsov)

*Suppose that it is true that, given a set  $\Sigma \subseteq M_n(K)$  of  $n \times n$  matrices over a field  $K$  such that*

- ❶  $|\langle \Sigma \rangle| < \infty$
- ❷  $0 \in \langle \Sigma \rangle$ ,

*there is a word  $w \in \Sigma^*$  of length at most  $n^2$  representing the zero matrix. Then the Černý-Pin conjecture is true.*

- Unfortunately, Rystsov's conjecture is false.
- Paterson showed that it is undecidable whether the monoid generated by a finite subset of  $M_3(\mathbb{Z})$  contains 0 (The Matrix Mortality Problem).
- If Rystsov's conjecture were true, then by considering reduction modulo primes this problem would be decidable.

## Theorem (Rystsov)

*Suppose that it is true that, given a set  $\Sigma \subseteq M_n(K)$  of  $n \times n$  matrices over a field  $K$  such that*

- ❶  $|\langle \Sigma \rangle| < \infty$
- ❷  $0 \in \langle \Sigma \rangle$ ,

*there is a word  $w \in \Sigma^*$  of length at most  $n^2$  representing the zero matrix. Then the Černý-Pin conjecture is true.*

- Unfortunately, Rystsov's conjecture is false.
- Paterson showed that it is undecidable whether the monoid generated by a finite subset of  $M_3(\mathbb{Z})$  contains 0 (The Matrix Mortality Problem).
- If Rystsov's conjecture were true, then by considering reduction modulo primes this problem would be decidable.

## Theorem (Rystsov)

*Suppose that it is true that, given a set  $\Sigma \subseteq M_n(K)$  of  $n \times n$  matrices over a field  $K$  such that*

- ❶  $|\langle \Sigma \rangle| < \infty$
- ❷  $0 \in \langle \Sigma \rangle$ ,

*there is a word  $w \in \Sigma^*$  of length at most  $n^2$  representing the zero matrix. Then the Černý-Pin conjecture is true.*

- Unfortunately, Rystsov's conjecture is false.
- Paterson showed that it is undecidable whether the monoid generated by a finite subset of  $M_3(\mathbb{Z})$  contains 0 (**The Matrix Mortality Problem**).
- If Rystsov's conjecture were true, then by considering reduction modulo primes this problem would be decidable.

## Theorem (Rystsov)

*Suppose that it is true that, given a set  $\Sigma \subseteq M_n(K)$  of  $n \times n$  matrices over a field  $K$  such that*

- ❶  $|\langle \Sigma \rangle| < \infty$
- ❷  $0 \in \langle \Sigma \rangle$ ,

*there is a word  $w \in \Sigma^*$  of length at most  $n^2$  representing the zero matrix. Then the Černý-Pin conjecture is true.*

- Unfortunately, Rystsov's conjecture is false.
- Paterson showed that it is undecidable whether the monoid generated by a finite subset of  $M_3(\mathbb{Z})$  contains 0 (**The Matrix Mortality Problem**).
- If Rystsov's conjecture were true, then by considering reduction modulo primes this problem would be decidable.

## Theorem (Rystsov)

*Suppose that it is true that, given a set  $\Sigma \subseteq M_n(K)$  of  $n \times n$  matrices over a field  $K$  such that*

- ❶  $|\langle \Sigma \rangle| < \infty$
- ❷  $0 \in \langle \Sigma \rangle$ ,

*there is a word  $w \in \Sigma^*$  of length at most  $n^2$  representing the zero matrix. Then the Černý-Pin conjecture is true.*

- Unfortunately, Rystsov's conjecture is false.
- Paterson showed that it is undecidable whether the monoid generated by a finite subset of  $M_3(\mathbb{Z})$  contains 0 (**The Matrix Mortality Problem**).
- If Rystsov's conjecture were true, then by considering reduction modulo primes this problem would be decidable.

## Theorem (Rystsov)

*Suppose that it is true that, given a set  $\Sigma \subseteq M_n(K)$  of  $n \times n$  matrices over a field  $K$  such that*

- ❶  $|\langle \Sigma \rangle| < \infty$
- ❷  $0 \in \langle \Sigma \rangle$ ,

*there is a word  $w \in \Sigma^*$  of length at most  $n^2$  representing the zero matrix. Then the Černý-Pin conjecture is true.*

- Unfortunately, Rystsov's conjecture is false.
- Paterson showed that it is undecidable whether the monoid generated by a finite subset of  $M_3(\mathbb{Z})$  contains 0 (**The Matrix Mortality Problem**).
- If Rystsov's conjecture were true, then by considering reduction modulo primes this problem would be decidable.

# Mortality functions

- The proof of Rystsov's result uses only the field  $\mathbb{F}_2$ .
- We find it more convenient to work with the field  $\mathbb{Q}$  of rational numbers.
- A monoid homomorphism  $\rho: M \rightarrow M_n(\mathbb{Q})$  is called a **representation of degree  $n$** .
- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is a **mortality function** for the monoid  $M$  if, for all representations  $\rho: M \rightarrow M_n(\mathbb{Q})$  with  $0 \in \rho(M)$  and all generating sets  $\Sigma$  for  $M$ , there exists  $w \in \Sigma^*$  of length at most  $f(n)$  such that  $\rho(w) = 0$ .
- Of course  $f(n) = |M| - 1$  is a mortality function for a finite monoid  $M$ .
- We say  $f$  is a mortality function for a class of monoids  $\mathcal{C}$  if it is a mortality function for all monoids in  $\mathcal{C}$ .
- By a universal mortality function, we mean a mortality function for the class of all finite monoids.



# Mortality functions

- The proof of Rystsov's result uses only the field  $\mathbb{F}_2$ .
- We find it more convenient to work with the field  $\mathbb{Q}$  of rational numbers.
- A monoid homomorphism  $\rho: M \rightarrow M_n(\mathbb{Q})$  is called a **representation of degree  $n$** .
- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is a **mortality function** for the monoid  $M$  if, for all representations  $\rho: M \rightarrow M_n(\mathbb{Q})$  with  $0 \in \rho(M)$  and all generating sets  $\Sigma$  for  $M$ , there exists  $w \in \Sigma^*$  of length at most  $f(n)$  such that  $\rho(w) = 0$ .
- Of course  $f(n) = |M| - 1$  is a mortality function for a finite monoid  $M$ .
- We say  $f$  is a mortality function for a class of monoids  $\mathcal{C}$  if it is a mortality function for all monoids in  $\mathcal{C}$ .
- By a universal mortality function, we mean a mortality function for the class of all finite monoids.

# Mortality functions

- The proof of Rystsov's result uses only the field  $\mathbb{F}_2$ .
- We find it more convenient to work with the field  $\mathbb{Q}$  of rational numbers.
- A monoid homomorphism  $\rho: M \rightarrow M_n(\mathbb{Q})$  is called a **representation** of **degree**  $n$ .
- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is a **mortality function** for the monoid  $M$  if, for all representations  $\rho: M \rightarrow M_n(\mathbb{Q})$  with  $0 \in \rho(M)$  and all generating sets  $\Sigma$  for  $M$ , there exists  $w \in \Sigma^*$  of length at most  $f(n)$  such that  $\rho(w) = 0$ .
- Of course  $f(n) = |M| - 1$  is a mortality function for a finite monoid  $M$ .
- We say  $f$  is a **mortality function** for a class of monoids  $\mathcal{C}$  if it is a mortality function for all monoids in  $\mathcal{C}$ .
- By a **universal mortality function**, we mean a mortality function for the class of all finite monoids.

# Mortality functions

- The proof of Rystsov's result uses only the field  $\mathbb{F}_2$ .
- We find it more convenient to work with the field  $\mathbb{Q}$  of rational numbers.
- A monoid homomorphism  $\rho: M \rightarrow M_n(\mathbb{Q})$  is called a **representation** of **degree**  $n$ .
- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is a **mortality function** for the monoid  $M$  if, for all representations  $\rho: M \rightarrow M_n(\mathbb{Q})$  with  $0 \in \rho(M)$  and all generating sets  $\Sigma$  for  $M$ , there exists  $w \in \Sigma^*$  of length at most  $f(n)$  such that  $\rho(w) = 0$ .
- Of course  $f(n) = |M| - 1$  is a mortality function for a finite monoid  $M$ .
- We say  $f$  is a **mortality function** for a class of monoids  $\mathcal{C}$  if it is a mortality function for all monoids in  $\mathcal{C}$ .
- By a **universal mortality function**, we mean a mortality function for the class of all finite monoids.

# Mortality functions

- The proof of Rystsov's result uses only the field  $\mathbb{F}_2$ .
- We find it more convenient to work with the field  $\mathbb{Q}$  of rational numbers.
- A monoid homomorphism  $\rho: M \rightarrow M_n(\mathbb{Q})$  is called a **representation** of **degree**  $n$ .
- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is a **mortality function** for the monoid  $M$  if, for all representations  $\rho: M \rightarrow M_n(\mathbb{Q})$  with  $0 \in \rho(M)$  and all generating sets  $\Sigma$  for  $M$ , there exists  $w \in \Sigma^*$  of length at most  $f(n)$  such that  $\rho(w) = 0$ .
- Of course  $f(n) = |M| - 1$  is a mortality function for a finite monoid  $M$ .
- We say  $f$  is a **mortality function** for a class of monoids  $\mathcal{C}$  if it is a mortality function for all monoids in  $\mathcal{C}$ .
- By a **universal mortality function**, we mean a mortality function for the class of all finite monoids.

# Mortality functions

- The proof of Rystsov's result uses only the field  $\mathbb{F}_2$ .
- We find it more convenient to work with the field  $\mathbb{Q}$  of rational numbers.
- A monoid homomorphism  $\rho: M \rightarrow M_n(\mathbb{Q})$  is called a **representation** of **degree**  $n$ .
- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is a **mortality function** for the monoid  $M$  if, for all representations  $\rho: M \rightarrow M_n(\mathbb{Q})$  with  $0 \in \rho(M)$  and all generating sets  $\Sigma$  for  $M$ , there exists  $w \in \Sigma^*$  of length at most  $f(n)$  such that  $\rho(w) = 0$ .
- Of course  $f(n) = |M| - 1$  is a mortality function for a finite monoid  $M$ .
- We say  $f$  is a **mortality function** for a class of monoids  $\mathcal{C}$  if it is a mortality function for all monoids in  $\mathcal{C}$ .
- By a **universal mortality function**, we mean a mortality function for the class of all finite monoids.

# Mortality functions

- The proof of Rystsov's result uses only the field  $\mathbb{F}_2$ .
- We find it more convenient to work with the field  $\mathbb{Q}$  of rational numbers.
- A monoid homomorphism  $\rho: M \rightarrow M_n(\mathbb{Q})$  is called a **representation** of **degree**  $n$ .
- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is a **mortality function** for the monoid  $M$  if, for all representations  $\rho: M \rightarrow M_n(\mathbb{Q})$  with  $0 \in \rho(M)$  and all generating sets  $\Sigma$  for  $M$ , there exists  $w \in \Sigma^*$  of length at most  $f(n)$  such that  $\rho(w) = 0$ .
- Of course  $f(n) = |M| - 1$  is a mortality function for a finite monoid  $M$ .
- We say  $f$  is a **mortality function** for a class of monoids  $\mathcal{C}$  if it is a mortality function for all monoids in  $\mathcal{C}$ .
- By a **universal mortality function**, we mean a mortality function for the class of all finite monoids.

# Mortality functions and the Černý-Pin problem

- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is **superadditive** if  $f(m) + f(n) \leq f(m + n)$ .
- The following result was inspired by Rystsov's argument.

## Theorem (Almeida, BS)

*Let  $\mathcal{A}$  be an  $n$ -state automaton of rank  $r$  with transition monoid  $M$  and suppose that  $f$  is a superadditive mortality function for  $M$ . Then there is a word of length at most  $f(n - r)$  having rank  $r$ .*

- So if  $n^2$  is a universal mortality function (which I don't believe), then the Černý-Pin conjecture is true.
- Almeida and I have obtained quadratic mortality bounds for a large class of monoids.
- For this talk, we restrict our attention to a special case of our results.

# Mortality functions and the Černý-Pin problem

- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is **superadditive** if  $f(m) + f(n) \leq f(m + n)$ .
- The following result was inspired by Rystsov's argument.

## Theorem (Almeida, BS)

*Let  $\mathcal{A}$  be an  $n$ -state automaton of rank  $r$  with transition monoid  $M$  and suppose that  $f$  is a superadditive mortality function for  $M$ . Then there is a word of length at most  $f(n - r)$  having rank  $r$ .*

- So if  $n^2$  is a universal mortality function (which I don't believe), then the Černý-Pin conjecture is true.
- Almeida and I have obtained quadratic mortality bounds for a large class of monoids.
- For this talk, we restrict our attention to a special case of our results.



# Mortality functions and the Černý-Pin problem

- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is **superadditive** if  $f(m) + f(n) \leq f(m + n)$ .
- The following result was inspired by Rystsov's argument.

## Theorem (Almeida, BS)

*Let  $\mathcal{A}$  be an  $n$ -state automaton of rank  $r$  with transition monoid  $M$  and suppose that  $f$  is a superadditive mortality function for  $M$ . Then there is a word of length at most  $f(n - r)$  having rank  $r$ .*

- So if  $n^2$  is a universal mortality function (which I don't believe), then the Černý-Pin conjecture is true.
- Almeida and I have obtained quadratic mortality bounds for a large class of monoids.
- For this talk, we restrict our attention to a special case of our results.

# Mortality functions and the Černý-Pin problem

- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is **superadditive** if  $f(m) + f(n) \leq f(m + n)$ .
- The following result was inspired by Rystsov's argument.

## Theorem (Almeida, BS)

*Let  $\mathcal{A}$  be an  $n$ -state automaton of rank  $r$  with transition monoid  $M$  and suppose that  $f$  is a superadditive mortality function for  $M$ . Then there is a word of length at most  $f(n - r)$  having rank  $r$ .*

- So if  $n^2$  is a universal mortality function (which I don't believe), then the Černý-Pin conjecture is true.
- Almeida and I have obtained quadratic mortality bounds for a large class of monoids.
- For this talk, we restrict our attention to a special case of our results.

# Mortality functions and the Černý-Pin problem

- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is **superadditive** if  $f(m) + f(n) \leq f(m + n)$ .
- The following result was inspired by Rystsov's argument.

## Theorem (Almeida, BS)

*Let  $\mathcal{A}$  be an  $n$ -state automaton of rank  $r$  with transition monoid  $M$  and suppose that  $f$  is a superadditive mortality function for  $M$ . Then there is a word of length at most  $f(n - r)$  having rank  $r$ .*

- So if  $n^2$  is a universal mortality function (which I don't believe), then the Černý-Pin conjecture is true.
- Almeida and I have obtained quadratic mortality bounds for a large class of monoids.
- For this talk, we restrict our attention to a special case of our results.

# Mortality functions and the Černý-Pin problem

- A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is **superadditive** if  $f(m) + f(n) \leq f(m + n)$ .
- The following result was inspired by Rystsov's argument.

## Theorem (Almeida, BS)

*Let  $\mathcal{A}$  be an  $n$ -state automaton of rank  $r$  with transition monoid  $M$  and suppose that  $f$  is a superadditive mortality function for  $M$ . Then there is a word of length at most  $f(n - r)$  having rank  $r$ .*

- So if  $n^2$  is a universal mortality function (which I don't believe), then the Černý-Pin conjecture is true.
- Almeida and I have obtained quadratic mortality bounds for a large class of monoids.
- For this talk, we restrict our attention to a special case of our results.

# A proof for the synchronizing case

- We outline a proof of the theorem for the synchronizing case, as it is much easier.
- Let  $\mathcal{A} = (Q, \Sigma)$  be an  $n$ -state automaton with transition monoid  $M$  and assume  $Q = \{1, \dots, n\}$ .
- Let  $e_1, \dots, e_n$  be the standard basis of row vectors for  $\mathbb{Q}^n$ .
- To each  $a \in \Sigma$ , associate the linear transformation  $\rho(a)$  given by  $e_i \rho(a) = e_{i \cdot a}$ .
- This induces an action of  $M$  on  $\mathbb{Q}^n$  by linear maps.
- Let  $V_0 = \{(c_1, \dots, c_n) \in \mathbb{Q}^n \mid c_1 + \dots + c_n = 0\} = \text{Span}\{e_i - e_j \mid 1 \leq i < j \leq n\}$ .
- $V_0$  is a hyperplane with basis  $\{e_1 - e_2, e_1 - e_3, \dots, e_1 - e_n\}$ , so it has dimension  $n - 1$ .
- Moreover,  $V_0$  is invariant under  $M$ .

# A proof for the synchronizing case

- We outline a proof of the theorem for the synchronizing case, as it is much easier.
- Let  $\mathcal{A} = (Q, \Sigma)$  be an  $n$ -state automaton with transition monoid  $M$  and assume  $Q = \{1, \dots, n\}$ .
- Let  $e_1, \dots, e_n$  be the standard basis of row vectors for  $\mathbb{Q}^n$ .
- To each  $a \in \Sigma$ , associate the linear transformation  $\rho(a)$  given by  $e_i \rho(a) = e_{i \cdot a}$ .
- This induces an action of  $M$  on  $\mathbb{Q}^n$  by linear maps.
- Let  $V_0 = \{(c_1, \dots, c_n) \in \mathbb{Q}^n \mid c_1 + \dots + c_n = 0\} = \text{Span}\{e_i - e_j \mid 1 \leq i < j \leq n\}$ .
- $V_0$  is a hyperplane with basis  $\{e_1 - e_2, e_1 - e_3, \dots, e_1 - e_n\}$ , so it has dimension  $n - 1$ .
- Moreover,  $V_0$  is invariant under  $M$ .

# A proof for the synchronizing case

- We outline a proof of the theorem for the synchronizing case, as it is much easier.
- Let  $\mathcal{A} = (Q, \Sigma)$  be an  $n$ -state automaton with transition monoid  $M$  and assume  $Q = \{1, \dots, n\}$ .
- Let  $e_1, \dots, e_n$  be the standard basis of row vectors for  $\mathbb{Q}^n$ .
- To each  $a \in \Sigma$ , associate the linear transformation  $\rho(a)$  given by  $e_i \rho(a) = e_{i \cdot a}$ .
- This induces an action of  $M$  on  $\mathbb{Q}^n$  by linear maps.
- Let  $V_0 = \{(c_1, \dots, c_n) \in \mathbb{Q}^n \mid c_1 + \dots + c_n = 0\} = \text{Span}\{e_i - e_j \mid 1 \leq i < j \leq n\}$ .
- $V_0$  is a hyperplane with basis  $\{e_1 - e_2, e_1 - e_3, \dots, e_1 - e_n\}$ , so it has dimension  $n - 1$ .
- Moreover,  $V_0$  is invariant under  $M$ .

# A proof for the synchronizing case

- We outline a proof of the theorem for the synchronizing case, as it is much easier.
- Let  $\mathcal{A} = (Q, \Sigma)$  be an  $n$ -state automaton with transition monoid  $M$  and assume  $Q = \{1, \dots, n\}$ .
- Let  $e_1, \dots, e_n$  be the standard basis of row vectors for  $\mathbb{Q}^n$ .
- To each  $a \in \Sigma$ , associate the linear transformation  $\rho(a)$  given by  $e_i \rho(a) = e_{i \cdot a}$ .
- This induces an action of  $M$  on  $\mathbb{Q}^n$  by linear maps.
- Let  $V_0 = \{(c_1, \dots, c_n) \in \mathbb{Q}^n \mid c_1 + \dots + c_n = 0\} = \text{Span}\{e_i - e_j \mid 1 \leq i < j \leq n\}$ .
- $V_0$  is a hyperplane with basis  $\{e_1 - e_2, e_1 - e_3, \dots, e_1 - e_n\}$ , so it has dimension  $n - 1$ .
- Moreover,  $V_0$  is invariant under  $M$ .



# A proof for the synchronizing case

- We outline a proof of the theorem for the synchronizing case, as it is much easier.
- Let  $\mathcal{A} = (Q, \Sigma)$  be an  $n$ -state automaton with transition monoid  $M$  and assume  $Q = \{1, \dots, n\}$ .
- Let  $e_1, \dots, e_n$  be the standard basis of row vectors for  $\mathbb{Q}^n$ .
- To each  $a \in \Sigma$ , associate the linear transformation  $\rho(a)$  given by  $e_i \rho(a) = e_{i \cdot a}$ .
- This induces an action of  $M$  on  $\mathbb{Q}^n$  by linear maps.
- Let  $V_0 = \{(c_1, \dots, c_n) \in \mathbb{Q}^n \mid c_1 + \dots + c_n = 0\} = \text{Span}\{e_i - e_j \mid 1 \leq i < j \leq n\}$ .
- $V_0$  is a hyperplane with basis  $\{e_1 - e_2, e_1 - e_3, \dots, e_1 - e_n\}$ , so it has dimension  $n - 1$ .
- Moreover,  $V_0$  is invariant under  $M$ .

# A proof for the synchronizing case

- We outline a proof of the theorem for the synchronizing case, as it is much easier.
- Let  $\mathcal{A} = (Q, \Sigma)$  be an  $n$ -state automaton with transition monoid  $M$  and assume  $Q = \{1, \dots, n\}$ .
- Let  $e_1, \dots, e_n$  be the standard basis of row vectors for  $\mathbb{Q}^n$ .
- To each  $a \in \Sigma$ , associate the linear transformation  $\rho(a)$  given by  $e_i \rho(a) = e_{i \cdot a}$ .
- This induces an action of  $M$  on  $\mathbb{Q}^n$  by linear maps.
- Let  $V_0 = \{(c_1, \dots, c_n) \in \mathbb{Q}^n \mid c_1 + \dots + c_n = 0\} = \text{Span}\{e_i - e_j \mid 1 \leq i < j \leq n\}$ .
- $V_0$  is a hyperplane with basis  $\{e_1 - e_2, e_1 - e_3, \dots, e_1 - e_n\}$ , so it has dimension  $n - 1$ .
- Moreover,  $V_0$  is invariant under  $M$ .

# A proof for the synchronizing case

- We outline a proof of the theorem for the synchronizing case, as it is much easier.
- Let  $\mathcal{A} = (Q, \Sigma)$  be an  $n$ -state automaton with transition monoid  $M$  and assume  $Q = \{1, \dots, n\}$ .
- Let  $e_1, \dots, e_n$  be the standard basis of row vectors for  $\mathbb{Q}^n$ .
- To each  $a \in \Sigma$ , associate the linear transformation  $\rho(a)$  given by  $e_i \rho(a) = e_{i \cdot a}$ .
- This induces an action of  $M$  on  $\mathbb{Q}^n$  by linear maps.
- Let  $V_0 = \{(c_1, \dots, c_n) \in \mathbb{Q}^n \mid c_1 + \dots + c_n = 0\} = \text{Span}\{e_i - e_j \mid 1 \leq i < j \leq n\}$ .
- $V_0$  is a hyperplane with basis  $\{e_1 - e_2, e_1 - e_3, \dots, e_1 - e_n\}$ , so it has dimension  $n - 1$ .
- Moreover,  $V_0$  is invariant under  $M$ .

# A proof for the synchronizing case

- We outline a proof of the theorem for the synchronizing case, as it is much easier.
- Let  $\mathcal{A} = (Q, \Sigma)$  be an  $n$ -state automaton with transition monoid  $M$  and assume  $Q = \{1, \dots, n\}$ .
- Let  $e_1, \dots, e_n$  be the standard basis of row vectors for  $\mathbb{Q}^n$ .
- To each  $a \in \Sigma$ , associate the linear transformation  $\rho(a)$  given by  $e_i \rho(a) = e_{i \cdot a}$ .
- This induces an action of  $M$  on  $\mathbb{Q}^n$  by linear maps.
- Let  $V_0 = \{(c_1, \dots, c_n) \in \mathbb{Q}^n \mid c_1 + \dots + c_n = 0\} = \text{Span}\{e_i - e_j \mid 1 \leq i < j \leq n\}$ .
- $V_0$  is a hyperplane with basis  $\{e_1 - e_2, e_1 - e_3, \dots, e_1 - e_n\}$ , so it has dimension  $n - 1$ .
- Moreover,  $V_0$  is invariant under  $M$ .

# A proof for the synchronizing case II

- We claim that  $w \in \Sigma^*$  is a synchronizing word iff  $\rho(w)|_{V_0} = 0$ .
- Indeed,  $(e_i - e_j)\rho(w) = e_{i \cdot w} - e_{j \cdot w}$ .
- So  $\rho(w)$  annihilates  $V_0$  iff  $i \cdot w = j \cdot w$  for all  $1 \leq i < j \leq n$ .
- But this occurs iff  $|Q \cdot w| = 1$ , i.e.,  $w$  is a reset word.
- It now follows that if  $f$  is a mortality function for  $M$ , then there is a reset word  $w$  for  $\mathcal{M}$  of length at most  $f(n-1)$ .
- Notice that for the synchronizing case the superadditivity of  $f$  is not required.
- Our argument for the Pin conjecture requires it.

# A proof for the synchronizing case II

- We claim that  $w \in \Sigma^*$  is a synchronizing word iff  $\rho(w)|_{V_0} = 0$ .
- Indeed,  $(e_i - e_j)\rho(w) = e_{i \cdot w} - e_{j \cdot w}$ .
- So  $\rho(w)$  annihilates  $V_0$  iff  $i \cdot w = j \cdot w$  for all  $1 \leq i < j \leq n$ .
- But this occurs iff  $|Q \cdot w| = 1$ , i.e.,  $w$  is a reset word.
- It now follows that if  $f$  is a mortality function for  $M$ , then there is a reset word  $w$  for  $\mathcal{A}$  of length at most  $f(n-1)$ .
- Notice that for the synchronizing case the superadditivity of  $f$  is not required.
- Our argument for the Pin conjecture requires it.

# A proof for the synchronizing case II

- We claim that  $w \in \Sigma^*$  is a synchronizing word iff  $\rho(w)|_{V_0} = 0$ .
- Indeed,  $(e_i - e_j)\rho(w) = e_{i \cdot w} - e_{j \cdot w}$ .
- So  $\rho(w)$  annihilates  $V_0$  iff  $i \cdot w = j \cdot w$  for all  $1 \leq i < j \leq n$ .
- But this occurs iff  $|Q \cdot w| = 1$ , i.e.,  $w$  is a reset word.
- It now follows that if  $f$  is a mortality function for  $M$ , then there is a reset word  $w$  for  $\mathcal{A}$  of length at most  $f(n-1)$ .
- Notice that for the synchronizing case the superadditivity of  $f$  is not required.
- Our argument for the Pin conjecture requires it.

# A proof for the synchronizing case II

- We claim that  $w \in \Sigma^*$  is a synchronizing word iff  $\rho(w)|_{V_0} = 0$ .
- Indeed,  $(e_i - e_j)\rho(w) = e_{i \cdot w} - e_{j \cdot w}$ .
- So  $\rho(w)$  annihilates  $V_0$  iff  $i \cdot w = j \cdot w$  for all  $1 \leq i < j \leq n$ .
- But this occurs iff  $|Q \cdot w| = 1$ , i.e.,  $w$  is a reset word.
- It now follows that if  $f$  is a mortality function for  $M$ , then there is a reset word  $w$  for  $\mathcal{A}$  of length at most  $f(n-1)$ .
- Notice that for the synchronizing case the superadditivity of  $f$  is not required.
- Our argument for the Pin conjecture requires it.



# A proof for the synchronizing case II

- We claim that  $w \in \Sigma^*$  is a synchronizing word iff  $\rho(w)|_{V_0} = 0$ .
- Indeed,  $(e_i - e_j)\rho(w) = e_{i \cdot w} - e_{j \cdot w}$ .
- So  $\rho(w)$  annihilates  $V_0$  iff  $i \cdot w = j \cdot w$  for all  $1 \leq i < j \leq n$ .
- But this occurs iff  $|Q \cdot w| = 1$ , i.e.,  $w$  is a reset word.
- It now follows that if  $f$  is a mortality function for  $M$ , then there is a reset word  $w$  for  $\mathcal{A}$  of length at most  $f(n - 1)$ .
- Notice that for the synchronizing case the superadditivity of  $f$  is not required.
- Our argument for the Pin conjecture requires it.

# A proof for the synchronizing case II

- We claim that  $w \in \Sigma^*$  is a synchronizing word iff  $\rho(w)|_{V_0} = 0$ .
- Indeed,  $(e_i - e_j)\rho(w) = e_{i \cdot w} - e_{j \cdot w}$ .
- So  $\rho(w)$  annihilates  $V_0$  iff  $i \cdot w = j \cdot w$  for all  $1 \leq i < j \leq n$ .
- But this occurs iff  $|Q \cdot w| = 1$ , i.e.,  $w$  is a reset word.
- It now follows that if  $f$  is a mortality function for  $M$ , then there is a reset word  $w$  for  $\mathcal{A}$  of length at most  $f(n - 1)$ .
- Notice that for the synchronizing case the superadditivity of  $f$  is not required.
- Our argument for the Pin conjecture requires it.

# A proof for the synchronizing case II

- We claim that  $w \in \Sigma^*$  is a synchronizing word iff  $\rho(w)|_{V_0} = 0$ .
- Indeed,  $(e_i - e_j)\rho(w) = e_{i \cdot w} - e_{j \cdot w}$ .
- So  $\rho(w)$  annihilates  $V_0$  iff  $i \cdot w = j \cdot w$  for all  $1 \leq i < j \leq n$ .
- But this occurs iff  $|Q \cdot w| = 1$ , i.e.,  $w$  is a reset word.
- It now follows that if  $f$  is a mortality function for  $M$ , then there is a reset word  $w$  for  $\mathcal{A}$  of length at most  $f(n - 1)$ .
- Notice that for the synchronizing case the superadditivity of  $f$  is not required.
- Our argument for the Pin conjecture requires it.

# Representation theory

- Let  $\rho: M \rightarrow M_n(\mathbb{Q})$  be a representation and put  $V = \mathbb{Q}^n$ .
- A subspace  $W \leq V$  is said to be  **$M$ -invariant** if  $W\rho(M) \subseteq W$ .
- For example,  $V_0$  from the above proof is an  $M$ -invariant subspace of  $V$ .
- A representation is **irreducible** if  $\{0\}$  and  $V$  are the only  $M$ -invariant subspaces.
- Every representation can be 'built up' from irreducible representations in much the same way that every finite group can be 'built up' from finite simple groups.
- This allowed us to prove the following result by induction on the dimension.

## Theorem (Almeida, BS)

*Let  $f$  be a superadditive function. Then  $f$  is a mortality function for  $M$  iff  $f$  is a mortality function for each irreducible representation of  $M$ .*

# Representation theory

- Let  $\rho: M \rightarrow M_n(\mathbb{Q})$  be a representation and put  $V = \mathbb{Q}^n$ .
- A subspace  $W \leq V$  is said to be  **$M$ -invariant** if  $W\rho(M) \subseteq W$ .
- For example,  $V_0$  from the above proof is an  $M$ -invariant subspace of  $V$ .
- A representation is **irreducible** if  $\{0\}$  and  $V$  are the only  $M$ -invariant subspaces.
- Every representation can be 'built up' from irreducible representations in much the same way that every finite group can be 'built up' from finite simple groups.
- This allowed us to prove the following result by induction on the dimension.

## Theorem (Almeida, BS)

*Let  $f$  be a superadditive function. Then  $f$  is a mortality function for  $M$  iff  $f$  is a mortality function for each irreducible representation of  $M$ .*

# Representation theory

- Let  $\rho: M \rightarrow M_n(\mathbb{Q})$  be a representation and put  $V = \mathbb{Q}^n$ .
- A subspace  $W \leq V$  is said to be  **$M$ -invariant** if  $W\rho(M) \subseteq W$ .
- For example,  $V_0$  from the above proof is an  $M$ -invariant subspace of  $V$ .
- A representation is **irreducible** if  $\{0\}$  and  $V$  are the only  $M$ -invariant subspaces.
- Every representation can be 'built up' from irreducible representations in much the same way that every finite group can be 'built up' from finite simple groups.
- This allowed us to prove the following result by induction on the dimension.

## Theorem (Almeida BS)

*Let  $f$  be a superadditive function. Then  $f$  is a mortality function for  $M$  iff  $f$  is a mortality function for each irreducible representation of  $M$ .*

# Representation theory

- Let  $\rho: M \rightarrow M_n(\mathbb{Q})$  be a representation and put  $V = \mathbb{Q}^n$ .
- A subspace  $W \leq V$  is said to be  **$M$ -invariant** if  $W\rho(M) \subseteq W$ .
- For example,  $V_0$  from the above proof is an  $M$ -invariant subspace of  $V$ .
- A representation is **irreducible** if  $\{0\}$  and  $V$  are the only  $M$ -invariant subspaces.
- Every representation can be 'built up' from irreducible representations in much the same way that every finite group can be 'built up' from finite simple groups.
- This allowed us to prove the following result by induction on the dimension.

## Theorem (Almeida,BS)

*Let  $f$  be a superadditive function. Then  $f$  is a mortality function for  $M$  iff  $f$  is a mortality function for each irreducible representation of  $M$ .*

# Representation theory

- Let  $\rho: M \rightarrow M_n(\mathbb{Q})$  be a representation and put  $V = \mathbb{Q}^n$ .
- A subspace  $W \leq V$  is said to be  **$M$ -invariant** if  $W\rho(M) \subseteq W$ .
- For example,  $V_0$  from the above proof is an  $M$ -invariant subspace of  $V$ .
- A representation is **irreducible** if  $\{0\}$  and  $V$  are the only  $M$ -invariant subspaces.
- Every representation can be 'built up' from irreducible representations in much the same way that every finite group can be 'built up' from finite simple groups.
- This allowed us to prove the following result by induction on the dimension.

## Theorem (Almeida,BS)

*Let  $f$  be a superadditive function. Then  $f$  is a mortality function for  $M$  iff  $f$  is a mortality function for each irreducible representation of  $M$ .*



# Representation theory

- Let  $\rho: M \rightarrow M_n(\mathbb{Q})$  be a representation and put  $V = \mathbb{Q}^n$ .
- A subspace  $W \leq V$  is said to be  **$M$ -invariant** if  $W\rho(M) \subseteq W$ .
- For example,  $V_0$  from the above proof is an  $M$ -invariant subspace of  $V$ .
- A representation is **irreducible** if  $\{0\}$  and  $V$  are the only  $M$ -invariant subspaces.
- Every representation can be 'built up' from irreducible representations in much the same way that every finite group can be 'built up' from finite simple groups.
- This allowed us to prove the following result by induction on the dimension.

## Theorem (Almeida,BS)

*Let  $f$  be a superadditive function. Then  $f$  is a mortality function for  $M$  iff  $f$  is a mortality function for each irreducible representation of  $M$ .*

# Representation theory

- Let  $\rho: M \rightarrow M_n(\mathbb{Q})$  be a representation and put  $V = \mathbb{Q}^n$ .
- A subspace  $W \leq V$  is said to be  **$M$ -invariant** if  $W\rho(M) \subseteq W$ .
- For example,  $V_0$  from the above proof is an  $M$ -invariant subspace of  $V$ .
- A representation is **irreducible** if  $\{0\}$  and  $V$  are the only  $M$ -invariant subspaces.
- Every representation can be 'built up' from irreducible representations in much the same way that every finite group can be 'built up' from finite simple groups.
- This allowed us to prove the following result by induction on the dimension.

## Theorem (Almeida,BS)

*Let  $f$  be a superadditive function. Then  $f$  is a mortality function for  $M$  iff  $f$  is a mortality function for each irreducible representation of  $M$ .*

# Irreducible representations

- There is a well-developed theory of irreducible representations of finite monoids due to Munn-Ponizovsky and further elaborated by Rhodes and Zalcstein.
- In particular, the irreducible representations of a finite monoid  $M$  can be constructed from the irreducible representations of its maximal subgroups.
- The degrees of the irreducible representations are intimately connected to the Rees matrix representations of the principal factors of  $M$ , that is, the semigroups of the form  $J^0$  with  $J$  a regular  $\mathcal{J}$ -class of  $M$ .

# Irreducible representations

- There is a well-developed theory of irreducible representations of finite monoids due to Munn-Ponizovsky and further elaborated by Rhodes and Zalcstein.
- In particular, the irreducible representations of a finite monoid  $M$  can be constructed from the irreducible representations of its **maximal subgroups**.
- The degrees of the irreducible representations are intimately connected to the Rees matrix representations of the principal factors of  $M$ , that is, the semigroups of the form  $J^0$  with  $J$  a regular  $\mathcal{J}$ -class of  $M$ .

- There is a well-developed theory of irreducible representations of finite monoids due to Munn-Ponizovsky and further elaborated by Rhodes and Zalcstein.
- In particular, the irreducible representations of a finite monoid  $M$  can be constructed from the irreducible representations of its **maximal subgroups**.
- The degrees of the irreducible representations are intimately connected to the Rees matrix representations of the principal factors of  $M$ , that is, the semigroups of the form  $J^0$  with  $J$  a regular  $\mathcal{J}$ -class of  $M$ .

# The case of DS

- A finite monoid belongs to the class **DS** if  $e \in MaM \cap MbM$  implies  $e \in MabM$  for all idempotents  $e \in M$ .
- Recall that  $e$  is **idempotent** if  $e^2 = e$ .
- Equivalently,  $M \in \text{DS}$  iff  $M \times M$  cannot recognize the language  $(ab)^*$ .
- This class was introduced independently by Putcha and Schützenberger.
- Examples of monoids in DS include:
  - commutative monoids (obvious);
  - monoids satisfying an identity  $x^m = x$  (Clifford);
  - monoids of upper triangular matrices over a finite field (Putcha).
- The variety DS appears frequently in the algebraic theory of automata.

# The case of DS

- A finite monoid belongs to the class **DS** if  $e \in MaM \cap MbM$  implies  $e \in MabM$  for all idempotents  $e \in M$ .
- Recall that  $e$  is **idempotent** if  $e^2 = e$ .
- Equivalently,  $M \in \text{DS}$  iff  $M \times M$  cannot recognize the language  $(ab)^*$ .
- This class was introduced independently by Putcha and Schützenberger.
- Examples of monoids in DS include:
  - commutative monoids (obvious);
  - monoids satisfying an identity  $x^m = x$  (Clifford);
  - monoids of upper triangular matrices over a finite field (Putcha).
- The variety DS appears frequently in the algebraic theory of automata.

# The case of DS

- A finite monoid belongs to the class **DS** if  $e \in MaM \cap MbM$  implies  $e \in MabM$  for all idempotents  $e \in M$ .
- Recall that  $e$  is **idempotent** if  $e^2 = e$ .
- Equivalently,  $M \in \text{DS}$  iff  $M \times M$  cannot recognize the language  $(ab)^*$ .
- This class was introduced independently by Putcha and Schützenberger.
- Examples of monoids in DS include:
  - commutative monoids (obvious);
  - monoids satisfying an identity  $x^m = x$  (Clifford);
  - monoids of upper triangular matrices over a finite field (Putcha).
- The variety DS appears frequently in the algebraic theory of automata.



# The case of DS

- A finite monoid belongs to the class **DS** if  $e \in MaM \cap MbM$  implies  $e \in MabM$  for all idempotents  $e \in M$ .
- Recall that  $e$  is **idempotent** if  $e^2 = e$ .
- Equivalently,  $M \in \text{DS}$  iff  $M \times M$  cannot recognize the language  $(ab)^*$ .
- This class was introduced independently by Putcha and Schützenberger.
- Examples of monoids in DS include:
  - commutative monoids (obvious);
  - monoids satisfying an identity  $x^m = x$  (Clifford);
  - monoids of upper triangular matrices over a finite field (Putcha).
- The variety DS appears frequently in the algebraic theory of automata.

# The case of DS

- A finite monoid belongs to the class **DS** if  $e \in MaM \cap MbM$  implies  $e \in MabM$  for all idempotents  $e \in M$ .
- Recall that  $e$  is **idempotent** if  $e^2 = e$ .
- Equivalently,  $M \in \text{DS}$  iff  $M \times M$  cannot recognize the language  $(ab)^*$ .
- This class was introduced independently by Putcha and Schützenberger.
- Examples of monoids in DS include:
  - commutative monoids (obvious);
  - monoids satisfying an identity  $x^m = x$  (Clifford);
  - monoids of upper triangular matrices over a finite field (Putcha).
- The variety DS appears frequently in the algebraic theory of automata.

# The case of DS

- A finite monoid belongs to the class **DS** if  $e \in MaM \cap MbM$  implies  $e \in MabM$  for all idempotents  $e \in M$ .
- Recall that  $e$  is **idempotent** if  $e^2 = e$ .
- Equivalently,  $M \in \text{DS}$  iff  $M \times M$  cannot recognize the language  $(ab)^*$ .
- This class was introduced independently by Putcha and Schützenberger.
- Examples of monoids in DS include:
  - commutative monoids (obvious);
  - monoids satisfying an identity  $x^m = x$  (Clifford);
  - monoids of upper triangular matrices over a finite field (Putcha).
- The variety DS appears frequently in the algebraic theory of automata.

# The case of DS

- A finite monoid belongs to the class **DS** if  $e \in MaM \cap MbM$  implies  $e \in MabM$  for all idempotents  $e \in M$ .
- Recall that  $e$  is **idempotent** if  $e^2 = e$ .
- Equivalently,  $M \in \text{DS}$  iff  $M \times M$  cannot recognize the language  $(ab)^*$ .
- This class was introduced independently by Putcha and Schützenberger.
- Examples of monoids in DS include:
  - commutative monoids (obvious);
  - monoids satisfying an identity  $x^m = x$  (Clifford);
  - monoids of upper triangular matrices over a finite field (Putcha).
- The variety DS appears frequently in the algebraic theory of automata.

# The case of DS

- A finite monoid belongs to the class **DS** if  $e \in MaM \cap MbM$  implies  $e \in MabM$  for all idempotents  $e \in M$ .
- Recall that  $e$  is **idempotent** if  $e^2 = e$ .
- Equivalently,  $M \in \text{DS}$  iff  $M \times M$  cannot recognize the language  $(ab)^*$ .
- This class was introduced independently by Putcha and Schützenberger.
- Examples of monoids in DS include:
  - commutative monoids (obvious);
  - monoids satisfying an identity  $x^m = x$  (Clifford);
  - monoids of upper triangular matrices over a finite field (Putcha).
- The variety DS appears frequently in the algebraic theory of automata.

# The case of DS

- A finite monoid belongs to the class **DS** if  $e \in MaM \cap MbM$  implies  $e \in MabM$  for all idempotents  $e \in M$ .
- Recall that  $e$  is **idempotent** if  $e^2 = e$ .
- Equivalently,  $M \in \text{DS}$  iff  $M \times M$  cannot recognize the language  $(ab)^*$ .
- This class was introduced independently by Putcha and Schützenberger.
- Examples of monoids in DS include:
  - commutative monoids (obvious);
  - monoids satisfying an identity  $x^m = x$  (Clifford);
  - monoids of upper triangular matrices over a finite field (Putcha).
- The variety DS appears frequently in the algebraic theory of automata.

# The case of DS II

- In previous work, we showed with Almeida, Margolis and Volkov that the Černý conjecture holds for automata with transition monoid in DS using representation theory.
- Volkov asked in LATA 2008 whether the bound of  $(n - 1)^2$  was tight for this class.
- The main ingredient in our proof was the following result.

## Lemma (AMSV)

*If  $\rho: M \rightarrow M_n(\mathbb{Q})$  is an irreducible representation of a monoid in DS such that  $0 \in \rho(M)$  and  $\Sigma$  is a generating set for  $M$ , then there is a **letter**  $a \in \Sigma$  with  $\rho(a) = 0$ .*

- Consequently,  $f(n) = n$  is a superadditive mortality function for DS.

# The case of DS II

- In previous work, we showed with Almeida, Margolis and Volkov that the Černý conjecture holds for automata with transition monoid in DS using representation theory.
- Volkov asked in LATA 2008 whether the bound of  $(n - 1)^2$  was tight for this class.
- The main ingredient in our proof was the following result.

## Lemma (AMSV)

*If  $\rho: M \rightarrow M_n(\mathbb{Q})$  is an irreducible representation of a monoid in DS such that  $0 \in \rho(M)$  and  $\Sigma$  is a generating set for  $M$ , then there is a **letter**  $a \in \Sigma$  with  $\rho(a) = 0$ .*

- Consequently,  $f(n) = n$  is a superadditive mortality function for DS.



# The case of DS II

- In previous work, we showed with Almeida, Margolis and Volkov that the Černý conjecture holds for automata with transition monoid in DS using representation theory.
- Volkov asked in LATA 2008 whether the bound of  $(n - 1)^2$  was tight for this class.
- The main ingredient in our proof was the following result.

## Lemma (AMSV)

*If  $\rho: M \rightarrow M_n(\mathbb{Q})$  is an irreducible representation of a monoid in DS such that  $0 \in \rho(M)$  and  $\Sigma$  is a generating set for  $M$ , then there is a **letter**  $a \in \Sigma$  with  $\rho(a) = 0$ .*

- Consequently,  $f(n) = n$  is a superadditive mortality function for DS.

- In previous work, we showed with Almeida, Margolis and Volkov that the Černý conjecture holds for automata with transition monoid in DS using representation theory.
- Volkov asked in LATA 2008 whether the bound of  $(n - 1)^2$  was tight for this class.
- The main ingredient in our proof was the following result.

## Lemma (AMSV)

*If  $\rho: M \rightarrow M_n(\mathbb{Q})$  is an irreducible representation of a monoid in DS such that  $0 \in \rho(M)$  and  $\Sigma$  is a generating set for  $M$ , then there is a **letter**  $a \in \Sigma$  with  $\rho(a) = 0$ .*

- Consequently,  $f(n) = n$  is a superadditive mortality function for DS.

- In previous work, we showed with Almeida, Margolis and Volkov that the Černý conjecture holds for automata with transition monoid in DS using representation theory.
- Volkov asked in LATA 2008 whether the bound of  $(n - 1)^2$  was tight for this class.
- The main ingredient in our proof was the following result.

## Lemma (AMSV)

*If  $\rho: M \rightarrow M_n(\mathbb{Q})$  is an irreducible representation of a monoid in DS such that  $0 \in \rho(M)$  and  $\Sigma$  is a generating set for  $M$ , then there is a **letter**  $a \in \Sigma$  with  $\rho(a) = 0$ .*

- Consequently,  $f(n) = n$  is a superadditive mortality function for DS.

# The Černý-Pin conjecture for DS

## Theorem (Almeida, BS)

*If  $\mathcal{A}$  is an  $n$ -state, rank  $r$  automaton with transition monoid in DS, then there is a word  $w$  of length at most  $n - r$  with  $\text{rk}(w) = r$ .*

- This bound is easily seen to be sharp by considering automata over unary alphabets.
- For instance,



is synchronizing with minimum length reset word  $a^{n-1}$  and the transition monoid is commutative.

# The Černý-Pin conjecture for DS

## Theorem (Almeida, BS)

*If  $\mathcal{A}$  is an  $n$ -state, rank  $r$  automaton with transition monoid in DS, then there is a word  $w$  of length at most  $n - r$  with  $\text{rk}(w) = r$ .*

- This bound is easily seen to be sharp by considering automata over unary alphabets.
- For instance,




is synchronizing with minimum length reset word  $a^{n-1}$  and the transition monoid is commutative.

# The Černý-Pin conjecture for DS

## Theorem (Almeida, BS)

*If  $\mathcal{A}$  is an  $n$ -state, rank  $r$  automaton with transition monoid in DS, then there is a word  $w$  of length at most  $n - r$  with  $\text{rk}(w) = r$ .*

- This bound is easily seen to be sharp by considering automata over unary alphabets.
- For instance,



is synchronizing with minimum length reset word  $a^{n-1}$  and the transition monoid is commutative.

- A finite monoid belongs to the class EDS if its idempotents generate a submonoid in DS.
- $DS \subseteq EDS$ .
- Monoids with commuting idempotents (such as inverse monoids) belong to EDS.
- More generally, monoids whose idempotents form a submonoid belong to EDS.
- It is known that a monoid  $M$  belongs to EDS iff it cannot recognize the language  $\{a, b\}^*ab\{a, b\}^*$ .
- Equivalently, EDS is the largest variety of monoids that cannot recognize all 2-testable languages.

- A finite monoid belongs to the class EDS if its idempotents generate a submonoid in DS.
- $DS \subseteq EDS$ .
- Monoids with commuting idempotents (such as inverse monoids) belong to EDS.
- More generally, monoids whose idempotents form a submonoid belong to EDS.
- It is known that a monoid  $M$  belongs to EDS iff it cannot recognize the language  $\{a, b\}^*ab\{a, b\}^*$ .
- Equivalently, EDS is the largest variety of monoids that cannot recognize all 2-testable languages.



# The class EDS

- A finite monoid belongs to the class EDS if its idempotents generate a submonoid in DS.
- $DS \subseteq EDS$ .
- Monoids with commuting idempotents (such as inverse monoids) belong to EDS.
- More generally, monoids whose idempotents form a submonoid belong to EDS.
- It is known that a monoid  $M$  belongs to EDS iff it cannot recognize the language  $\{a, b\}^*ab\{a, b\}^*$ .
- Equivalently, EDS is the largest variety of monoids that cannot recognize all 2-testable languages.

# The class EDS

- A finite monoid belongs to the class EDS if its idempotents generate a submonoid in DS.
- $DS \subseteq EDS$ .
- Monoids with commuting idempotents (such as inverse monoids) belong to EDS.
- More generally, monoids whose idempotents form a submonoid belong to EDS.
- It is known that a monoid  $M$  belongs to EDS iff it cannot recognize the language  $\{a, b\}^*ab\{a, b\}^*$ .
- Equivalently, EDS is the largest variety of monoids that cannot recognize all 2-testable languages.

# The class EDS

- A finite monoid belongs to the class EDS if its idempotents generate a submonoid in DS.
- $DS \subseteq EDS$ .
- Monoids with commuting idempotents (such as inverse monoids) belong to EDS.
- More generally, monoids whose idempotents form a submonoid belong to EDS.
- It is known that a monoid  $M$  belongs to EDS iff it cannot recognize the language  $\{a, b\}^* ab \{a, b\}^*$ .
- Equivalently, EDS is the largest variety of monoids that cannot recognize all 2-testable languages.

# The class EDS

- A finite monoid belongs to the class EDS if its idempotents generate a submonoid in DS.
- $DS \subseteq EDS$ .
- Monoids with commuting idempotents (such as inverse monoids) belong to EDS.
- More generally, monoids whose idempotents form a submonoid belong to EDS.
- It is known that a monoid  $M$  belongs to EDS iff it cannot recognize the language  $\{a, b\}^* ab \{a, b\}^*$ .
- Equivalently, EDS is the largest variety of monoids that cannot recognize all 2-testable languages.

# The case of EDS

- Suppose that  $M$  is a  $\Sigma$ -generated finite monoid.
- The Rhodes-Zalcstein theory allows us to associate to each irreducible representation  $\rho: M \rightarrow M_n(\mathbb{Q})$  of  $M$  a finite automaton  $\mathcal{A}(\rho)$  over  $\Sigma$ .
- $0 \in \rho(M)$  iff  $\mathcal{A}(\rho)$  is synchronizing with a sink state.
- The 0-words for the representation are precisely the reset words for the automaton.
- When  $M \in \text{EDS}$ , Almeida and I showed that  $\mathcal{A}(\rho)$  has at most  $n + 1$  states.
- Rystsov showed that, for  $m$ -state synchronizing automata with sink state, there is a synchronizing word of length at most  $m(m - 1)/2$  (and this bound is sharp).
- Consequently,  $f(n) = n(n + 1)/2$  is a superadditive mortality function for EDS.

# The case of EDS

- Suppose that  $M$  is a  $\Sigma$ -generated finite monoid.
- The Rhodes-Zalcstein theory allows us to associate to each irreducible representation  $\rho: M \rightarrow M_n(\mathbb{Q})$  of  $M$  a finite automaton  $\mathcal{A}(\rho)$  over  $\Sigma$ .
- $0 \in \rho(M)$  iff  $\mathcal{A}(\rho)$  is synchronizing with a sink state.
- The 0-words for the representation are precisely the reset words for the automaton.
- When  $M \in \text{EDS}$ , Almeida and I showed that  $\mathcal{A}(\rho)$  has at most  $n + 1$  states.
- Rystsov showed that, for  $m$ -state synchronizing automata with sink state, there is a synchronizing word of length at most  $m(m - 1)/2$  (and this bound is sharp).
- Consequently,  $f(n) = n(n + 1)/2$  is a superadditive mortality function for EDS.

# The case of EDS

- Suppose that  $M$  is a  $\Sigma$ -generated finite monoid.
- The Rhodes-Zalcstein theory allows us to associate to each irreducible representation  $\rho: M \rightarrow M_n(\mathbb{Q})$  of  $M$  a finite automaton  $\mathcal{A}(\rho)$  over  $\Sigma$ .
- $0 \in \rho(M)$  iff  $\mathcal{A}(\rho)$  is synchronizing with a sink state.
- The 0-words for the representation are precisely the reset words for the automaton.
- When  $M \in \text{EDS}$ , Almeida and I showed that  $\mathcal{A}(\rho)$  has at most  $n + 1$  states.
- Rystsov showed that, for  $m$ -state synchronizing automata with sink state, there is a synchronizing word of length at most  $m(m - 1)/2$  (and this bound is sharp).
- Consequently,  $f(n) = n(n + 1)/2$  is a superadditive mortality function for EDS.

# The case of EDS

- Suppose that  $M$  is a  $\Sigma$ -generated finite monoid.
- The Rhodes-Zalcstein theory allows us to associate to each irreducible representation  $\rho: M \rightarrow M_n(\mathbb{Q})$  of  $M$  a finite automaton  $\mathcal{A}(\rho)$  over  $\Sigma$ .
- $0 \in \rho(M)$  iff  $\mathcal{A}(\rho)$  is synchronizing with a sink state.
- The 0-words for the representation are precisely the reset words for the automaton.
- When  $M \in \text{EDS}$ , Almeida and I showed that  $\mathcal{A}(\rho)$  has at most  $n + 1$  states.
- Rystsov showed that, for  $m$ -state synchronizing automata with sink state, there is a synchronizing word of length at most  $m(m - 1)/2$  (and this bound is sharp).
- Consequently,  $f(n) = n(n + 1)/2$  is a superadditive mortality function for EDS.



# The case of EDS

- Suppose that  $M$  is a  $\Sigma$ -generated finite monoid.
- The Rhodes-Zalcstein theory allows us to associate to each irreducible representation  $\rho: M \rightarrow M_n(\mathbb{Q})$  of  $M$  a finite automaton  $\mathcal{A}(\rho)$  over  $\Sigma$ .
- $0 \in \rho(M)$  iff  $\mathcal{A}(\rho)$  is synchronizing with a sink state.
- The 0-words for the representation are precisely the reset words for the automaton.
- When  $M \in \text{EDS}$ , Almeida and I showed that  $\mathcal{A}(\rho)$  has at most  $n + 1$  states.
- Rystsov showed that, for  $m$ -state synchronizing automata with sink state, there is a synchronizing word of length at most  $m(m - 1)/2$  (and this bound is sharp).
- Consequently,  $f(n) = n(n + 1)/2$  is a superadditive mortality function for EDS.

# The case of EDS

- Suppose that  $M$  is a  $\Sigma$ -generated finite monoid.
- The Rhodes-Zalcstein theory allows us to associate to each irreducible representation  $\rho: M \rightarrow M_n(\mathbb{Q})$  of  $M$  a finite automaton  $\mathcal{A}(\rho)$  over  $\Sigma$ .
- $0 \in \rho(M)$  iff  $\mathcal{A}(\rho)$  is synchronizing with a sink state.
- The 0-words for the representation are precisely the reset words for the automaton.
- When  $M \in \text{EDS}$ , Almeida and I showed that  $\mathcal{A}(\rho)$  has at most  $n + 1$  states.
- Rystsov showed that, for  $m$ -state synchronizing automata with sink state, there is a synchronizing word of length at most  $m(m - 1)/2$  (and this bound is sharp).
- Consequently,  $f(n) = n(n + 1)/2$  is a superadditive mortality function for EDS.

# The case of EDS

- Suppose that  $M$  is a  $\Sigma$ -generated finite monoid.
- The Rhodes-Zalcstein theory allows us to associate to each irreducible representation  $\rho: M \rightarrow M_n(\mathbb{Q})$  of  $M$  a finite automaton  $\mathcal{A}(\rho)$  over  $\Sigma$ .
- $0 \in \rho(M)$  iff  $\mathcal{A}(\rho)$  is synchronizing with a sink state.
- The 0-words for the representation are precisely the reset words for the automaton.
- When  $M \in \text{EDS}$ , Almeida and I showed that  $\mathcal{A}(\rho)$  has at most  $n + 1$  states.
- Rystsov showed that, for  $m$ -state synchronizing automata with sink state, there is a synchronizing word of length at most  $m(m - 1)/2$  (and this bound is sharp).
- Consequently,  $f(n) = n(n + 1)/2$  is a superadditive mortality function for EDS.

# The Černý-Pin conjecture for EDS

## Theorem (Almeida, BS)

*If  $\mathcal{A}$  is an  $n$ -state, rank  $r$  automaton with transition monoid in EDS, then there is a word  $w$  of length at most*

$$\frac{(n-r)(n-r+1)}{2}$$

*with  $\text{rk}(w) = r$ .*

- The bound of  $n(n-1)/2$  is sharp for the Černý conjecture.
- Rystsov has an example of an  $n$ -state synchronizing automaton with minimal length reset word of length  $n(n-1)/2$  whose transition monoid has commuting idempotents.
- Our method works much more generally than for EDS.
- For instance  $n(n+1)/2$  is a mortality function for  $M_k(\mathbb{F}_q)$  and the partial transformation monoids  $PT_k$ .

# The Černý-Pin conjecture for EDS

## Theorem (Almeida, BS)

*If  $\mathcal{A}$  is an  $n$ -state, rank  $r$  automaton with transition monoid in EDS, then there is a word  $w$  of length at most*

$$\frac{(n - r)(n - r + 1)}{2}$$

*with  $\text{rk}(w) = r$ .*

- The bound of  $n(n - 1)/2$  is sharp for the Černý conjecture.
- Rystsov has an example of an  $n$ -state synchronizing automaton with minimal length reset word of length  $n(n - 1)/2$  whose transition monoid has commuting idempotents.
- Our method works much more generally than for EDS.
- For instance  $n(n + 1)/2$  is a mortality function for  $M_k(\mathbb{F}_q)$  and the partial transformation monoids  $PT_k$ .

# The Černý-Pin conjecture for EDS

## Theorem (Almeida, BS)

*If  $\mathcal{A}$  is an  $n$ -state, rank  $r$  automaton with transition monoid in EDS, then there is a word  $w$  of length at most*

$$\frac{(n - r)(n - r + 1)}{2}$$

*with  $\text{rk}(w) = r$ .*

- The bound of  $n(n - 1)/2$  is sharp for the Černý conjecture.
- Rystsov has an example of an  $n$ -state synchronizing automaton with minimal length reset word of length  $n(n - 1)/2$  whose transition monoid has commuting idempotents.
- Our method works much more generally than for EDS.
- For instance  $n(n + 1)/2$  is a mortality function for  $M_k(\mathbb{F}_q)$  and the partial transformation monoids  $PT_k$ .

# The Černý-Pin conjecture for EDS

## Theorem (Almeida, BS)

*If  $\mathcal{A}$  is an  $n$ -state, rank  $r$  automaton with transition monoid in EDS, then there is a word  $w$  of length at most*

$$\frac{(n - r)(n - r + 1)}{2}$$

*with  $\text{rk}(w) = r$ .*

- The bound of  $n(n - 1)/2$  is sharp for the Černý conjecture.
- Rystsov has an example of an  $n$ -state synchronizing automaton with minimal length reset word of length  $n(n - 1)/2$  whose transition monoid has commuting idempotents.
- Our method works much more generally than for EDS.
- For instance  $n(n + 1)/2$  is a mortality function for  $M_k(\mathbb{F}_q)$  and the partial transformation monoids  $PT_k$ .

# The Černý-Pin conjecture for EDS

## Theorem (Almeida, BS)

*If  $\mathcal{A}$  is an  $n$ -state, rank  $r$  automaton with transition monoid in EDS, then there is a word  $w$  of length at most*

$$\frac{(n - r)(n - r + 1)}{2}$$

*with  $\text{rk}(w) = r$ .*

- The bound of  $n(n - 1)/2$  is sharp for the Černý conjecture.
- Rystsov has an example of an  $n$ -state synchronizing automaton with minimal length reset word of length  $n(n - 1)/2$  whose transition monoid has commuting idempotents.
- Our method works much more generally than for EDS.
- For instance  $n(n + 1)/2$  is a mortality function for  $M_k(\mathbb{F}_q)$  and the partial transformation monoids  $PT_k$ .



# A universal mortality function

- Given the undecidability of the Matrix Mortality Problem for  $3 \times 3$  integer matrices, it is not altogether clear that a universal mortality function exists.
- On the other hand, Simon/Mandel and Jacob independently proved that there is a recursive bound on the order of a finite  $k$ -generated submonoid of  $M_n(\mathbb{Q})$ .
- If a monoid  $M$  contains 0, then 0 can be represented by a word of length  $|M| - 1$ .
- So if we can remove the dependence on the number of generators, we are done.
- The results of Simon/Mandel and Jacob rely on the solution to the Burnside problem for matrix semigroups.
- For the irreducible case, the number of generators is irrelevant.
- So by working a little harder to get a superadditive bound in the irreducible case, we proved the following theorem.

# A universal mortality function

- Given the undecidability of the Matrix Mortality Problem for  $3 \times 3$  integer matrices, it is not altogether clear that a universal mortality function exists.
- On the other hand, Simon/Mandel and Jacob independently proved that there is a recursive bound on the order of a finite  $k$ -generated submonoid of  $M_n(\mathbb{Q})$ .
- If a monoid  $M$  contains 0, then 0 can be represented by a word of length  $|M| - 1$ .
- So if we can remove the dependence on the number of generators, we are done.
- The results of Simon/Mandel and Jacob rely on the solution to the Burnside problem for matrix semigroups.
- For the irreducible case, the number of generators is irrelevant.
- So by working a little harder to get a superadditive bound in the irreducible case, we proved the following theorem.

# A universal mortality function

- Given the undecidability of the Matrix Mortality Problem for  $3 \times 3$  integer matrices, it is not altogether clear that a universal mortality function exists.
- On the other hand, Simon/Mandel and Jacob independently proved that there is a recursive bound on the order of a finite  $k$ -generated submonoid of  $M_n(\mathbb{Q})$ .
- If a monoid  $M$  contains 0, then 0 can be represented by a word of length  $|M| - 1$ .
- So if we can remove the dependence on the number of generators, we are done.
- The results of Simon/Mandel and Jacob rely on the solution to the Burnside problem for matrix semigroups.
- For the irreducible case, the number of generators is irrelevant.
- So by working a little harder to get a superadditive bound in the irreducible case, we proved the following theorem.

# A universal mortality function

- Given the undecidability of the Matrix Mortality Problem for  $3 \times 3$  integer matrices, it is not altogether clear that a universal mortality function exists.
- On the other hand, Simon/Mandel and Jacob independently proved that there is a recursive bound on the order of a finite  $k$ -generated submonoid of  $M_n(\mathbb{Q})$ .
- If a monoid  $M$  contains 0, then 0 can be represented by a word of length  $|M| - 1$ .
- So if we can remove the dependence on the number of generators, we are done.
- The results of Simon/Mandel and Jacob rely on the solution to the Burnside problem for matrix semigroups.
- For the irreducible case, the number of generators is irrelevant.
- So by working a little harder to get a superadditive bound in the irreducible case, we proved the following theorem.

# A universal mortality function

- Given the undecidability of the Matrix Mortality Problem for  $3 \times 3$  integer matrices, it is not altogether clear that a universal mortality function exists.
- On the other hand, Simon/Mandel and Jacob independently proved that there is a recursive bound on the order of a finite  $k$ -generated submonoid of  $M_n(\mathbb{Q})$ .
- If a monoid  $M$  contains 0, then 0 can be represented by a word of length  $|M| - 1$ .
- So if we can remove the dependence on the number of generators, we are done.
- The results of Simon/Mandel and Jacob rely on the solution to the Burnside problem for matrix semigroups.
- For the irreducible case, the number of generators is irrelevant.
- So by working a little harder to get a superadditive bound in the irreducible case, we proved the following theorem.

# A universal mortality function

- Given the undecidability of the Matrix Mortality Problem for  $3 \times 3$  integer matrices, it is not altogether clear that a universal mortality function exists.
- On the other hand, Simon/Mandel and Jacob independently proved that there is a recursive bound on the order of a finite  $k$ -generated submonoid of  $M_n(\mathbb{Q})$ .
- If a monoid  $M$  contains 0, then 0 can be represented by a word of length  $|M| - 1$ .
- So if we can remove the dependence on the number of generators, we are done.
- The results of Simon/Mandel and Jacob rely on the solution to the Burnside problem for matrix semigroups.
- For the irreducible case, the number of generators is irrelevant.
- So by working a little harder to get a superadditive bound in the irreducible case, we proved the following theorem.

# A universal mortality function

- Given the undecidability of the Matrix Mortality Problem for  $3 \times 3$  integer matrices, it is not altogether clear that a universal mortality function exists.
- On the other hand, Simon/Mandel and Jacob independently proved that there is a recursive bound on the order of a finite  $k$ -generated submonoid of  $M_n(\mathbb{Q})$ .
- If a monoid  $M$  contains 0, then 0 can be represented by a word of length  $|M| - 1$ .
- So if we can remove the dependence on the number of generators, we are done.
- The results of Simon/Mandel and Jacob rely on the solution to the Burnside problem for matrix semigroups.
- For the irreducible case, the number of generators is irrelevant.
- So by working a little harder to get a superadditive bound in the irreducible case, we proved the following theorem.

# A universal mortality function II

## Theorem (Almeida, BS)

*The function*

$$f(n) = \begin{cases} 1 & n = 1 \\ (2n - 1)^{n^2} - 1 & n > 1 \end{cases}$$

*is a superadditive universal mortality function.*

- We know this upper bound is not tight.
- The best lower bound we have is  $n^2$ .
- For aperiodic monoids, we can now prove  $2^n - 1$  is a mortality function (the article in the Proceedings has  $2^{n^2} - 1$ ).



# A universal mortality function II

## Theorem (Almeida, BS)

*The function*

$$f(n) = \begin{cases} 1 & n = 1 \\ (2n - 1)^{n^2} - 1 & n > 1 \end{cases}$$

*is a superadditive universal mortality function.*

- We know this upper bound is not tight.
- The best lower bound we have is  $n^2$ .
- For aperiodic monoids, we can now prove  $2^n - 1$  is a mortality function (the article in the Proceedings has  $2^{n^2} - 1$ ).

# A universal mortality function II

## Theorem (Almeida, BS)

*The function*

$$f(n) = \begin{cases} 1 & n = 1 \\ (2n - 1)^{n^2} - 1 & n > 1 \end{cases}$$

*is a superadditive universal mortality function.*

- We know this upper bound is not tight.
- The best lower bound we have is  $n^2$ .
- For aperiodic monoids, we can now prove  $2^n - 1$  is a mortality function (the article in the Proceedings has  $2^{n^2} - 1$ ).

# A universal mortality function II

## Theorem (Almeida, BS)

*The function*

$$f(n) = \begin{cases} 1 & n = 1 \\ (2n - 1)^{n^2} - 1 & n > 1 \end{cases}$$

*is a superadditive universal mortality function.*

- We know this upper bound is not tight.
- The best lower bound we have is  $n^2$ .
- For aperiodic monoids, we can now prove  $2^n - 1$  is a mortality function (the article in the Proceedings has  $2^{n^2} - 1$ ).

And Now For Something  
Completely Different  
A Larch...

And Now For Something  
Completely Different

A Larch...

And Now For Something  
Completely Different  
A Larch...

- Now I want to focus on some aspects of Černý's conjecture related to Pin and Dubuc's theorems.
- This part of the talk is not in the Proceedings and is my own work.

## Theorem (Pin '78)

*Let  $\mathcal{A} = (Q, \Sigma)$  be a synchronizing automaton so that  $|Q|$  is a prime  $p$  and some element of  $\Sigma$  cyclically permutes  $Q$ . Then:*

1.  *$\mathcal{A}$  is synchronizing if and only if  $\Sigma$  contains a non-permutation;*
2. *In this case,  $\mathcal{A}$  has a reset word of length at most  $(p - 1)^2$ .*

- Now I want to focus on some aspects of Černý's conjecture related to Pin and Dubuc's theorems.
- This part of the talk is not in the Proceedings and is my own work.

## Theorem (Pin '78)

*Let  $\mathcal{A} = (Q, \Sigma)$  be a synchronizing automaton so that  $|Q|$  is a prime  $p$  and some element of  $\Sigma$  cyclically permutes  $Q$ . Then:*

- 1  $\mathcal{A}$  is synchronizing if and only if  $\Sigma$  contains a non-permutation;*
- 2 In this case,  $\mathcal{A}$  has a reset word of length at most  $(p - 1)^2$ .*



- Now I want to focus on some aspects of Černý's conjecture related to Pin and Dubuc's theorems.
- This part of the talk is not in the Proceedings and is my own work.

## Theorem (Pin '78)

*Let  $\mathcal{A} = (Q, \Sigma)$  be a synchronizing automaton so that  $|Q|$  is a prime  $p$  and some element of  $\Sigma$  cyclically permutes  $Q$ . Then:*

- 1  $\mathcal{A}$  is synchronizing if and only if  $\Sigma$  contains a non-permutation;*
- 2 In this case,  $\mathcal{A}$  has a reset word of length at most  $(p - 1)^2$ .*

- Now I want to focus on some aspects of Černý's conjecture related to Pin and Dubuc's theorems.
- This part of the talk is not in the Proceedings and is my own work.

## Theorem (Pin '78)

*Let  $\mathcal{A} = (Q, \Sigma)$  be a synchronizing automaton so that  $|Q|$  is a prime  $p$  and some element of  $\Sigma$  cyclically permutes  $Q$ . Then:*

- 1  *$\mathcal{A}$  is synchronizing if and only if  $\Sigma$  contains a non-permutation;*
- 2 *In this case,  $\mathcal{A}$  has a reset word of length at most  $(p - 1)^2$ .*

- Now I want to focus on some aspects of Černý's conjecture related to Pin and Dubuc's theorems.
- This part of the talk is not in the Proceedings and is my own work.

## Theorem (Pin '78)

*Let  $\mathcal{A} = (Q, \Sigma)$  be a synchronizing automaton so that  $|Q|$  is a prime  $p$  and some element of  $\Sigma$  cyclically permutes  $Q$ . Then:*

- 1  *$\mathcal{A}$  is synchronizing if and only if  $\Sigma$  contains a non-permutation;*
- 2 *In this case,  $\mathcal{A}$  has a reset word of length at most  $(p - 1)^2$ .*

# Synchronizing groups

- Motivated by the first part of Pin's Theorem, I defined in 2004 the notion of a synchronizing group.
- A **permutation group**  $G \subseteq S_n$  is called **synchronizing** if, for all non-permutations  $t$  of  $\{1, \dots, n\}$ , the automaton  $(\{1, \dots, n\}, G \cup \{t\})$  is synchronizing.
- Pin's Theorem implies that cyclic groups of prime order are synchronizing.
- It is easy to see that **2-transitive groups** are synchronizing.
- With Arnold, I proved synchronizing groups are primitive and gave a sufficient condition for a group to be synchronizing in terms of representation theory that covers the above results.
- João Araújo independently came up with the notion in 2006 and found a beautiful group theoretic reformulation.
- Synchronizing groups have recently received quite a bit of attention from prominent group theorists including Peter Cameron, Peter Neumann and Jan Saxl.

# Synchronizing groups

- Motivated by the first part of Pin's Theorem, I defined in 2004 the notion of a synchronizing group.
- A **permutation group**  $G \subseteq S_n$  is called **synchronizing** if, for all non-permutations  $t$  of  $\{1, \dots, n\}$ , the automaton  $(\{1, \dots, n\}, G \cup \{t\})$  is synchronizing.
- Pin's Theorem implies that cyclic groups of prime order are synchronizing.
- It is easy to see that **2-transitive groups** are synchronizing.
- With Arnold, I proved synchronizing groups are **primitive** and gave a sufficient condition for a group to be synchronizing in terms of representation theory that covers the above results.
- João Araújo independently came up with the notion in 2006 and found a beautiful group theoretic reformulation.
- Synchronizing groups have recently received quite a bit of attention from prominent group theorists including Peter Cameron, Peter Neumann and Jan Saxl.

# Synchronizing groups

- Motivated by the first part of Pin's Theorem, I defined in 2004 the notion of a synchronizing group.
- A **permutation group**  $G \subseteq S_n$  is called **synchronizing** if, for all non-permutations  $t$  of  $\{1, \dots, n\}$ , the automaton  $(\{1, \dots, n\}, G \cup \{t\})$  is synchronizing.
- Pin's Theorem implies that cyclic groups of prime order are synchronizing.
- It is easy to see that **2-transitive groups** are synchronizing.
- With Arnold, I proved synchronizing groups are **primitive** and gave a sufficient condition for a group to be synchronizing in terms of representation theory that covers the above results.
- João Araújo independently came up with the notion in 2006 and found a beautiful group theoretic reformulation.
- Synchronizing groups have recently received quite a bit of attention from prominent group theorists including Peter Cameron, Peter Neumann and Jan Saxl.

# Synchronizing groups

- Motivated by the first part of Pin's Theorem, I defined in 2004 the notion of a synchronizing group.
- A **permutation group**  $G \subseteq S_n$  is called **synchronizing** if, for all non-permutations  $t$  of  $\{1, \dots, n\}$ , the automaton  $(\{1, \dots, n\}, G \cup \{t\})$  is synchronizing.
- Pin's Theorem implies that cyclic groups of prime order are synchronizing.
- It is easy to see that **2-transitive groups** are synchronizing.
- With Arnold, I proved synchronizing groups are **primitive** and gave a sufficient condition for a group to be synchronizing in terms of representation theory that covers the above results.
- João Araújo independently came up with the notion in 2006 and found a beautiful group theoretic reformulation.
- Synchronizing groups have recently received quite a bit of attention from prominent group theorists including Peter Cameron, Peter Neumann and Jan Saxl.

# Synchronizing groups

- Motivated by the first part of Pin's Theorem, I defined in 2004 the notion of a synchronizing group.
- A **permutation group**  $G \subseteq S_n$  is called **synchronizing** if, for all non-permutations  $t$  of  $\{1, \dots, n\}$ , the automaton  $(\{1, \dots, n\}, G \cup \{t\})$  is synchronizing.
- Pin's Theorem implies that cyclic groups of prime order are synchronizing.
- It is easy to see that **2-transitive groups** are synchronizing.
- With Arnold, I proved synchronizing groups are **primitive** and gave a sufficient condition for a group to be synchronizing in terms of representation theory that covers the above results.
- João Araújo independently came up with the notion in 2006 and found a beautiful group theoretic reformulation.
- Synchronizing groups have recently received quite a bit of attention from prominent group theorists including Peter Cameron, Peter Neumann and Jan Saxl.



# Synchronizing groups

- Motivated by the first part of Pin's Theorem, I defined in 2004 the notion of a synchronizing group.
- A **permutation group**  $G \subseteq S_n$  is called **synchronizing** if, for all non-permutations  $t$  of  $\{1, \dots, n\}$ , the automaton  $(\{1, \dots, n\}, G \cup \{t\})$  is synchronizing.
- Pin's Theorem implies that cyclic groups of prime order are synchronizing.
- It is easy to see that **2-transitive groups** are synchronizing.
- With Arnold, I proved synchronizing groups are **primitive** and gave a sufficient condition for a group to be synchronizing in terms of representation theory that covers the above results.
- João Araújo independently came up with the notion in 2006 and found a beautiful group theoretic reformulation.
- Synchronizing groups have recently received quite a bit of attention from prominent group theorists including Peter Cameron, Peter Neumann and Jan Saxl.

# Synchronizing groups

- Motivated by the first part of Pin's Theorem, I defined in 2004 the notion of a synchronizing group.
- A **permutation group**  $G \subseteq S_n$  is called **synchronizing** if, for all non-permutations  $t$  of  $\{1, \dots, n\}$ , the automaton  $(\{1, \dots, n\}, G \cup \{t\})$  is synchronizing.
- Pin's Theorem implies that cyclic groups of prime order are synchronizing.
- It is easy to see that **2-transitive groups** are synchronizing.
- With Arnold, I proved synchronizing groups are **primitive** and gave a sufficient condition for a group to be synchronizing in terms of representation theory that covers the above results.
- João Araújo independently came up with the notion in 2006 and found a beautiful group theoretic reformulation.
- Synchronizing groups have recently received quite a bit of attention from prominent group theorists including Peter Cameron, Peter Neumann and Jan Saxl.

- Dubuc extended the second part of Pin's Theorem to arbitrary automata containing a cyclic permutation via an ingenious linear algebraic argument.

## Theorem (Dubuc '98)

*Let  $\mathcal{A} = (Q, \Sigma)$  be a synchronizing automaton on  $n$  states such that  $\Sigma$  contains a cyclic permutation of the states. Then  $\mathcal{A}$  has a reset word of length at most  $(n - 1)^2$ .*

- The results of Dubuc and Pin make it natural to consider more general groups than cyclic groups.

- Dubuc extended the second part of Pin's Theorem to arbitrary automata containing a cyclic permutation via an ingenious linear algebraic argument.

## Theorem (Dubuc '98)

*Let  $\mathcal{A} = (Q, \Sigma)$  be a synchronizing automaton on  $n$  states such that  $\Sigma$  contains a cyclic permutation of the states. Then  $\mathcal{A}$  has a reset word of length at most  $(n - 1)^2$ .*

- The results of Dubuc and Pin make it natural to consider more general groups than cyclic groups.

- Dubuc extended the second part of Pin's Theorem to arbitrary automata containing a cyclic permutation via an ingenious linear algebraic argument.

## Theorem (Dubuc '98)

*Let  $\mathcal{A} = (Q, \Sigma)$  be a synchronizing automaton on  $n$  states such that  $\Sigma$  contains a cyclic permutation of the states. Then  $\mathcal{A}$  has a reset word of length at most  $(n - 1)^2$ .*

- The results of Dubuc and Pin make it natural to consider more general groups than cyclic groups.

# Černý Cayley graphs

- Let  $G$  be a group of order  $n$  and  $\Delta$  a generating set of  $G$ .
- The automaton  $(G, \Delta)$  is called the **Cayley graph** of  $G$  with respect to  $\Delta$ . A typical transition is of the form  $g \xrightarrow{a} ga$  with  $g \in G$ ,  $a \in \Sigma$ .
- Let us say that an automaton  $\mathcal{A}$  **contains** the Cayley graph  $(G, \Delta)$  if  $\mathcal{A} = (G, \Sigma)$  where  $\Delta \subseteq \Sigma$ .
- So  $\mathcal{A}$  is obtained from the Cayley graph by adding new transitions but no new states.
- Call  $(G, \Delta)$  a Černý Cayley graph if every synchronizing automaton containing it has a reset word of length at most  $(n-1)^2$ .
- Let's say  $G$  is a Černý group if all its Cayley graphs are Černý Cayley graphs.
- Dubuc's theorem says that  $(\mathbb{Z}_n, \{1\})$  is a Černý Cayley graph.
- Cyclic groups of prime power order are Černý groups.

# Černý Cayley graphs

- Let  $G$  be a group of order  $n$  and  $\Delta$  a generating set of  $G$ .
- The automaton  $(G, \Delta)$  is called the **Cayley graph** of  $G$  with respect to  $\Delta$ . A typical transition is of the form  $g \xrightarrow{a} ga$  with  $g \in G$ ,  $a \in \Sigma$ .
- Let us say that an automaton  $\mathcal{A}$  **contains** the Cayley graph  $(G, \Delta)$  if  $\mathcal{A} = (G, \Sigma)$  where  $\Delta \subseteq \Sigma$ .
- So  $\mathcal{A}$  is obtained from the Cayley graph by adding new transitions but no new states.
- Call  $(G, \Delta)$  a **Černý Cayley graph** if every synchronizing automaton containing it has a reset word of length at most  $(n - 1)^2$ .
- Let's say  $G$  is a **Černý group** if all its Cayley graphs are Černý Cayley graphs.
- Dubuc's theorem says that  $(\mathbb{Z}_n, \{1\})$  is a Černý Cayley graph.
- Cyclic groups of prime power order are Černý groups.

# Černý Cayley graphs

- Let  $G$  be a group of order  $n$  and  $\Delta$  a generating set of  $G$ .
- The automaton  $(G, \Delta)$  is called the **Cayley graph** of  $G$  with respect to  $\Delta$ . A typical transition is of the form  $g \xrightarrow{a} ga$  with  $g \in G$ ,  $a \in \Sigma$ .
- Let us say that an automaton  $\mathcal{A}$  **contains** the Cayley graph  $(G, \Delta)$  if  $\mathcal{A} = (G, \Sigma)$  where  $\Delta \subseteq \Sigma$ .
- So  $\mathcal{A}$  is obtained from the Cayley graph by adding new transitions but no new states.
- Call  $(G, \Delta)$  a **Černý Cayley graph** if every synchronizing automaton containing it has a reset word of length at most  $(n - 1)^2$ .
- Let's say  $G$  is a **Černý group** if all its Cayley graphs are Černý Cayley graphs.
- Dubuc's theorem says that  $(\mathbb{Z}_n, \{1\})$  is a Černý Cayley graph.
- Cyclic groups of prime power order are Černý groups.



# Černý Cayley graphs

- Let  $G$  be a group of order  $n$  and  $\Delta$  a generating set of  $G$ .
- The automaton  $(G, \Delta)$  is called the **Cayley graph** of  $G$  with respect to  $\Delta$ . A typical transition is of the form  $g \xrightarrow{a} ga$  with  $g \in G$ ,  $a \in \Sigma$ .
- Let us say that an automaton  $\mathcal{A}$  **contains** the Cayley graph  $(G, \Delta)$  if  $\mathcal{A} = (G, \Sigma)$  where  $\Delta \subseteq \Sigma$ .
- So  $\mathcal{A}$  is obtained from the Cayley graph by adding new transitions but no new states.
- Call  $(G, \Delta)$  a **Černý Cayley graph** if every synchronizing automaton containing it has a reset word of length at most  $(n - 1)^2$ .
- Let's say  $G$  is a **Černý group** if all its Cayley graphs are Černý Cayley graphs.
- Dubuc's theorem says that  $(\mathbb{Z}_n, \{1\})$  is a Černý Cayley graph.
- Cyclic groups of prime power order are Černý groups.

# Černý Cayley graphs

- Let  $G$  be a group of order  $n$  and  $\Delta$  a generating set of  $G$ .
- The automaton  $(G, \Delta)$  is called the **Cayley graph** of  $G$  with respect to  $\Delta$ . A typical transition is of the form  $g \xrightarrow{a} ga$  with  $g \in G$ ,  $a \in \Sigma$ .
- Let us say that an automaton  $\mathcal{A}$  **contains** the Cayley graph  $(G, \Delta)$  if  $\mathcal{A} = (G, \Sigma)$  where  $\Delta \subseteq \Sigma$ .
- So  $\mathcal{A}$  is obtained from the Cayley graph by adding new transitions but no new states.
- Call  $(G, \Delta)$  a **Černý Cayley graph** if every synchronizing automaton containing it has a reset word of length at most  $(n - 1)^2$ .
- Let's say  $G$  is a **Černý group** if all its Cayley graphs are Černý Cayley graphs.
- Dubuc's theorem says that  $(\mathbb{Z}_n, \{1\})$  is a Černý Cayley graph.
- Cyclic groups of prime power order are Černý groups.

# Černý Cayley graphs

- Let  $G$  be a group of order  $n$  and  $\Delta$  a generating set of  $G$ .
- The automaton  $(G, \Delta)$  is called the **Cayley graph** of  $G$  with respect to  $\Delta$ . A typical transition is of the form  $g \xrightarrow{a} ga$  with  $g \in G$ ,  $a \in \Sigma$ .
- Let us say that an automaton  $\mathcal{A}$  **contains** the Cayley graph  $(G, \Delta)$  if  $\mathcal{A} = (G, \Sigma)$  where  $\Delta \subseteq \Sigma$ .
- So  $\mathcal{A}$  is obtained from the Cayley graph by adding new transitions but no new states.
- Call  $(G, \Delta)$  a **Černý Cayley graph** if every synchronizing automaton containing it has a reset word of length at most  $(n - 1)^2$ .
- Let's say  $G$  is a **Černý group** if all its Cayley graphs are Černý Cayley graphs.
- Dubuc's theorem says that  $(\mathbb{Z}_n, \{1\})$  is a Černý Cayley graph.
- Cyclic groups of prime power order are Černý groups.

# Černý Cayley graphs

- Let  $G$  be a group of order  $n$  and  $\Delta$  a generating set of  $G$ .
- The automaton  $(G, \Delta)$  is called the **Cayley graph** of  $G$  with respect to  $\Delta$ . A typical transition is of the form  $g \xrightarrow{a} ga$  with  $g \in G$ ,  $a \in \Sigma$ .
- Let us say that an automaton  $\mathcal{A}$  **contains** the Cayley graph  $(G, \Delta)$  if  $\mathcal{A} = (G, \Sigma)$  where  $\Delta \subseteq \Sigma$ .
- So  $\mathcal{A}$  is obtained from the Cayley graph by adding new transitions but no new states.
- Call  $(G, \Delta)$  a **Černý Cayley graph** if every synchronizing automaton containing it has a reset word of length at most  $(n - 1)^2$ .
- Let's say  $G$  is a **Černý group** if all its Cayley graphs are Černý Cayley graphs.
- Dubuc's theorem says that  $(\mathbb{Z}_n, \{1\})$  is a Černý Cayley graph.
- Cyclic groups of prime power order are Černý groups.

# Černý Cayley graphs

- Let  $G$  be a group of order  $n$  and  $\Delta$  a generating set of  $G$ .
- The automaton  $(G, \Delta)$  is called the **Cayley graph** of  $G$  with respect to  $\Delta$ . A typical transition is of the form  $g \xrightarrow{a} ga$  with  $g \in G$ ,  $a \in \Sigma$ .
- Let us say that an automaton  $\mathcal{A}$  **contains** the Cayley graph  $(G, \Delta)$  if  $\mathcal{A} = (G, \Sigma)$  where  $\Delta \subseteq \Sigma$ .
- So  $\mathcal{A}$  is obtained from the Cayley graph by adding new transitions but no new states.
- Call  $(G, \Delta)$  a **Černý Cayley graph** if every synchronizing automaton containing it has a reset word of length at most  $(n - 1)^2$ .
- Let's say  $G$  is a **Černý group** if all its Cayley graphs are Černý Cayley graphs.
- Dubuc's theorem says that  $(\mathbb{Z}_n, \{1\})$  is a Černý Cayley graph.
- Cyclic groups of prime power order are Černý groups.

# Rystsov's Theorem

- The above notion was implicitly considered by Rystsov.
- In 1995, he proved a synchronizing automaton containing the Cayley graph of a group of order  $n$  admits a reset word of length  $\leq 2(n-1)^2$ .
- He proved in fact a slightly better result.
- Let  $(G, \Delta)$  be a Cayley graph with  $|G| = n > 1$ .
- Define  $\text{diam}_\Delta(G)$  to be the least  $m$  so that any two states of  $(G, \Delta)$  can be connected by a word of length at most  $m$ .
- $1 \leq \text{diam}_\Delta(G) \leq n-1$ .

## Theorem (Rystsov '95)

*A synchronizing automaton containing the Cayley graph  $(G, \Delta)$  has a reset word of length at most  $1 + (n-1 + \text{diam}_\Delta(G))(n-2)$ .*

- $(n-1)^2 = 1 + n(n-2)$ .

# Rystsov's Theorem

- The above notion was implicitly considered by Rystsov.
- In 1995, he proved a synchronizing automaton containing the Cayley graph of a group of order  $n$  admits a reset word of length  $\leq 2(n-1)^2$ .
- He proved in fact a slightly better result.
- Let  $(G, \Delta)$  be a Cayley graph with  $|G| = n > 1$ .
- Define  $\text{diam}_\Delta(G)$  to be the least  $m$  so that any two states of  $(G, \Delta)$  can be connected by a word of length at most  $m$ .
- $1 \leq \text{diam}_\Delta(G) \leq n-1$ .

## Theorem (Rystsov '95)

*A synchronizing automaton containing the Cayley graph  $(G, \Delta)$  has a reset word of length at most  $1 + (n-1 + \text{diam}_\Delta(G))(n-2)$ .*

- $(n-1)^2 = 1 + n(n-2)$ .

# Rystsov's Theorem

- The above notion was implicitly considered by Rystsov.
- In 1995, he proved a synchronizing automaton containing the Cayley graph of a group of order  $n$  admits a reset word of length  $\leq 2(n-1)^2$ .
- He proved in fact a slightly better result.
- Let  $(G, \Delta)$  be a Cayley graph with  $|G| = n > 1$ .
- Define  $\text{diam}_\Delta(G)$  to be the least  $m$  so that any two states of  $(G, \Delta)$  can be connected by a word of length at most  $m$ .
- $1 \leq \text{diam}_\Delta(G) \leq n-1$ .

## Theorem (Rystsov '95)

*A synchronizing automaton containing the Cayley graph  $(G, \Delta)$  has a reset word of length at most  $1 + (n-1 + \text{diam}_\Delta(G))(n-2)$ .*

- $(n-1)^2 = 1 + n(n-2)$ .



# Rystsov's Theorem

- The above notion was implicitly considered by Rystsov.
- In 1995, he proved a synchronizing automaton containing the Cayley graph of a group of order  $n$  admits a reset word of length  $\leq 2(n-1)^2$ .
- He proved in fact a slightly better result.
- Let  $(G, \Delta)$  be a Cayley graph with  $|G| = n > 1$ .
- Define  $\text{diam}_\Delta(G)$  to be the least  $m$  so that any two states of  $(G, \Delta)$  can be connected by a word of length at most  $m$ .
- $1 \leq \text{diam}_\Delta(G) \leq n-1$ .

## Theorem (Rystsov '95)

*A synchronizing automaton containing the Cayley graph  $(G, \Delta)$  has a reset word of length at most  $1 + (n-1 + \text{diam}_\Delta(G))(n-2)$ .*

- $(n-1)^2 = 1 + n(n-2)$ .

# Rystsov's Theorem

- The above notion was implicitly considered by Rystsov.
- In 1995, he proved a synchronizing automaton containing the Cayley graph of a group of order  $n$  admits a reset word of length  $\leq 2(n-1)^2$ .
- He proved in fact a slightly better result.
- Let  $(G, \Delta)$  be a Cayley graph with  $|G| = n > 1$ .
- Define  $\text{diam}_\Delta(G)$  to be the least  $m$  so that any two states of  $(G, \Delta)$  can be connected by a word of length at most  $m$ .
- $1 \leq \text{diam}_\Delta(G) \leq n-1$ .

## Theorem (Rystsov '95)

*A synchronizing automaton containing the Cayley graph  $(G, \Delta)$  has a reset word of length at most  $1 + (n-1 + \text{diam}_\Delta(G))(n-2)$ .*

- $(n-1)^2 = 1 + n(n-2)$ .

# Rystsov's Theorem

- The above notion was implicitly considered by Rystsov.
- In 1995, he proved a synchronizing automaton containing the Cayley graph of a group of order  $n$  admits a reset word of length  $\leq 2(n-1)^2$ .
- He proved in fact a slightly better result.
- Let  $(G, \Delta)$  be a Cayley graph with  $|G| = n > 1$ .
- Define  $\text{diam}_\Delta(G)$  to be the least  $m$  so that any two states of  $(G, \Delta)$  can be connected by a word of length at most  $m$ .
- $1 \leq \text{diam}_\Delta(G) \leq n-1$ .

## Theorem (Rystsov '95)

*A synchronizing automaton containing the Cayley graph  $(G, \Delta)$  has a reset word of length at most  $1 + (n-1 + \text{diam}_\Delta(G))(n-2)$ .*

- $(n-1)^2 = 1 + n(n-2)$ .

# Rystsov's Theorem

- The above notion was implicitly considered by Rystsov.
- In 1995, he proved a synchronizing automaton containing the Cayley graph of a group of order  $n$  admits a reset word of length  $\leq 2(n-1)^2$ .
- He proved in fact a slightly better result.
- Let  $(G, \Delta)$  be a Cayley graph with  $|G| = n > 1$ .
- Define  $\text{diam}_\Delta(G)$  to be the least  $m$  so that any two states of  $(G, \Delta)$  can be connected by a word of length at most  $m$ .
- $1 \leq \text{diam}_\Delta(G) \leq n-1$ .

## Theorem (Rystsov '95)

*A synchronizing automaton containing the Cayley graph  $(G, \Delta)$  has a reset word of length at most  $1 + (n-1 + \text{diam}_\Delta(G))(n-2)$ .*

•  $(n-1)^2 = 1 + n(n-2)$ .

# Rystsov's Theorem

- The above notion was implicitly considered by Rystsov.
- In 1995, he proved a synchronizing automaton containing the Cayley graph of a group of order  $n$  admits a reset word of length  $\leq 2(n-1)^2$ .
- He proved in fact a slightly better result.
- Let  $(G, \Delta)$  be a Cayley graph with  $|G| = n > 1$ .
- Define  $\text{diam}_\Delta(G)$  to be the least  $m$  so that any two states of  $(G, \Delta)$  can be connected by a word of length at most  $m$ .
- $1 \leq \text{diam}_\Delta(G) \leq n-1$ .

## Theorem (Rystsov '95)

*A synchronizing automaton containing the Cayley graph  $(G, \Delta)$  has a reset word of length at most  $1 + (n-1 + \text{diam}_\Delta(G))(n-2)$ .*

- $(n-1)^2 = 1 + n(n-2)$ .

# Our goal

- Recall Rystsov's bound is  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  and  $(n - 1)^2 = 1 + n(n - 2)$ .
- So Rystsov's bound only achieves the Černý bound when the diameter is 1, i.e., all non-trivial elements of  $G$  belong to the generating set.
- We aim to improve his bound so that in many cases we achieve the Černý bound.
- Even when we do not achieve the Černý bound with our main result, our techniques often suffice to establish a family of Cayley graphs is Černý.
- Our results lead to several new families of Černý groups.
- Our main tool is still representation theory.

# Our goal

- Recall Rystsov's bound is  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  and  $(n - 1)^2 = 1 + n(n - 2)$ .
- So Rystsov's bound only achieves the Černý bound when the diameter is 1, i.e., all non-trivial elements of  $G$  belong to the generating set.
- We aim to improve his bound so that in many cases we achieve the Černý bound.
- Even when we do not achieve the Černý bound with our main result, our techniques often suffice to establish a family of Cayley graphs is Černý.
- Our results lead to several new families of Černý groups.
- Our main tool is still representation theory.

# Our goal

- Recall Rystsov's bound is  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  and  $(n - 1)^2 = 1 + n(n - 2)$ .
- So Rystsov's bound only achieves the Černý bound when the diameter is 1, i.e., all non-trivial elements of  $G$  belong to the generating set.
- We aim to improve his bound so that in many cases we achieve the Černý bound.
- Even when we do not achieve the Černý bound with our main result, our techniques often suffice to establish a family of Cayley graphs is Černý.
- Our results lead to several new families of Černý groups.
- Our main tool is still representation theory.



# Our goal

- Recall Rystsov's bound is  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  and  $(n - 1)^2 = 1 + n(n - 2)$ .
- So Rystsov's bound only achieves the Černý bound when the diameter is 1, i.e., all non-trivial elements of  $G$  belong to the generating set.
- We aim to improve his bound so that in many cases we achieve the Černý bound.
- Even when we do not achieve the Černý bound with our main result, our techniques often suffice to establish a family of Cayley graphs is Černý.
- Our results lead to several new families of Černý groups.
- Our main tool is still representation theory.

# Our goal

- Recall Rystsov's bound is  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  and  $(n - 1)^2 = 1 + n(n - 2)$ .
- So Rystsov's bound only achieves the Černý bound when the diameter is 1, i.e., all non-trivial elements of  $G$  belong to the generating set.
- We aim to improve his bound so that in many cases we achieve the Černý bound.
- Even when we do not achieve the Černý bound with our main result, our techniques often suffice to establish a family of Cayley graphs is Černý.
- Our results lead to several new families of Černý groups.
- Our main tool is still representation theory.

# Our goal

- Recall Rystsov's bound is  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  and  $(n - 1)^2 = 1 + n(n - 2)$ .
- So Rystsov's bound only achieves the Černý bound when the diameter is 1, i.e., all non-trivial elements of  $G$  belong to the generating set.
- We aim to improve his bound so that in many cases we achieve the Černý bound.
- Even when we do not achieve the Černý bound with our main result, our techniques often suffice to establish a family of Cayley graphs is Černý.
- Our results lead to several new families of Černý groups.
- Our main tool is still representation theory.

# Irreducible representations of groups

- We shall call an irreducible representation of a group an **irrep**.
- For groups, an arbitrary representation is a direct sum of irreps, which is not the case for monoids.
- If  $|G| = n$ , then the degree of any irrep is between 1 and  $n - 1$ .

## Definition

For a finite group  $G$ , define  $m(G)$  to be the maximal degree of an irrep of  $G$  over  $\mathbb{Q}$ .

- $1 \leq m(G) \leq |G| - 1$ .

# Irreducible representations of groups

- We shall call an irreducible representation of a group an **irrep**.
- For groups, an arbitrary representation is a direct sum of irreps, which is not the case for monoids.
- If  $|G| = n$ , then the degree of any irrep is between 1 and  $n - 1$ .

## Definition

For a finite group  $G$ , define  $m(G)$  to be the maximal degree of an irrep of  $G$  over  $\mathbb{Q}$ .

- $1 \leq m(G) \leq |G| - 1$ .

# Irreducible representations of groups

- We shall call an irreducible representation of a group an **irrep**.
- For groups, an arbitrary representation is a direct sum of irreps, which is not the case for monoids.
- If  $|G| = n$ , then the degree of any irrep is between 1 and  $n - 1$ .

## Definition

For a finite group  $G$ , define  $m(G)$  to be the maximal degree of an irrep of  $G$  over  $\mathbb{Q}$ .

- $1 \leq m(G) \leq |G| - 1$ .

# Irreducible representations of groups

- We shall call an irreducible representation of a group an **irrep**.
- For groups, an arbitrary representation is a direct sum of irreps, which is not the case for monoids.
- If  $|G| = n$ , then the degree of any irrep is between 1 and  $n - 1$ .

## Definition

For a finite group  $G$ , define  $m(G)$  to be the maximal degree of an irrep of  $G$  over  $\mathbb{Q}$ .

- $1 \leq m(G) \leq |G| - 1$ .

# Irreducible representations of groups

- We shall call an irreducible representation of a group an **irrep**.
- For groups, an arbitrary representation is a direct sum of irreps, which is not the case for monoids.
- If  $|G| = n$ , then the degree of any irrep is between 1 and  $n - 1$ .

## Definition

For a finite group  $G$ , define  $m(G)$  to be the maximal degree of an irrep of  $G$  over  $\mathbb{Q}$ .

- $1 \leq m(G) \leq |G| - 1$ .



# The main result

## Theorem (BS)

*Let  $(G, \Delta)$  be a Cayley graph of a group of order  $n$ . Then any synchronizing automaton containing  $(G, \Delta)$  admits a reset word of length at most*

$$1 + (n - m(G) + \text{diam}_\Delta(G))(n - 2).$$

*In particular, if  $\text{diam}_\Delta(G) \leq m(G)$ , then  $(G, \Delta)$  is a Černý Cayley graph.*

- The last statement follows since  $(n - 1)^2 = 1 + n(n - 2)$ .
- $m(G) = 1$  iff  $G \cong \mathbb{Z}_2^k$  for some  $k$ .
- So we beat Rystsov's bound of  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  in essentially all cases.

# The main result

## Theorem (BS)

*Let  $(G, \Delta)$  be a Cayley graph of a group of order  $n$ . Then any synchronizing automaton containing  $(G, \Delta)$  admits a reset word of length at most*

$$1 + (n - m(G) + \text{diam}_\Delta(G))(n - 2).$$

*In particular, if  $\text{diam}_\Delta(G) \leq m(G)$ , then  $(G, \Delta)$  is a Černý Cayley graph.*

- The last statement follows since  $(n - 1)^2 = 1 + n(n - 2)$ .
- $m(G) = 1$  iff  $G \cong \mathbb{Z}_2^k$  for some  $k$ .
- So we beat Rystsov's bound of  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  in essentially all cases.

# The main result

## Theorem (BS)

*Let  $(G, \Delta)$  be a Cayley graph of a group of order  $n$ . Then any synchronizing automaton containing  $(G, \Delta)$  admits a reset word of length at most*

$$1 + (n - m(G) + \text{diam}_\Delta(G))(n - 2).$$

*In particular, if  $\text{diam}_\Delta(G) \leq m(G)$ , then  $(G, \Delta)$  is a Černý Cayley graph.*

- The last statement follows since  $(n - 1)^2 = 1 + n(n - 2)$ .
- $m(G) = 1$  iff  $G \cong \mathbb{Z}_2^k$  for some  $k$ .
- So we beat Rystsov's bound of  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  in essentially all cases.

# The main result

## Theorem (BS)

*Let  $(G, \Delta)$  be a Cayley graph of a group of order  $n$ . Then any synchronizing automaton containing  $(G, \Delta)$  admits a reset word of length at most*

$$1 + (n - m(G) + \text{diam}_\Delta(G))(n - 2).$$

*In particular, if  $\text{diam}_\Delta(G) \leq m(G)$ , then  $(G, \Delta)$  is a Černý Cayley graph.*

- The last statement follows since  $(n - 1)^2 = 1 + n(n - 2)$ .
- $m(G) = 1$  iff  $G \cong \mathbb{Z}_2^k$  for some  $k$ .
- So we beat Rystsov's bound of  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  in essentially all cases.

# The main result

## Theorem (BS)

*Let  $(G, \Delta)$  be a Cayley graph of a group of order  $n$ . Then any synchronizing automaton containing  $(G, \Delta)$  admits a reset word of length at most*

$$1 + (n - m(G) + \text{diam}_\Delta(G))(n - 2).$$

*In particular, if  $\text{diam}_\Delta(G) \leq m(G)$ , then  $(G, \Delta)$  is a Černý Cayley graph.*

- The last statement follows since  $(n - 1)^2 = 1 + n(n - 2)$ .
- $m(G) = 1$  iff  $G \cong \mathbb{Z}_2^k$  for some  $k$ .
- So we beat Rystsov's bound of  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  in essentially all cases.

# The main result

## Theorem (BS)

*Let  $(G, \Delta)$  be a Cayley graph of a group of order  $n$ . Then any synchronizing automaton containing  $(G, \Delta)$  admits a reset word of length at most*

$$1 + (n - m(G) + \text{diam}_\Delta(G))(n - 2).$$

*In particular, if  $\text{diam}_\Delta(G) \leq m(G)$ , then  $(G, \Delta)$  is a Černý Cayley graph.*

- The last statement follows since  $(n - 1)^2 = 1 + n(n - 2)$ .
- $m(G) = 1$  iff  $G \cong \mathbb{Z}_2^k$  for some  $k$ .
- So we beat Rystsov's bound of  $1 + (n - 1 + \text{diam}_\Delta(G))(n - 2)$  in essentially all cases.

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
- Then  $\text{diam}_{\{p,q\}}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
- $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
- So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
- This does not follow from Dubuc's result.

# Cyclic groups

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
- Then  $\text{diam}_{\{p,q\}}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
- $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
- So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
- This does not follow from Dubuc's result.



# Cyclic groups

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
- Then  $\text{diam}_{\{p,q\}}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
- $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
- So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
- This does not follow from Dubuc's result.

# Cyclic groups

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
- Then  $\text{diam}_{(p,q)}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
- $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
- So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
- This does not follow from Dubuc's result.

# Cyclic groups

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
- Then  $\text{diam}_{\{p,q\}}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
- $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
- So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
- This does not follow from Dubuc's result.

# Cyclic groups

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
- Then  $\text{diam}_{\{p,q\}}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
- $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
- So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
- This does not follow from Dubuc's result.

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
  - Then  $\text{diam}_{\{p,q\}}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
  - $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
  - So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
  - This does not follow from Dubuc's result.

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
- Then  $\text{diam}_{\{p,q\}}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
- $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
- So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
- This does not follow from Dubuc's result.

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
- Then  $\text{diam}_{\{p,q\}}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
- $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
- So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
- This does not follow from Dubuc's result.

- One can prove  $m(\mathbb{Z}_n) = \phi(n)$  (Euler's function).
- The irrep comes from the action of  $\mathbb{Z}_n$  on  $\mathbb{Q}(\zeta_n)$  ( $\zeta_n$  a primitive  $n^{\text{th}}$ -root of unity) by multiplication by  $\zeta_n$ .
- Of course,  $\text{diam}_{\{1\}}(\mathbb{Z}_n) = n - 1$ .
- So we achieve the Černý bound iff  $\phi(n) = n - 1$ .
- This occurs iff  $n$  is prime.
- In particular, our method recovers Pin's Theorem, but not Dubuc's Theorem (although we are very close).
- Suppose  $p < q$  are odd primes and  $n = pq$ .
- Then  $\text{diam}_{\{p,q\}}(\mathbb{Z}_n) = q - 1 + p - 1$  ( $\mathbb{Z}_n \cong \mathbb{Z}_q \times \mathbb{Z}_p$ ).
- $\phi(n) = (p - 1)(q - 1) \geq q - 1 + p - 1$ .
- So  $(\mathbb{Z}_{pq}, \{p, q\})$  is a Černý Cayley graph.
- This does not follow from Dubuc's result.



# Direct products of cyclic groups of prime order

- Let  $p$  be a prime.
- To show that  $\mathbb{Z}_p^k$  is a Černý group, it suffices to consider the Cayley graph with respect to a basis  $\Delta$ .
- $\text{diam}_\Delta(\mathbb{Z}_p^k) = k(p-1)$ .
- One can prove  $m(\mathbb{Z}_p^k) = p-1$ .
- Our bound therefore is not strong enough when  $k > 1$ .  
Nonetheless we can prove:

## Theorem (BS)

*The group  $\mathbb{Z}_p^k$  is a Černý group for  $p$  prime and all  $k \geq 1$ .*

# Direct products of cyclic groups of prime order

- Let  $p$  be a prime.
- To show that  $\mathbb{Z}_p^k$  is a Černý group, it suffices to consider the Cayley graph with respect to a basis  $\Delta$ .
- $\text{diam}_\Delta(\mathbb{Z}_p^k) = k(p-1)$ .
- One can prove  $m(\mathbb{Z}_p^k) = p-1$ .
- Our bound therefore is not strong enough when  $k > 1$ .  
Nonetheless we can prove:

## Theorem (BS)

*The group  $\mathbb{Z}_p^k$  is a Černý group for  $p$  prime and all  $k \geq 1$ .*

# Direct products of cyclic groups of prime order

- Let  $p$  be a prime.
- To show that  $\mathbb{Z}_p^k$  is a Černý group, it suffices to consider the Cayley graph with respect to a basis  $\Delta$ .
- $\text{diam}_\Delta(\mathbb{Z}_p^k) = k(p-1)$ .
- One can prove  $m(\mathbb{Z}_p^k) = p-1$ .
- Our bound therefore is not strong enough when  $k > 1$ .  
Nonetheless we can prove:

## Theorem (BS)

*The group  $\mathbb{Z}_p^k$  is a Černý group for  $p$  prime and all  $k \geq 1$ .*

# Direct products of cyclic groups of prime order

- Let  $p$  be a prime.
- To show that  $\mathbb{Z}_p^k$  is a Černý group, it suffices to consider the Cayley graph with respect to a basis  $\Delta$ .
- $\text{diam}_\Delta(\mathbb{Z}_p^k) = k(p - 1)$ .
- One can prove  $m(\mathbb{Z}_p^k) = p - 1$ .
- Our bound therefore is not strong enough when  $k > 1$ .  
Nonetheless we can prove:

## Theorem (BS)

*The group  $\mathbb{Z}_p^k$  is a Černý group for  $p$  prime and all  $k \geq 1$ .*

# Direct products of cyclic groups of prime order

- Let  $p$  be a prime.
- To show that  $\mathbb{Z}_p^k$  is a Černý group, it suffices to consider the Cayley graph with respect to a basis  $\Delta$ .
- $\text{diam}_\Delta(\mathbb{Z}_p^k) = k(p - 1)$ .
- One can prove  $m(\mathbb{Z}_p^k) = p - 1$ .
- Our bound therefore is not strong enough when  $k > 1$ .  
Nonetheless we can prove:

## Theorem (BS)

*The group  $\mathbb{Z}_p^k$  is a Černý group for  $p$  prime and all  $k \geq 1$ .*

# Direct products of cyclic groups of prime order

- Let  $p$  be a prime.
- To show that  $\mathbb{Z}_p^k$  is a Černý group, it suffices to consider the Cayley graph with respect to a basis  $\Delta$ .
- $\text{diam}_\Delta(\mathbb{Z}_p^k) = k(p-1)$ .
- One can prove  $m(\mathbb{Z}_p^k) = p-1$ .
- Our bound therefore is not strong enough when  $k > 1$ .  
Nonetheless we can prove:

## Theorem (BS)

*The group  $\mathbb{Z}_p^k$  is a Černý group for  $p$  prime and all  $k \geq 1$ .*

- Let  $D_n$  be the dihedral group of order  $2n$  (the symmetry group of a regular  $n$ -gon).
- Let  $\Delta$  consist of a reflection and a rotation by  $2\pi/n$ .
- Then  $\text{diam}_\Delta(D_n) \leq \lceil \frac{n+1}{2} \rceil$ .
- One can prove  $m(D_n) = \phi(n)$ .
- If  $n = p^a q^b$  where  $p \leq q$  are odd primes, then one verifies that  $\lceil \frac{n+1}{2} \rceil \leq \phi(n)$  and so we obtain a Černý Cayley graph.

## Theorem (BS)

*Let  $p$  be an odd prime. Then  $D_p$  and  $D_{p^2}$  are Černý groups.*

- Let  $D_n$  be the dihedral group of order  $2n$  (the symmetry group of a regular  $n$ -gon).
- Let  $\Delta$  consist of a reflection and a rotation by  $2\pi/n$ .
- Then  $\text{diam}_\Delta(D_n) \leq \lceil \frac{n+1}{2} \rceil$ .
- One can prove  $m(D_n) = \phi(n)$ .
- If  $n = p^a q^b$  where  $p \leq q$  are odd primes, then one verifies that  $\lceil \frac{n+1}{2} \rceil \leq \phi(n)$  and so we obtain a Černý Cayley graph.

## Theorem (BS)

*Let  $p$  be an odd prime. Then  $D_p$  and  $D_{p^2}$  are Černý groups.*



- Let  $D_n$  be the dihedral group of order  $2n$  (the symmetry group of a regular  $n$ -gon).
- Let  $\Delta$  consist of a reflection and a rotation by  $2\pi/n$ .
- Then  $\text{diam}_\Delta(D_n) \leq \lceil \frac{n+1}{2} \rceil$ .
- One can prove  $m(D_n) = \phi(n)$ .
- If  $n = p^a q^b$  where  $p \leq q$  are odd primes, then one verifies that  $\lceil \frac{n+1}{2} \rceil \leq \phi(n)$  and so we obtain a Černý Cayley graph.

## Theorem (BS)

*Let  $p$  be an odd prime. Then  $D_p$  and  $D_{p^2}$  are Černý groups.*

# Dihedral groups

- Let  $D_n$  be the dihedral group of order  $2n$  (the symmetry group of a regular  $n$ -gon).
- Let  $\Delta$  consist of a reflection and a rotation by  $2\pi/n$ .
- Then  $\text{diam}_\Delta(D_n) \leq \lceil \frac{n+1}{2} \rceil$ .
- One can prove  $m(D_n) = \phi(n)$ .
- If  $n = p^a q^b$  where  $p \leq q$  are odd primes, then one verifies that  $\lceil \frac{n+1}{2} \rceil \leq \phi(n)$  and so we obtain a Černý Cayley graph.

## Theorem (BS)

*Let  $p$  be an odd prime. Then  $D_p$  and  $D_{p^2}$  are Černý groups.*

# Dihedral groups

- Let  $D_n$  be the dihedral group of order  $2n$  (the symmetry group of a regular  $n$ -gon).
- Let  $\Delta$  consist of a reflection and a rotation by  $2\pi/n$ .
- Then  $\text{diam}_\Delta(D_n) \leq \lceil \frac{n+1}{2} \rceil$ .
- One can prove  $m(D_n) = \phi(n)$ .
- If  $n = p^a q^b$  where  $p \leq q$  are odd primes, then one verifies that  $\lceil \frac{n+1}{2} \rceil \leq \phi(n)$  and so we obtain a Černý Cayley graph.

## Theorem (BS)

*Let  $p$  be an odd prime. Then  $D_p$  and  $D_{p^2}$  are Černý groups.*

# Dihedral groups

- Let  $D_n$  be the dihedral group of order  $2n$  (the symmetry group of a regular  $n$ -gon).
- Let  $\Delta$  consist of a reflection and a rotation by  $2\pi/n$ .
- Then  $\text{diam}_\Delta(D_n) \leq \lceil \frac{n+1}{2} \rceil$ .
- One can prove  $m(D_n) = \phi(n)$ .
- If  $n = p^a q^b$  where  $p \leq q$  are odd primes, then one verifies that  $\lceil \frac{n+1}{2} \rceil \leq \phi(n)$  and so we obtain a Černý Cayley graph.

## Theorem (BS)

*Let  $p$  be an odd prime. Then  $D_p$  and  $D_{p^2}$  are Černý groups.*

# Symmetric groups

- It is known that the symmetric group  $S_n$  has  $p_n$  irreducible representations where  $p_n$  is the number of partitions of  $n$ .
- The sum of the squares of the degrees of the irreps of  $S_n$  is  $n!$ .
- Thus  $m(S_n)^2 p_n \geq n!$ , i.e.,  $m(S_n) \geq \sqrt{n!/p_n}$ .
- $p_n \sim \frac{\exp(\pi\sqrt{2n/3})}{4n\sqrt{3}}$  and  $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ .
- Therefore,  $m(S_n)$  grows extremely quickly as a function of  $n$ .
- With Coxeter-Moore generators  $(1\ 2), (2\ 3), \dots, (n-1\ n)$ , the diameter is  $\binom{n}{2}$  [think "Bubble Sort"] and so we obtain a Černý Cayley graph for  $n$  large enough.
- With the generating set  $(1\ 2), (1\ 2\ \dots\ n)$ , the diameter of  $S_n$  is at most  $\binom{n}{2}(n+1)$  and so we again get a Černý Cayley graph for  $n$  large enough.

# Symmetric groups

- It is known that the symmetric group  $S_n$  has  $p_n$  irreducible representations where  $p_n$  is the number of partitions of  $n$ .
- The sum of the squares of the degrees of the irreps of  $S_n$  is  $n!$ .
- Thus  $m(S_n)^2 p_n \geq n!$ , i.e.,  $m(S_n) \geq \sqrt{n!/p_n}$ .
- $p_n \sim \frac{\exp(\pi\sqrt{2n/3})}{4n\sqrt{3}}$  and  $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ .
- Therefore,  $m(S_n)$  grows extremely quickly as a function of  $n$ .
- With Coxeter-Moore generators  $(1\ 2), (2\ 3), \dots, (n-1\ n)$ , the diameter is  $\binom{n}{2}$  [think “Bubble Sort”] and so we obtain a Černý Cayley graph for  $n$  large enough.
- With the generating set  $(1\ 2), (1\ 2\ \dots\ n)$ , the diameter of  $S_n$  is at most  $\binom{n}{2}(n+1)$  and so we again get a Černý Cayley graph for  $n$  large enough.

# Symmetric groups

- It is known that the symmetric group  $S_n$  has  $p_n$  irreducible representations where  $p_n$  is the number of partitions of  $n$ .
- The sum of the squares of the degrees of the irreps of  $S_n$  is  $n!$ .
- Thus  $m(S_n)^2 p_n \geq n!$ , i.e.,  $m(S_n) \geq \sqrt{n!/p_n}$ .
- $p_n \sim \frac{\exp(\pi\sqrt{2n/3})}{4n\sqrt{3}}$  and  $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ .
- Therefore,  $m(S_n)$  grows extremely quickly as a function of  $n$ .
- With Coxeter-Moore generators  $(1\ 2), (2\ 3), \dots, (n-1\ n)$ , the diameter is  $\binom{n}{2}$  [think "Bubble Sort"] and so we obtain a Černý Cayley graph for  $n$  large enough.
- With the generating set  $(1\ 2), (1\ 2\ \dots\ n)$ , the diameter of  $S_n$  is at most  $\binom{n}{2}(n+1)$  and so we again get a Černý Cayley graph for  $n$  large enough.

# Symmetric groups

- It is known that the symmetric group  $S_n$  has  $p_n$  irreducible representations where  $p_n$  is the number of partitions of  $n$ .
- The sum of the squares of the degrees of the irreps of  $S_n$  is  $n!$ .
- Thus  $m(S_n)^2 p_n \geq n!$ , i.e.,  $m(S_n) \geq \sqrt{n!/p_n}$ .
- $p_n \sim \frac{\exp(\pi\sqrt{2n/3})}{4n\sqrt{3}}$  and  $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ .
- Therefore,  $m(S_n)$  grows extremely quickly as a function of  $n$ .
- With Coxeter-Moore generators  $(1\ 2), (2\ 3), \dots, (n-1\ n)$ , the diameter is  $\binom{n}{2}$  [think “Bubble Sort”] and so we obtain a Černý Cayley graph for  $n$  large enough.
- With the generating set  $(1\ 2), (1\ 2\ \dots\ n)$ , the diameter of  $S_n$  is at most  $\binom{n}{2}(n+1)$  and so we again get a Černý Cayley graph for  $n$  large enough.



# Symmetric groups

- It is known that the symmetric group  $S_n$  has  $p_n$  irreducible representations where  $p_n$  is the number of partitions of  $n$ .
- The sum of the squares of the degrees of the irreps of  $S_n$  is  $n!$ .
- Thus  $m(S_n)^2 p_n \geq n!$ , i.e.,  $m(S_n) \geq \sqrt{n!/p_n}$ .
- $p_n \sim \frac{\exp(\pi\sqrt{2n/3})}{4n\sqrt{3}}$  and  $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ .
- Therefore,  $m(S_n)$  grows extremely quickly as a function of  $n$ .
- With Coxeter-Moore generators  $(1\ 2), (2\ 3), \dots, (n-1\ n)$ , the diameter is  $\binom{n}{2}$  [think “Bubble Sort”] and so we obtain a Černý Cayley graph for  $n$  large enough.
- With the generating set  $(1\ 2), (1\ 2\ \dots\ n)$ , the diameter of  $S_n$  is at most  $\binom{n}{2}(n+1)$  and so we again get a Černý Cayley graph for  $n$  large enough.

# Symmetric groups

- It is known that the symmetric group  $S_n$  has  $p_n$  irreducible representations where  $p_n$  is the number of partitions of  $n$ .
- The sum of the squares of the degrees of the irreps of  $S_n$  is  $n!$ .
- Thus  $m(S_n)^2 p_n \geq n!$ , i.e.,  $m(S_n) \geq \sqrt{n!/p_n}$ .
- $p_n \sim \frac{\exp(\pi\sqrt{2n/3})}{4n\sqrt{3}}$  and  $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ .
- Therefore,  $m(S_n)$  grows extremely quickly as a function of  $n$ .
- With Coxeter-Moore generators  $(1\ 2), (2\ 3), \dots, (n-1\ n)$ , the diameter is  $\binom{n}{2}$  [think “Bubble Sort”] and so we obtain a Černý Cayley graph for  $n$  large enough.
- With the generating set  $(1\ 2), (1\ 2\ \dots\ n)$ , the diameter of  $S_n$  is at most  $\binom{n}{2}(n+1)$  and so we again get a Černý Cayley graph for  $n$  large enough.

# Symmetric groups

- It is known that the symmetric group  $S_n$  has  $p_n$  irreducible representations where  $p_n$  is the number of partitions of  $n$ .
- The sum of the squares of the degrees of the irreps of  $S_n$  is  $n!$ .
- Thus  $m(S_n)^2 p_n \geq n!$ , i.e.,  $m(S_n) \geq \sqrt{n!/p_n}$ .
- $p_n \sim \frac{\exp(\pi\sqrt{2n/3})}{4n\sqrt{3}}$  and  $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ .
- Therefore,  $m(S_n)$  grows extremely quickly as a function of  $n$ .
- With Coxeter-Moore generators  $(1\ 2), (2\ 3), \dots, (n-1\ n)$ , the diameter is  $\binom{n}{2}$  [think “Bubble Sort”] and so we obtain a Černý Cayley graph for  $n$  large enough.
- With the generating set  $(1\ 2), (1\ 2\ \dots\ n)$ , the diameter of  $S_n$  is at most  $\binom{n}{2}(n+1)$  and so we again get a Černý Cayley graph for  $n$  large enough.

- Let  $p$  be a prime.
- $SL(2, p) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a, b, c, d \in \mathbb{Z}_p, ad - bc = 1 \right\}$ .
- A standard generating set  $\Delta$  for  $SL(2, p)$  consists of the matrices

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}.$$

- The diameter with this generating set is no more than  $3p - 2$ .
- Estimating  $m(SL(2, p))$  is a bit more complicated.

- Let  $p$  be a prime.
- $SL(2, p) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a, b, c, d \in \mathbb{Z}_p, ad - bc = 1 \right\}$ .
- A standard generating set  $\Delta$  for  $SL(2, p)$  consists of the matrices

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}.$$

- The diameter with this generating set is no more than  $3p - 2$ .
- Estimating  $m(SL(2, p))$  is a bit more complicated.

- Let  $p$  be a prime.
- $SL(2, p) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a, b, c, d \in \mathbb{Z}_p, ad - bc = 1 \right\}$ .
- A standard generating set  $\Delta$  for  $SL(2, p)$  consists of the matrices

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}.$$

- The diameter with this generating set is no more than  $3p - 2$ .
- Estimating  $m(SL(2, p))$  is a bit more complicated.

- Let  $p$  be a prime.
- $SL(2, p) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a, b, c, d \in \mathbb{Z}_p, ad - bc = 1 \right\}$ .
- A standard generating set  $\Delta$  for  $SL(2, p)$  consists of the matrices

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}.$$

- The diameter with this generating set is no more than  $3p - 2$ .
- Estimating  $m(SL(2, p))$  is a bit more complicated.

- Let  $p$  be a prime.
- $SL(2, p) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a, b, c, d \in \mathbb{Z}_p, ad - bc = 1 \right\}$ .
- A standard generating set  $\Delta$  for  $SL(2, p)$  consists of the matrices

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}.$$

- The diameter with this generating set is no more than  $3p - 2$ .
- Estimating  $m(SL(2, p))$  is a bit more complicated.



# Special linear groups II

- In the group theory literature there is much more detailed information about representations over  $\mathbb{C}$  than over  $\mathbb{Q}$ .
- Schur index theory allows one to use Galois theory in order to understand irreps over  $\mathbb{Q}$  in terms of irreps over  $\mathbb{C}$ .
- Via these methods, we computed
$$m(SL(2, p)) \geq \max \left\{ (p+1)^{\frac{\phi(p-1)}{2}}, (p-1)^{\frac{\phi(p+1)}{2}} \right\}.$$
- The diameter of the Cayley graph of  $SL(2, p)$  with our generators was at most  $3p - 2$ .

## Theorem (BS)

*Let  $p \geq 17$  be a prime. Then the Cayley graph of  $SL(2, p)$  with respect to the generators  $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  is a Černý Cayley graph.*

# Special linear groups II

- In the group theory literature there is much more detailed information about representations over  $\mathbb{C}$  than over  $\mathbb{Q}$ .
- Schur index theory allows one to use Galois theory in order to understand irreps over  $\mathbb{Q}$  in terms of irreps over  $\mathbb{C}$ .
- Via these methods, we computed
$$m(SL(2, p)) \geq \max \left\{ (p+1)^{\frac{\phi(p-1)}{2}}, (p-1)^{\frac{\phi(p+1)}{2}} \right\}.$$
- The diameter of the Cayley graph of  $SL(2, p)$  with our generators was at most  $3p - 2$ .

## Theorem (BS)

*Let  $p \geq 17$  be a prime. Then the Cayley graph of  $SL(2, p)$  with respect to the generators  $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  is a Černý Cayley graph.*

# Special linear groups II

- In the group theory literature there is much more detailed information about representations over  $\mathbb{C}$  than over  $\mathbb{Q}$ .
- Schur index theory allows one to use Galois theory in order to understand irreps over  $\mathbb{Q}$  in terms of irreps over  $\mathbb{C}$ .
- Via these methods, we computed
$$m(SL(2, p)) \geq \max \left\{ (p+1)^{\frac{\phi(p-1)}{2}}, (p-1)^{\frac{\phi(p+1)}{2}} \right\}.$$
- The diameter of the Cayley graph of  $SL(2, p)$  with our generators was at most  $3p - 2$ .

## Theorem (BS)

*Let  $p \geq 17$  be a prime. Then the Cayley graph of  $SL(2, p)$  with respect to the generators  $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  is a Černý Cayley graph.*

# Special linear groups II

- In the group theory literature there is much more detailed information about representations over  $\mathbb{C}$  than over  $\mathbb{Q}$ .
- Schur index theory allows one to use Galois theory in order to understand irreps over  $\mathbb{Q}$  in terms of irreps over  $\mathbb{C}$ .
- Via these methods, we computed
$$m(SL(2, p)) \geq \max \left\{ (p+1)^{\frac{\phi(p-1)}{2}}, (p-1)^{\frac{\phi(p+1)}{2}} \right\}.$$
- The diameter of the Cayley graph of  $SL(2, p)$  with our generators was at most  $3p - 2$ .

## Theorem (BS)

*Let  $p \geq 17$  be a prime. Then the Cayley graph of  $SL(2, p)$  with respect to the generators  $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  is a Černý Cayley graph.*

# Special linear groups II

- In the group theory literature there is much more detailed information about representations over  $\mathbb{C}$  than over  $\mathbb{Q}$ .
- Schur index theory allows one to use Galois theory in order to understand irreps over  $\mathbb{Q}$  in terms of irreps over  $\mathbb{C}$ .
- Via these methods, we computed
$$m(SL(2, p)) \geq \max \left\{ (p+1)^{\frac{\phi(p-1)}{2}}, (p-1)^{\frac{\phi(p+1)}{2}} \right\}.$$
- The diameter of the Cayley graph of  $SL(2, p)$  with our generators was at most  $3p - 2$ .

## Theorem (BS)

*Let  $p \geq 17$  be a prime. Then the Cayley graph of  $SL(2, p)$  with respect to the generators  $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$  is a Černý Cayley graph.*

- A number of questions remain open.
- Is every cyclic group a Černý group?
- Is every abelian group a Černý group?
- Is every dihedral group a Černý group?
- Is every group a Černý group?

- A number of questions remain open.
- Is every cyclic group a Černý group?
- Is every abelian group a Černý group?
- Is every dihedral group a Černý group?
- Is every group a Černý group?

- A number of questions remain open.
- Is every cyclic group a Černý group?
- Is every abelian group a Černý group?
- Is every dihedral group a Černý group?
- Is every group a Černý group?



- A number of questions remain open.
- Is every cyclic group a Černý group?
- Is every abelian group a Černý group?
- Is every dihedral group a Černý group?
- Is every group a Černý group?

# Open questions

- A number of questions remain open.
- Is every cyclic group a Černý group?
- Is every abelian group a Černý group?
- Is every dihedral group a Černý group?
- Is every group a Černý group?

Vielen Dank für Ihre Aufmerksamkeit!