

Quivers of monoids with basic algebras

Stuart Margolis¹ Benjamin Steinberg²

¹Bar-Ilan University

²Carleton University

bsteinbg@math.carleton.ca

<http://www.mathstat.carleton.ca/~bsteinbg>

February 16, 2011

Outline

- 1 Motivation
 - History
- 2 Finite dimensional algebras
 - Quivers of Algebras
- 3 History
 - Left regular bands
 - Left regular bands of groups
 - \mathcal{R} -trivial monoids
- 4 Rectangular Monoids
 - Monoids with Basic Algebras
 - Quivers of rectangular monoids

Group Representation Theory and its Applications

- Group representation theory originated in the late 1800s with the work of Frobenius.
- It took some time before its usefulness was seen.
- The preface of the first edition of Burnside's book on finite groups asserts:

"it would be difficult to find a result that could be most directly obtained by the consideration of groups of linear transformations."

- Nowadays, group representation theory is routinely applied to such diverse areas as physics, probability, data analysis, graph theory and signal processing.

Group Representation Theory and its Applications

- Group representation theory originated in the late 1800s with the work of Frobenius.
- It took some time before its usefulness was seen.
- The preface of the first edition of Burnside's book on finite groups asserts:

"it would be difficult to find a result that could be most directly obtained by the consideration of groups of linear transformations."

- Nowadays, group representation theory is routinely applied to such diverse areas as physics, probability, data analysis, graph theory and signal processing.

Group Representation Theory and its Applications

- Group representation theory originated in the late 1800s with the work of Frobenius.
- It took some time before its usefulness was seen.
- The preface of the first edition of Burnside's book on finite groups asserts:

"it would be difficult to find a result that could be most directly obtained by the consideration of groups of linear transformations."

- Nowadays, group representation theory is routinely applied to such diverse areas as physics, probability, data analysis, graph theory and signal processing.

Group Representation Theory and its Applications

- Group representation theory originated in the late 1800s with the work of Frobenius.
- It took some time before its usefulness was seen.
- The preface of the first edition of Burnside's book on finite groups asserts:

"it would be difficult to find a result that could be most directly obtained by the consideration of groups of linear transformations."

- Nowadays, group representation theory is routinely applied to such diverse areas as physics, probability, data analysis, graph theory and signal processing.

Group Representation Theory and its Applications

- Group representation theory originated in the late 1800s with the work of Frobenius.
- It took some time before its usefulness was seen.
- The preface of the first edition of Burnside's book on finite groups asserts:

"it would be difficult to find a result that could be most directly obtained by the consideration of groups of linear transformations."

- Nowadays, group representation theory is routinely applied to such diverse areas as physics, probability, data analysis, graph theory and signal processing.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as real and complex hyperplane arrangements, matroids, oriented matroids and interval greedoids have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be *more* easily analyzed as semigroup walks than as group walks.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as real and complex hyperplane arrangements, matroids, oriented matroids and interval greedoids have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be *more* easily analyzed as semigroup walks than as group walks.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as real and complex hyperplane arrangements, matroids, oriented matroids and interval greedoids have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be *more* easily analyzed as semigroup walks than as group walks.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as real and complex hyperplane arrangements, matroids, oriented matroids and interval greedoids have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be *more* easily analyzed as semigroup walks than as group walks.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as **real and complex hyperplane arrangements**, **matroids**, **oriented matroids** and **interval greedoids** have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be *more easily* analyzed as semigroup walks than as group walks.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as **real and complex hyperplane arrangements**, **matroids**, **oriented matroids** and **interval greedoids** have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be *more easily* analyzed as semigroup walks than as group walks.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as **real and complex hyperplane arrangements**, **matroids**, **oriented matroids** and **interval greedoids** have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be *more* easily analyzed as semigroup walks than as group walks.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as **real and complex hyperplane arrangements**, **matroids**, **oriented matroids** and **interval greedoids** have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be *more* easily analyzed as semigroup walks than as group walks.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as **real and complex hyperplane arrangements**, **matroids**, **oriented matroids** and **interval greedoids** have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be *more* easily analyzed as semigroup walks than as group walks.

Semigroup Representation Theory and Markov Chains

- Clifford initiated semigroup representation theory in the 1940s.
- The first applications appeared in 1998 in the work of Bidigare, Hanlon and Rockmore (Duke J. Math).
- This was followed by work of Brown and Diaconis, Björner and others.
- Many combinatorial objects, such as **real and complex hyperplane arrangements**, **matroids**, **oriented matroids** and **interval greedoids** have the structure of a semigroup.
- Semigroup representation theory can be used to analyze random walks on these combinatorial objects.
- Important examples, like card shuffling, can be **more** easily analyzed as semigroup walks than as group walks.

Further Applications

- Bidigare also established a strong connection between hyperplane face semigroups and **Solomon's descent algebra**.
- The descent algebra is at the heart of the non-commutative character theory of the symmetric group.
- Saliola used the semigroup approach in his computation of the quiver of the descent algebra in types A and B .
- Semigroups play an important role in the study of the 0-Hecke algebra of a Coxeter group and related algebras (Bergeron, Denton, Hivert, Thiéry, Schilling and others).
- Malandro and Rockmore have used fast Fourier transforms for semigroup algebras to study partially ranked data (Trans. AMS).

Further Applications

- Bidigare also established a strong connection between hyperplane face semigroups and **Solomon's descent algebra**.
- The descent algebra is at the heart of the non-commutative character theory of the symmetric group.
- Saliola used the semigroup approach in his computation of the quiver of the descent algebra in types A and B .
- Semigroups play an important role in the study of the 0-Hecke algebra of a Coxeter group and related algebras (Bergeron, Denton, Hivert, Thiéry, Schilling and others).
- Malandro and Rockmore have used fast Fourier transforms for semigroup algebras to study partially ranked data (Trans. AMS).

Further Applications

- Bidigare also established a strong connection between hyperplane face semigroups and **Solomon's descent algebra**.
- The descent algebra is at the heart of the non-commutative character theory of the symmetric group.
- Saliola used the semigroup approach in his computation of the quiver of the descent algebra in types A and B .
- Semigroups play an important role in the study of the 0-Hecke algebra of a Coxeter group and related algebras (Bergeron, Denton, Hivert, Thiéry, Schilling and others).
- Malandro and Rockmore have used fast Fourier transforms for semigroup algebras to study partially ranked data (Trans. AMS).

Further Applications

- Bidigare also established a strong connection between hyperplane face semigroups and **Solomon's descent algebra**.
- The descent algebra is at the heart of the non-commutative character theory of the symmetric group.
- Saliola used the semigroup approach in his computation of the quiver of the descent algebra in types A and B .
- Semigroups play an important role in the study of the 0-Hecke algebra of a Coxeter group and related algebras (Bergeron, Denton, Hivert, Thiéry, Schilling and others).
- Malandro and Rockmore have used fast Fourier transforms for semigroup algebras to study partially ranked data (Trans. AMS).

Further Applications

- Bidigare also established a strong connection between hyperplane face semigroups and **Solomon's descent algebra**.
- The descent algebra is at the heart of the non-commutative character theory of the symmetric group.
- Saliola used the semigroup approach in his computation of the quiver of the descent algebra in types A and B .
- Semigroups play an important role in the study of the 0-Hecke algebra of a Coxeter group and related algebras (Bergeron, Denton, Hivert, Thiéry, Schilling and others).
- Malandro and Rockmore have used fast Fourier transforms for semigroup algebras to study partially ranked data (Trans. AMS).

Basic Algebras

- Fix an algebraically closed field k of characteristic 0.
- All algebras are assumed finite dimensional over k and unital.
- A is **basic** if $A/\text{rad}(A) \cong k^n$.
- In other words, all irreducible representations of A are one-dimensional.
- Equivalently, A is an algebra of upper triangular matrices.
- Each algebra is Morita equivalent to a unique basic algebra.
- So for much of representation theory, it is enough to consider basic algebras.
- Caveat: This viewpoint says nothing about group algebras and so one must analyze one's goals before passing to Morita equivalence.

Basic Algebras

- Fix an algebraically closed field k of characteristic 0.
- All algebras are assumed finite dimensional over k and unital.
- A is **basic** if $A/\text{rad}(A) \cong k^n$.
- In other words, all irreducible representations of A are one-dimensional.
- Equivalently, A is an algebra of upper triangular matrices.
- Each algebra is Morita equivalent to a unique basic algebra.
- So for much of representation theory, it is enough to consider basic algebras.
- Caveat: This viewpoint says nothing about group algebras and so one must analyze one's goals before passing to Morita equivalence.

Basic Algebras

- Fix an algebraically closed field k of characteristic 0.
- All algebras are assumed finite dimensional over k and unital.
- A is **basic** if $A/\text{rad}(A) \cong k^n$.
- In other words, all irreducible representations of A are one-dimensional.
- Equivalently, A is an algebra of upper triangular matrices.
- Each algebra is Morita equivalent to a unique basic algebra.
- So for much of representation theory, it is enough to consider basic algebras.
- Caveat: This viewpoint says nothing about group algebras and so one must analyze one's goals before passing to Morita equivalence.

Basic Algebras

- Fix an algebraically closed field k of characteristic 0.
- All algebras are assumed finite dimensional over k and unital.
- A is **basic** if $A/\text{rad}(A) \cong k^n$.
- In other words, all irreducible representations of A are one-dimensional.
- Equivalently, A is an algebra of upper triangular matrices.
- Each algebra is Morita equivalent to a unique basic algebra.
- So for much of representation theory, it is enough to consider basic algebras.
- Caveat: This viewpoint says nothing about group algebras and so one must analyze one's goals before passing to Morita equivalence.

Basic Algebras

- Fix an algebraically closed field k of characteristic 0.
- All algebras are assumed finite dimensional over k and unital.
- A is **basic** if $A/\text{rad}(A) \cong k^n$.
- In other words, all irreducible representations of A are one-dimensional.
- Equivalently, A is an algebra of upper triangular matrices.
- Each algebra is Morita equivalent to a unique basic algebra.
- So for much of representation theory, it is enough to consider basic algebras.
- **Caveat:** This viewpoint says nothing about group algebras and so one must analyze one's goals before passing to Morita equivalence.

Basic Algebras

- Fix an algebraically closed field k of characteristic 0.
- All algebras are assumed finite dimensional over k and unital.
- A is **basic** if $A/\text{rad}(A) \cong k^n$.
- In other words, all irreducible representations of A are one-dimensional.
- Equivalently, A is an algebra of upper triangular matrices.
- Each algebra is Morita equivalent to a unique basic algebra.
- So for much of representation theory, it is enough to consider basic algebras.
- **Caveat:** This viewpoint says nothing about group algebras and so one must analyze one's goals before passing to Morita equivalence.

Basic Algebras

- Fix an algebraically closed field k of characteristic 0.
- All algebras are assumed finite dimensional over k and unital.
- A is **basic** if $A/\text{rad}(A) \cong k^n$.
- In other words, all irreducible representations of A are one-dimensional.
- Equivalently, A is an algebra of upper triangular matrices.
- Each algebra is Morita equivalent to a unique basic algebra.
- So for much of representation theory, it is enough to consider basic algebras.
- **Caveat:** This viewpoint says nothing about group algebras and so one must analyze one's goals before passing to Morita equivalence.

Basic Algebras

- Fix an algebraically closed field k of characteristic 0.
- All algebras are assumed finite dimensional over k and unital.
- A is **basic** if $A/\text{rad}(A) \cong k^n$.
- In other words, all irreducible representations of A are one-dimensional.
- Equivalently, A is an algebra of upper triangular matrices.
- Each algebra is Morita equivalent to a unique basic algebra.
- So for much of representation theory, it is enough to consider basic algebras.
- **Caveat:** This viewpoint says nothing about group algebras and so one must analyze one's goals before passing to Morita equivalence.

Quivers

- A **quiver** Q is a directed graph.
- The path algebra kQ has basis all paths in Q and multiplication extending concatenation.

Example

The path algebra of A_n is the algebra of $n \times n$ upper triangular matrices.



- Gabriel proved every basic algebra is a quotient of a path algebra by an admissible ideal.

Quivers

- A **quiver** Q is a directed graph.
- The path algebra kQ has basis all paths in Q and multiplication extending concatenation.

Example

The path algebra of A_n is the algebra of $n \times n$ upper triangular matrices.




- Gabriel proved every basic algebra is a quotient of a path algebra by an admissible ideal.

Quivers

- A **quiver** Q is a directed graph.
- The path algebra kQ has basis all paths in Q and multiplication extending concatenation.

Example

The path algebra of A_n is the algebra of $n \times n$ upper triangular matrices.



- Gabriel proved every basic algebra is a quotient of a path algebra by an admissible ideal.

Quivers

- A **quiver** Q is a directed graph.
- The path algebra kQ has basis all paths in Q and multiplication extending concatenation.

Example

The path algebra of A_n is the algebra of $n \times n$ upper triangular matrices.



- Gabriel proved every basic algebra is a quotient of a path algebra by an admissible ideal.

The Quiver of an Algebra

- The set of isoclasses of simple left A -modules is denoted $\text{Irr}(A)$.
- The **quiver** $Q(A)$ of A has vertex set $\text{Irr}(A)$.
- The number of edges from S to S' is $\dim \text{Ext}_A^1(S, S')$.
- The basic algebra of A is a quotient $kQ(A)$ by an admissible ideal.
- $Q(A)$ encodes the category of projective indecomposable A -modules.
- When A is basic, then $Q(A)$ encodes the two-dimensional representation theory of A .
- In this talk we are interested in the representation theory of monoids with basic algebras.

The Quiver of an Algebra

- The set of isoclasses of simple left A -modules is denoted $\text{Irr}(A)$.
- The **quiver** $Q(A)$ of A has vertex set $\text{Irr}(A)$.
- The number of edges from S to S' is $\dim \text{Ext}_A^1(S, S')$.
- The basic algebra of A is a quotient $kQ(A)$ by an admissible ideal.
- $Q(A)$ encodes the category of projective indecomposable A -modules.
- When A is basic, then $Q(A)$ encodes the two-dimensional representation theory of A .
- In this talk we are interested in the representation theory of monoids with basic algebras.

The Quiver of an Algebra

- The set of isoclasses of simple left A -modules is denoted $\text{Irr}(A)$.
- The **quiver** $Q(A)$ of A has vertex set $\text{Irr}(A)$.
- The number of edges from S to S' is $\dim \text{Ext}_A^1(S, S')$.
- The basic algebra of A is a quotient $kQ(A)$ by an admissible ideal.
- $Q(A)$ encodes the category of projective indecomposable A -modules.
- When A is basic, then $Q(A)$ encodes the two-dimensional representation theory of A .
- In this talk we are interested in the representation theory of monoids with basic algebras.

The Quiver of an Algebra

- The set of isoclasses of simple left A -modules is denoted $\text{Irr}(A)$.
- The **quiver** $Q(A)$ of A has vertex set $\text{Irr}(A)$.
- The number of edges from S to S' is $\dim \text{Ext}_A^1(S, S')$.
- The basic algebra of A is a quotient $kQ(A)$ by an admissible ideal.
- $Q(A)$ encodes the category of projective indecomposable A -modules.
- When A is basic, then $Q(A)$ encodes the two-dimensional representation theory of A .
- In this talk we are interested in the representation theory of monoids with basic algebras.

The Quiver of an Algebra

- The set of isoclasses of simple left A -modules is denoted $\text{Irr}(A)$.
- The **quiver** $Q(A)$ of A has vertex set $\text{Irr}(A)$.
- The number of edges from S to S' is $\dim \text{Ext}_A^1(S, S')$.
- The basic algebra of A is a quotient $kQ(A)$ by an admissible ideal.
- $Q(A)$ encodes the category of projective indecomposable A -modules.
- When A is basic, then $Q(A)$ encodes the two-dimensional representation theory of A .
- In this talk we are interested in the representation theory of monoids with basic algebras.

The Quiver of an Algebra


- The set of isoclasses of simple left A -modules is denoted $\text{Irr}(A)$.
- The **quiver** $Q(A)$ of A has vertex set $\text{Irr}(A)$.
- The number of edges from S to S' is $\dim \text{Ext}_A^1(S, S')$.
- The basic algebra of A is a quotient $kQ(A)$ by an admissible ideal.
- $Q(A)$ encodes the category of projective indecomposable A -modules.
- When A is basic, then $Q(A)$ encodes the two-dimensional representation theory of A .
- In this talk we are interested in the representation theory of monoids with basic algebras.

The Quiver of an Algebra

- The set of isoclasses of simple left A -modules is denoted $\text{Irr}(A)$.
- The **quiver** $Q(A)$ of A has vertex set $\text{Irr}(A)$.
- The number of edges from S to S' is $\dim \text{Ext}_A^1(S, S')$.
- The basic algebra of A is a quotient $kQ(A)$ by an admissible ideal.
- $Q(A)$ encodes the category of projective indecomposable A -modules.
- When A is basic, then $Q(A)$ encodes the two-dimensional representation theory of A .
- In this talk we are interested in the representation theory of monoids with basic algebras.

Hyperplane Face Semigroups


- Consider a central hyperplane arrangement \mathcal{H} , e.g.,



- The hyperplane arrangement carves \mathbb{R}^n into chambers, faces, etc.
- The Tits projections gives the set $F(\mathcal{H})$ of faces the structure of a semigroup.

Hyperplane Face Semigroups


- Consider a central hyperplane arrangement \mathcal{H} , e.g.,



- The hyperplane arrangement carves \mathbb{R}^n into chambers, faces, etc.
- The Tits projections gives the set $F(\mathcal{H})$ of faces the structure of a semigroup.

Hyperplane Face Semigroups

- Consider a central hyperplane arrangement \mathcal{H} , e.g.,



- The hyperplane arrangement carves \mathbb{R}^n into chambers, faces, etc.
- The Tits projections gives the set $F(\mathcal{H})$ of faces the structure of a semigroup.

Left Regular Bands

- $F(\mathcal{H})$ satisfies the identities $x^2 = x$ and $xyx = xy$.
- Semigroups satisfying these identities are called **left regular bands** (LRBs).
- Semigroups satisfying just $x^2 = x$ are called **bands**.
- All bands have basic algebras.
- Brown developed a theory of random walks on LRBs.
- These walks include card shuffling, random walks on bases of matroids and the Tsetlin library.
- Using the structure of the semigroup, he showed all LRB random walks are diagonalizable.
- This gave a uniform explanation for diagonalizability of a number of classical Markov chains.

Left Regular Bands

- $F(\mathcal{H})$ satisfies the identities $x^2 = x$ and $xyx = xy$.
- Semigroups satisfying these identities are called **left regular bands** (LRBs).
- Semigroups satisfying just $x^2 = x$ are called **bands**.
- All bands have basic algebras.
- Brown developed a theory of random walks on LRBs.
- These walks include card shuffling, random walks on bases of matroids and the Tsetlin library.
- Using the structure of the semigroup, he showed all LRB random walks are diagonalizable.
- This gave a uniform explanation for diagonalizability of a number of classical Markov chains.

Left Regular Bands

- $F(\mathcal{H})$ satisfies the identities $x^2 = x$ and $xyx = xy$.
- Semigroups satisfying these identities are called **left regular bands** (LRBs).
- Semigroups satisfying just $x^2 = x$ are called **bands**.
- All bands have basic algebras.
- Brown developed a theory of random walks on LRBs.
- These walks include card shuffling, random walks on bases of matroids and the Tsetlin library.
- Using the structure of the semigroup, he showed all LRB random walks are diagonalizable.
- This gave a uniform explanation for diagonalizability of a number of classical Markov chains.

Left Regular Bands

- $F(\mathcal{H})$ satisfies the identities $x^2 = x$ and $xyx = xy$.
- Semigroups satisfying these identities are called **left regular bands** (LRBs).
- Semigroups satisfying just $x^2 = x$ are called **bands**.
- All bands have basic algebras.
- Brown developed a theory of random walks on LRBs.
- These walks include card shuffling, random walks on bases of matroids and the Tsetlin library.
- Using the structure of the semigroup, he showed all LRB random walks are diagonalizable.
- This gave a uniform explanation for diagonalizability of a number of classical Markov chains.

Left Regular Bands

- $F(\mathcal{H})$ satisfies the identities $x^2 = x$ and $xyx = xy$.
- Semigroups satisfying these identities are called **left regular bands** (LRBs).
- Semigroups satisfying just $x^2 = x$ are called **bands**.
- All bands have basic algebras.
- Brown developed a theory of random walks on LRBs.
- These walks include card shuffling, random walks on bases of matroids and the Tsetlin library.
- Using the structure of the semigroup, he showed all LRB random walks are diagonalizable.
- This gave a uniform explanation for diagonalizability of a number of classical Markov chains.

Left Regular Bands

- $F(\mathcal{H})$ satisfies the identities $x^2 = x$ and $xyx = xy$.
- Semigroups satisfying these identities are called **left regular bands** (LRBs).
- Semigroups satisfying just $x^2 = x$ are called **bands**.
- All bands have basic algebras.
- Brown developed a theory of random walks on LRBs.
- These walks include card shuffling, random walks on bases of matroids and the Tsetlin library.
- Using the structure of the semigroup, he showed all LRB random walks are diagonalizable.
- This gave a uniform explanation for diagonalizability of a number of classical Markov chains.

Left Regular Bands

- $F(\mathcal{H})$ satisfies the identities $x^2 = x$ and $xyx = xy$.
- Semigroups satisfying these identities are called **left regular bands** (LRBs).
- Semigroups satisfying just $x^2 = x$ are called **bands**.
- All bands have basic algebras.
- Brown developed a theory of random walks on LRBs.
- These walks include card shuffling, random walks on bases of matroids and the Tsetlin library.
- Using the structure of the semigroup, he showed all LRB random walks are diagonalizable.
- This gave a uniform explanation for diagonalizability of a number of classical Markov chains.

Left Regular Bands

- $F(\mathcal{H})$ satisfies the identities $x^2 = x$ and $xyx = xy$.
- Semigroups satisfying these identities are called **left regular bands** (LRBs).
- Semigroups satisfying just $x^2 = x$ are called **bands**.
- All bands have basic algebras.
- Brown developed a theory of random walks on LRBs.
- These walks include card shuffling, random walks on bases of matroids and the Tsetlin library.
- Using the structure of the semigroup, he showed all LRB random walks are diagonalizable.
- This gave a uniform explanation for diagonalizability of a number of classical Markov chains.

Solomon's Descent Algebra

- If W is a finite Coxeter group, there is an associated reflection arrangement \mathcal{H}_W .
- W acts by automorphisms on $F(\mathcal{H}_W)$ and hence on $kF(\mathcal{H}_W)$.
- Bidigare proved that the algebra $kF(\mathcal{H}_W)^W$ of W -invariants is isomorphic to Solomon's descent algebra.
- The descent algebra is a subalgebra of kW .
- It is a basic algebra with semisimple quotient a subalgebra of the character ring of W .
- In type A , the semisimple quotient is the character ring.
- The descent algebra is at the heart of the non-commutative character theory of Coxeter groups.

Solomon's Descent Algebra

- If W is a finite Coxeter group, there is an associated reflection arrangement \mathcal{H}_W .
- W acts by automorphisms on $F(\mathcal{H}_W)$ and hence on $kF(\mathcal{H}_W)$.
- Bidigare proved that the algebra $kF(\mathcal{H}_W)^W$ of W -invariants is isomorphic to Solomon's descent algebra.
- The descent algebra is a subalgebra of kW .
- It is a basic algebra with semisimple quotient a subalgebra of the character ring of W .
- In type A , the semisimple quotient is the character ring.
- The descent algebra is at the heart of the non-commutative character theory of Coxeter groups.

Solomon's Descent Algebra

- If W is a finite Coxeter group, there is an associated reflection arrangement \mathcal{H}_W .
- W acts by automorphisms on $F(\mathcal{H}_W)$ and hence on $kF(\mathcal{H}_W)$.
- Bidigare proved that the algebra $kF(\mathcal{H}_W)^W$ of W -invariants is isomorphic to Solomon's descent algebra.
- The descent algebra is a subalgebra of kW .
- It is a basic algebra with semisimple quotient a subalgebra of the character ring of W .
- In type A , the semisimple quotient is the character ring.
- The descent algebra is at the heart of the non-commutative character theory of Coxeter groups.

Solomon's Descent Algebra

- If W is a finite Coxeter group, there is an associated reflection arrangement \mathcal{H}_W .
- W acts by automorphisms on $F(\mathcal{H}_W)$ and hence on $kF(\mathcal{H}_W)$.
- Bidigare proved that the algebra $kF(\mathcal{H}_W)^W$ of W -invariants is isomorphic to Solomon's descent algebra.
- The descent algebra is a subalgebra of kW .
- It is a basic algebra with semisimple quotient a subalgebra of the character ring of W .
- In type A , the semisimple quotient is the character ring.
- The descent algebra is at the heart of the non-commutative character theory of Coxeter groups.

Solomon's Descent Algebra

- If W is a finite Coxeter group, there is an associated reflection arrangement \mathcal{H}_W .
- W acts by automorphisms on $F(\mathcal{H}_W)$ and hence on $kF(\mathcal{H}_W)$.
- Bidigare proved that the algebra $kF(\mathcal{H}_W)^W$ of W -invariants is isomorphic to Solomon's descent algebra.
- The descent algebra is a subalgebra of kW .
- It is a basic algebra with semisimple quotient a subalgebra of the character ring of W .
- In type A , the semisimple quotient is the character ring.
- The descent algebra is at the heart of the non-commutative character theory of Coxeter groups.

Solomon's Descent Algebra

- If W is a finite Coxeter group, there is an associated reflection arrangement \mathcal{H}_W .
- W acts by automorphisms on $F(\mathcal{H}_W)$ and hence on $kF(\mathcal{H}_W)$.
- Bidigare proved that the algebra $kF(\mathcal{H}_W)^W$ of W -invariants is isomorphic to Solomon's descent algebra.
- The descent algebra is a subalgebra of kW .
- It is a basic algebra with semisimple quotient a subalgebra of the character ring of W .
- In type A , the semisimple quotient is the character ring.
- The descent algebra is at the heart of the non-commutative character theory of Coxeter groups.

Solomon's Descent Algebra

- If W is a finite Coxeter group, there is an associated reflection arrangement \mathcal{H}_W .
- W acts by automorphisms on $F(\mathcal{H}_W)$ and hence on $kF(\mathcal{H}_W)$.
- Bidigare proved that the algebra $kF(\mathcal{H}_W)^W$ of W -invariants is isomorphic to Solomon's descent algebra.
- The descent algebra is a subalgebra of kW .
- It is a basic algebra with semisimple quotient a subalgebra of the character ring of W .
- In type A , the semisimple quotient is the character ring.
- The descent algebra is at the heart of the non-commutative character theory of Coxeter groups.

The Representation Theory of LRBs

- Saliola developed in his thesis the representation theory of LRBs.
- He computed the quiver, the Cartan invariants and a complete set of primitive idempotents.
- He also gave a semigroup theoretic model of the projective indecomposables.
- That is, he gave a basis B for the projective indecomposables such that $B \cup \{0\}$ is invariant under the action of the semigroup.
- He computed primitive idempotents and the quiver for the descent algebra in type A and B based on his LRB results.

The Representation Theory of LRBs

- Saliola developed in his thesis the representation theory of LRBs.
- He computed the quiver, the Cartan invariants and a complete set of primitive idempotents.
- He also gave a semigroup theoretic model of the projective indecomposables.
- That is, he gave a basis B for the projective indecomposables such that $B \cup \{0\}$ is invariant under the action of the semigroup.
- He computed primitive idempotents and the quiver for the descent algebra in type A and B based on his LRB results.

The Representation Theory of LRBs

- Saliola developed in his thesis the representation theory of LRBs.
- He computed the quiver, the Cartan invariants and a complete set of primitive idempotents.
- He also gave a semigroup theoretic model of the projective indecomposables.
- That is, he gave a basis B for the projective indecomposables such that $B \cup \{0\}$ is invariant under the action of the semigroup.
- He computed primitive idempotents and the quiver for the descent algebra in type A and B based on his LRB results.

The Representation Theory of LRBs

- Saliola developed in his thesis the representation theory of LRBs.
- He computed the quiver, the Cartan invariants and a complete set of primitive idempotents.
- He also gave a semigroup theoretic model of the projective indecomposables.
- That is, he gave a basis B for the projective indecomposables such that $B \cup \{0\}$ is invariant under the action of the semigroup.
- He computed primitive idempotents and the quiver for the descent algebra in type A and B based on his LRB results.

The Representation Theory of LRBs

- Saliola developed in his thesis the representation theory of LRBs.
- He computed the quiver, the Cartan invariants and a complete set of primitive idempotents.
- He also gave a semigroup theoretic model of the projective indecomposables.
- That is, he gave a basis B for the projective indecomposables such that $B \cup \{0\}$ is invariant under the action of the semigroup.
- He computed primitive idempotents and the quiver for the descent algebra in type A and B based on his LRB results.

Hereditary Algebras and Left Regular Bands

- An algebra A is **hereditary** if each left ideal of A is projective.
- Equivalently, each submodule of a projective A -module is projective.
- Gabriel proved that a basic algebra is hereditary iff it is the path algebra of an acyclic quiver.
- Recently, Margolis, Saliola and I proved that if M is an LRB such that the Hasse diagram of the poset of principal right ideals aM of M is a tree, then kM is hereditary.
- This includes the algebras of nearly all LRBs considered by Brown except the hyperplane face semigroups!
- We also showed that every basic hereditary algebra is an LRB algebra.

Hereditary Algebras and Left Regular Bands

- An algebra A is **hereditary** if each left ideal of A is projective.
- Equivalently, each submodule of a projective A -module is projective.
- Gabriel proved that a basic algebra is hereditary iff it is the path algebra of an acyclic quiver.
- Recently, Margolis, Saliola and I proved that if M is an LRB such that the Hasse diagram of the poset of principal right ideals aM of M is a tree, then kM is hereditary.
- This includes the algebras of nearly all LRBs considered by Brown except the hyperplane face semigroups!
- We also showed that every basic hereditary algebra is an LRB algebra.

Hereditary Algebras and Left Regular Bands

- An algebra A is **hereditary** if each left ideal of A is projective.
- Equivalently, each submodule of a projective A -module is projective.
- Gabriel proved that a basic algebra is hereditary iff it is the path algebra of an acyclic quiver.
- Recently, Margolis, Saliola and I proved that if M is an LRB such that the Hasse diagram of the poset of principal right ideals aM of M is a tree, then kM is hereditary.
- This includes the algebras of nearly all LRBs considered by Brown except the hyperplane face semigroups!
- We also showed that **every** basic hereditary algebra is an LRB algebra.

Hereditary Algebras and Left Regular Bands

- An algebra A is **hereditary** if each left ideal of A is projective.
- Equivalently, each submodule of a projective A -module is projective.
- Gabriel proved that a basic algebra is hereditary iff it is the path algebra of an acyclic quiver.
- Recently, Margolis, Saliola and I proved that if M is an LRB such that the Hasse diagram of the poset of principal right ideals aM of M is a tree, then kM is hereditary.
- This includes the algebras of nearly all LRBs considered by Brown except the hyperplane face semigroups!
- We also showed that **every** basic hereditary algebra is an LRB algebra.

Hereditary Algebras and Left Regular Bands

- An algebra A is **hereditary** if each left ideal of A is projective.
- Equivalently, each submodule of a projective A -module is projective.
- Gabriel proved that a basic algebra is hereditary iff it is the path algebra of an acyclic quiver.
- Recently, Margolis, Saliola and I proved that if M is an LRB such that the Hasse diagram of the poset of principal right ideals aM of M is a tree, then kM is hereditary.
- This includes the algebras of nearly all LRBs considered by Brown except the hyperplane face semigroups!
- We also showed that **every** basic hereditary algebra is an LRB algebra.

Hereditary Algebras and Left Regular Bands

- An algebra A is **hereditary** if each left ideal of A is projective.
- Equivalently, each submodule of a projective A -module is projective.
- Gabriel proved that a basic algebra is hereditary iff it is the path algebra of an acyclic quiver.
- Recently, Margolis, Saliola and I proved that if M is an LRB such that the Hasse diagram of the poset of principal right ideals aM of M is a tree, then kM is hereditary.
- This includes the algebras of nearly all LRBs considered by Brown except the hyperplane face semigroups!
- We also showed that **every** basic hereditary algebra is an LRB algebra.

The Mantaci-Reutenauer Descent Algebra

- Mantaci and Reutenauer introduced a generalization of the descent algebra for wreath products $G \wr S_n$ with G abelian.
- Generalizations have since been considered for G non-abelian, independently, by Baumann and Hohlweg, and by Novelli and Thibon.
- Again the descent algebra can be viewed as a non-commutative ring of characters.
- Hsiao introduced a certain subsemigroup M of $F(\mathcal{H}_{S_n}) \times G^n$.
- S_n acts naturally on kM and the algebra of S_n -invariants is the Mantaci-Reutenauer descent algebra.
- M has a grading by an LRB with homogeneous components groups and hence is a left regular band of groups.

The Mantaci-Reutenauer Descent Algebra

- Mantaci and Reutenauer introduced a generalization of the descent algebra for wreath products $G \wr S_n$ with G abelian.
- Generalizations have since been considered for G non-abelian, independently, by Baumann and Hohlweg, and by Novelli and Thibon.
- Again the descent algebra can be viewed as a non-commutative ring of characters.
- Hsiao introduced a certain subsemigroup M of $F(\mathcal{H}_{S_n}) \times G^n$.
- S_n acts naturally on kM and the algebra of S_n -invariants is the Mantaci-Reutenauer descent algebra.
- M has a grading by an LRB with homogeneous components groups and hence is a left regular band of groups.

The Mantaci-Reutenauer Descent Algebra

- Mantaci and Reutenauer introduced a generalization of the descent algebra for wreath products $G \wr S_n$ with G abelian.
- Generalizations have since been considered for G non-abelian, independently, by Baumann and Hohlweg, and by Novelli and Thibon.
- Again the descent algebra can be viewed as a non-commutative ring of characters.
- Hsiao introduced a certain subsemigroup M of $F(\mathcal{H}_{S_n}) \times G^n$.
- S_n acts naturally on kM and the algebra of S_n -invariants is the Mantaci-Reutenauer descent algebra.
- M has a grading by an LRB with homogeneous components groups and hence is a left regular band of groups.

The Mantaci-Reutenauer Descent Algebra

- Mantaci and Reutenauer introduced a generalization of the descent algebra for wreath products $G \wr S_n$ with G abelian.
- Generalizations have since been considered for G non-abelian, independently, by Baumann and Hohlweg, and by Novelli and Thibon.
- Again the descent algebra can be viewed as a non-commutative ring of characters.
- Hsiao introduced a certain subsemigroup M of $F(\mathcal{H}_{S_n}) \times G^n$.
- S_n acts naturally on kM and the algebra of S_n -invariants is the Mantaci-Reutenauer descent algebra.
- M has a grading by an LRB with homogeneous components groups and hence is a left regular band of groups.

The Mantaci-Reutenauer Descent Algebra

- Mantaci and Reutenauer introduced a generalization of the descent algebra for wreath products $G \wr S_n$ with G abelian.
- Generalizations have since been considered for G non-abelian, independently, by Baumann and Hohlweg, and by Novelli and Thibon.
- Again the descent algebra can be viewed as a non-commutative ring of characters.
- Hsiao introduced a certain subsemigroup M of $F(\mathcal{H}_{S_n}) \times G^n$.
- S_n acts naturally on kM and the algebra of S_n -invariants is the Mantaci-Reutenauer descent algebra.
- M has a grading by an LRB with homogeneous components groups and hence is a left regular band of groups.

The Mantaci-Reutenauer Descent Algebra

- Mantaci and Reutenauer introduced a generalization of the descent algebra for wreath products $G \wr S_n$ with G abelian.
- Generalizations have since been considered for G non-abelian, independently, by Baumann and Hohlweg, and by Novelli and Thibon.
- Again the descent algebra can be viewed as a non-commutative ring of characters.
- Hsiao introduced a certain subsemigroup M of $F(\mathcal{H}_{S_n}) \times G^n$.
- S_n acts naturally on kM and the algebra of S_n -invariants is the Mantaci-Reutenauer descent algebra.
- M has a grading by an LRB with homogeneous components groups and hence is a **left regular band of groups**.

Representation Theory of Left Regular Bands of Groups

- In previous work, we studied the representation theory of left regular bands of groups.
- We computed the quiver, Cartan invariants and gave an explicit construction of the projective indecomposables.
- In particular, we computed the quiver of Hsiao's semigroup.
- The computation reduced to decomposing certain modules over group algebras.

Representation Theory of Left Regular Bands of Groups

- In previous work, we studied the representation theory of left regular bands of groups.
- We computed the quiver, Cartan invariants and gave an explicit construction of the projective indecomposables.
- In particular, we computed the quiver of Hsiao's semigroup.
- The computation reduced to decomposing certain modules over group algebras.

Representation Theory of Left Regular Bands of Groups

- In previous work, we studied the representation theory of left regular bands of groups.
- We computed the quiver, Cartan invariants and gave an explicit construction of the projective indecomposables.
- In particular, we computed the quiver of Hsiao's semigroup.
- The computation reduced to decomposing certain modules over group algebras.

Representation Theory of Left Regular Bands of Groups

- In previous work, we studied the representation theory of left regular bands of groups.
- We computed the quiver, Cartan invariants and gave an explicit construction of the projective indecomposables.
- In particular, we computed the quiver of Hsiao's semigroup.
- The computation reduced to decomposing certain modules over group algebras.

0-Hecke Monoids

- Let W be a finite Coxeter group with Coxeter generators S .
- The **Hecke algebra** $\mathcal{H}(W, S)$ has generators S and relations the braid relations of W and the relations

$$s^2 = (q - 1)s + 1, \text{ for } s \in S.$$

- When $q = 1$, one recovers the group algebra kW .
- When $q = 0$, one obtains the monoid algebra of the 0-Hecke monoid $M(W)$.
- $M(W)$ is the monoid generated by S subject to the braid relations of W and the idempotent relations $s^2 = s$ for $s \in S$.

0-Hecke Monoids

- Let W be a finite Coxeter group with Coxeter generators S .
- The **Hecke algebra** $\mathcal{H}(W, S)$ has generators S and relations the braid relations of W and the relations

$$s^2 = (q - 1)s + 1, \text{ for } s \in S.$$

- When $q = 1$, one recovers the group algebra kW .
- When $q = 0$, one obtains the monoid algebra of the **0-Hecke monoid** $M(W)$.
- $M(W)$ is the monoid generated by S subject to the braid relations of W and the idempotent relations $s^2 = s$ for $s \in S$.

0-Hecke Monoids

- Let W be a finite Coxeter group with Coxeter generators S .
- The **Hecke algebra** $\mathcal{H}(W, S)$ has generators S and relations the braid relations of W and the relations

$$s^2 = (q - 1)s + 1, \text{ for } s \in S.$$

- When $q = 1$, one recovers the group algebra kW .
- When $q = 0$, one obtains the monoid algebra of the **0-Hecke monoid** $M(W)$.
- $M(W)$ is the monoid generated by S subject to the braid relations of W and the idempotent relations $s^2 = s$ for $s \in S$.

0-Hecke Monoids

- Let W be a finite Coxeter group with Coxeter generators S .
- The **Hecke algebra** $\mathcal{H}(W, S)$ has generators S and relations the braid relations of W and the relations

$$s^2 = (q - 1)s + 1, \text{ for } s \in S.$$

- When $q = 1$, one recovers the group algebra kW .
- When $q = 0$, one obtains the monoid algebra of the **0-Hecke monoid** $M(W)$.
- $M(W)$ is the monoid generated by S subject to the braid relations of W and the idempotent relations $s^2 = s$ for $s \in S$.

0-Hecke Monoids

- Let W be a finite Coxeter group with Coxeter generators S .
- The **Hecke algebra** $\mathcal{H}(W, S)$ has generators S and relations the braid relations of W and the relations

$$s^2 = (q - 1)s + 1, \text{ for } s \in S.$$

- When $q = 1$, one recovers the group algebra kW .
- When $q = 0$, one obtains the monoid algebra of the **0-Hecke monoid** $M(W)$.
- $M(W)$ is the monoid generated by S subject to the braid relations of W and the idempotent relations $s^2 = s$ for $s \in S$.

0-Hecke Monoids Continued

- Tits's solution to the word problem in Coxeter groups also solves the word problem in $M(W)$.
- In particular, two reduced words are equal in W iff they are equal in $M(W)$.
- So $M(W)$ is in bijection with W .
- The strong Bruhat ordering is compatible on $M(W)$.
- From this one deduces $M(W)$ is \mathcal{J} -trivial.
- A monoid M is \mathcal{J} -trivial if $MaM = MbM \implies a = b$.
- The representation theory of 0-Hecke algebras was first studied by Norton in the 70s.
- Denton, Hivert, Thiéry and Schilling obtained the classical results, and new ones, as a special case of their theory of representations of \mathcal{J} -trivial monoids.

0-Hecke Monoids Continued

- Tits's solution to the word problem in Coxeter groups also solves the word problem in $M(W)$.
- In particular, two reduced words are equal in W iff they are equal in $M(W)$.
- So $M(W)$ is in bijection with W .
- The strong Bruhat ordering is compatible on $M(W)$.
- From this one deduces $M(W)$ is \mathcal{J} -trivial.
- A monoid M is \mathcal{J} -trivial if $MaM = MbM \implies a = b$.
- The representation theory of 0-Hecke algebras was first studied by Norton in the 70s.
- Denton, Hivert, Thiéry and Schilling obtained the classical results, and new ones, as a special case of their theory of representations of \mathcal{J} -trivial monoids.

0-Hecke Monoids Continued

- Tits's solution to the word problem in Coxeter groups also solves the word problem in $M(W)$.
- In particular, two reduced words are equal in W iff they are equal in $M(W)$.
- So $M(W)$ is in bijection with W .
- The strong Bruhat ordering is compatible on $M(W)$.
- From this one deduces $M(W)$ is \mathcal{J} -trivial.
- A monoid M is \mathcal{J} -trivial if $MaM = MbM \implies a = b$.
- The representation theory of 0-Hecke algebras was first studied by Norton in the 70s.
- Denton, Hivert, Thiéry and Schilling obtained the classical results, and new ones, as a special case of their theory of representations of \mathcal{J} -trivial monoids.

0-Hecke Monoids Continued

- Tits's solution to the word problem in Coxeter groups also solves the word problem in $M(W)$.
- In particular, two reduced words are equal in W iff they are equal in $M(W)$.
- So $M(W)$ is in bijection with W .
- The strong Bruhat ordering is compatible on $M(W)$.
- From this one deduces $M(W)$ is \mathcal{J} -trivial.
- A monoid M is \mathcal{J} -trivial if $MaM = MbM \implies a = b$.
- The representation theory of 0-Hecke algebras was first studied by Norton in the 70s.
- Denton, Hivert, Thiéry and Schilling obtained the classical results, and new ones, as a special case of their theory of representations of \mathcal{J} -trivial monoids.

0-Hecke Monoids Continued

- Tits's solution to the word problem in Coxeter groups also solves the word problem in $M(W)$.
- In particular, two reduced words are equal in W iff they are equal in $M(W)$.
- So $M(W)$ is in bijection with W .
- The strong Bruhat ordering is compatible on $M(W)$.
- From this one deduces $M(W)$ is \mathcal{J} -trivial.
- A monoid M is \mathcal{J} -trivial if $MaM = MbM \implies a = b$.
- The representation theory of 0-Hecke algebras was first studied by Norton in the 70s.
- Denton, Hivert, Thiéry and Schilling obtained the classical results, and new ones, as a special case of their theory of representations of \mathcal{J} -trivial monoids.

0-Hecke Monoids Continued

- Tits's solution to the word problem in Coxeter groups also solves the word problem in $M(W)$.
- In particular, two reduced words are equal in W iff they are equal in $M(W)$.
- So $M(W)$ is in bijection with W .
- The strong Bruhat ordering is compatible on $M(W)$.
- From this one deduces $M(W)$ is \mathcal{J} -trivial.
- A monoid M is \mathcal{J} -trivial if $MaM = MbM \implies a = b$.
- The representation theory of 0-Hecke algebras was first studied by Norton in the 70s.
- Denton, Hivert, Thiéry and Schilling obtained the classical results, and new ones, as a special case of their theory of representations of \mathcal{J} -trivial monoids.

0-Hecke Monoids Continued

- Tits's solution to the word problem in Coxeter groups also solves the word problem in $M(W)$.
- In particular, two reduced words are equal in W iff they are equal in $M(W)$.
- So $M(W)$ is in bijection with W .
- The strong Bruhat ordering is compatible on $M(W)$.
- From this one deduces $M(W)$ is \mathcal{J} -trivial.
- A monoid M is \mathcal{J} -trivial if $MaM = MbM \implies a = b$.
- The representation theory of 0-Hecke algebras was first studied by Norton in the 70s.
- Denton, Hivert, Thiéry and Schilling obtained the classical results, and new ones, as a special case of their theory of representations of \mathcal{J} -trivial monoids.

0-Hecke Monoids Continued

- Tits's solution to the word problem in Coxeter groups also solves the word problem in $M(W)$.
- In particular, two reduced words are equal in W iff they are equal in $M(W)$.
- So $M(W)$ is in bijection with W .
- The strong Bruhat ordering is compatible on $M(W)$.
- From this one deduces $M(W)$ is \mathcal{J} -trivial.
- A monoid M is \mathcal{J} -trivial if $MaM = MbM \implies a = b$.
- The representation theory of 0-Hecke algebras was first studied by Norton in the 70s.
- Denton, Hivert, Thiéry and Schilling obtained the classical results, and new ones, as a special case of their theory of representations of \mathcal{J} -trivial monoids.

\mathcal{J} -trivial Semigroups

- Path algebras and truncated path algebras are semigroup algebras of \mathcal{J} -trivial semigroups.
- So are incidence algebras of posets.
- Bautista, Gabriel, Roiter and Salmerón have shown that every basic algebra of finite representation type is the algebra of a \mathcal{J} -trivial semigroup.
- The Catalan monoid C_n is \mathcal{J} -trivial.
- It consists of all maps $f: [n] \rightarrow [n]$ such that f is order-preserving and non-increasing.
- $|C_n|$ is the n^{th} -Catalan number.
- Denton *et al.* have computed quivers, primitive idempotents, Cartan invariants and projective indecomposables for \mathcal{J} -trivial monoids.

\mathcal{J} -trivial Semigroups

- Path algebras and truncated path algebras are semigroup algebras of \mathcal{J} -trivial semigroups.
- So are incidence algebras of posets.
- Bautista, Gabriel, Roiter and Salmerón have shown that every basic algebra of finite representation type is the algebra of a \mathcal{J} -trivial semigroup.
- The Catalan monoid C_n is \mathcal{J} -trivial.
- It consists of all maps $f: [n] \rightarrow [n]$ such that f is order-preserving and non-increasing.
- $|C_n|$ is the n^{th} -Catalan number.
- Denton *et al.* have computed quivers, primitive idempotents, Cartan invariants and projective indecomposables for \mathcal{J} -trivial monoids.

\mathcal{J} -trivial Semigroups

- Path algebras and truncated path algebras are semigroup algebras of \mathcal{J} -trivial semigroups.
- So are incidence algebras of posets.
- Bautista, Gabriel, Roiter and Salmerón have shown that every basic algebra of finite representation type is the algebra of a \mathcal{J} -trivial semigroup.
- The Catalan monoid C_n is \mathcal{J} -trivial.
- It consists of all maps $f: [n] \rightarrow [n]$ such that f is order-preserving and non-increasing.
- $|C_n|$ is the n^{th} -Catalan number.
- Denton *et al.* have computed quivers, primitive idempotents, Cartan invariants and projective indecomposables for \mathcal{J} -trivial monoids.

\mathcal{J} -trivial Semigroups

- Path algebras and truncated path algebras are semigroup algebras of \mathcal{J} -trivial semigroups.
- So are incidence algebras of posets.
- Bautista, Gabriel, Roiter and Salmerón have shown that every basic algebra of finite representation type is the algebra of a \mathcal{J} -trivial semigroup.
- The **Catalan monoid** C_n is \mathcal{J} -trivial.
- It consists of all maps $f: [n] \rightarrow [n]$ such that f is order-preserving and non-increasing.
- $|C_n|$ is the n^{th} -Catalan number.
- Denton *et al.* have computed quivers, primitive idempotents, Cartan invariants and projective indecomposables for \mathcal{J} -trivial monoids.

\mathcal{J} -trivial Semigroups

- Path algebras and truncated path algebras are semigroup algebras of \mathcal{J} -trivial semigroups.
- So are incidence algebras of posets.
- Bautista, Gabriel, Roiter and Salmerón have shown that every basic algebra of finite representation type is the algebra of a \mathcal{J} -trivial semigroup.
- The **Catalan monoid** C_n is \mathcal{J} -trivial.
- It consists of all maps $f: [n] \rightarrow [n]$ such that f is order-preserving and non-increasing.
- $|C_n|$ is the n^{th} -Catalan number.
- Denton *et al.* have computed quivers, primitive idempotents, Cartan invariants and projective indecomposables for \mathcal{J} -trivial monoids.

\mathcal{J} -trivial Semigroups

- Path algebras and truncated path algebras are semigroup algebras of \mathcal{J} -trivial semigroups.
- So are incidence algebras of posets.
- Bautista, Gabriel, Roiter and Salmerón have shown that every basic algebra of finite representation type is the algebra of a \mathcal{J} -trivial semigroup.
- The **Catalan monoid** C_n is \mathcal{J} -trivial.
- It consists of all maps $f: [n] \rightarrow [n]$ such that f is order-preserving and non-increasing.
- $|C_n|$ is the n^{th} -Catalan number.
- Denton *et al.* have computed quivers, primitive idempotents, Cartan invariants and projective indecomposables for \mathcal{J} -trivial monoids.

\mathcal{J} -trivial Semigroups

- Path algebras and truncated path algebras are semigroup algebras of \mathcal{J} -trivial semigroups.
- So are incidence algebras of posets.
- Bautista, Gabriel, Roiter and Salmerón have shown that every basic algebra of finite representation type is the algebra of a \mathcal{J} -trivial semigroup.
- The **Catalan monoid** C_n is \mathcal{J} -trivial.
- It consists of all maps $f: [n] \rightarrow [n]$ such that f is order-preserving and non-increasing.
- $|C_n|$ is the n^{th} -Catalan number.
- Denton *et al.* have computed quivers, primitive idempotents, Cartan invariants and projective indecomposables for \mathcal{J} -trivial monoids.

\mathcal{R} -trivial Monoids

- A monoid M is \mathcal{R} -trivial if $aM = bM \implies a = b$.
- LRBs are precisely the \mathcal{R} -trivial bands.
- \mathcal{J} -trivial implies \mathcal{R} -trivial.
- A typical example is E_n , the monoid of maps $f: [n] \rightarrow [n]$ such that $xf \leq x$ (with action on the right).
- $|E_n| = n!$.
- Each \mathcal{R} -trivial monoid of size n embeds in E_n .
- Berg, N. Bergeron, Bhargava and Salvi have computed primitive idempotents for \mathcal{R} -trivial monoids.
- We have computed quivers, Cartan invariants and semigroup theoretic models of the projective indecomposables.
- Our results are independent of Berg *et al.*

\mathcal{R} -trivial Monoids

- A monoid M is \mathcal{R} -trivial if $aM = bM \implies aM = bM$.
- LRBs are precisely the \mathcal{R} -trivial bands.
- \mathcal{J} -trivial implies \mathcal{R} -trivial.
- A typical example is E_n , the monoid of maps $f: [n] \rightarrow [n]$ such that $xf \leq x$ (with action on the right).
- $|E_n| = n!$.
- Each \mathcal{R} -trivial monoid of size n embeds in E_n .
- Berg, N. Bergeron, Bhargava and Salviola have computed primitive idempotents for \mathcal{R} -trivial monoids.
- We have computed quivers, Cartan invariants and semigroup theoretic models of the projective indecomposables.
- Our results are independent of Berg *et al.*

\mathcal{R} -trivial Monoids

- A monoid M is \mathcal{R} -trivial if $aM = bM \implies aM = bM$.
- LRBs are precisely the \mathcal{R} -trivial bands.
- \mathcal{J} -trivial implies \mathcal{R} -trivial.
- A typical example is E_n , the monoid of maps $f: [n] \rightarrow [n]$ such that $xf \leq x$ (with action on the right).
- $|E_n| = n!$.
- Each \mathcal{R} -trivial monoid of size n embeds in E_n .
- Berg, N. Bergeron, Bhargava and Salviola have computed primitive idempotents for \mathcal{R} -trivial monoids.
- We have computed quivers, Cartan invariants and semigroup theoretic models of the projective indecomposables.
- Our results are independent of Berg *et al.*

\mathcal{R} -trivial Monoids

- A monoid M is \mathcal{R} -trivial if $aM = bM \implies aM = bM$.
- LRBs are precisely the \mathcal{R} -trivial bands.
- \mathcal{J} -trivial implies \mathcal{R} -trivial.
- A typical example is E_n , the monoid of maps $f: [n] \rightarrow [n]$ such that $xf \leq x$ (with action on the right).
- $|E_n| = n!$.
- Each \mathcal{R} -trivial monoid of size n embeds in E_n .
- Berg, N. Bergeron, Bhargava and Salviola have computed primitive idempotents for \mathcal{R} -trivial monoids.
- We have computed quivers, Cartan invariants and semigroup theoretic models of the projective indecomposables.
- Our results are independent of Berg *et al.*

\mathcal{R} -trivial Monoids

- A monoid M is \mathcal{R} -trivial if $aM = bM \implies aM = bM$.
- LRBs are precisely the \mathcal{R} -trivial bands.
- \mathcal{J} -trivial implies \mathcal{R} -trivial.
- A typical example is E_n , the monoid of maps $f: [n] \rightarrow [n]$ such that $xf \leq x$ (with action on the right).
- $|E_n| = n!$.
- Each \mathcal{R} -trivial monoid of size n embeds in E_n .
- Berg, N. Bergeron, Bhargava and Saliola have computed primitive idempotents for \mathcal{R} -trivial monoids.
- We have computed quivers, Cartan invariants and semigroup theoretic models of the projective indecomposables.
- Our results are independent of Berg *et al.*

\mathcal{R} -trivial Monoids

- A monoid M is \mathcal{R} -trivial if $aM = bM \implies a = b$.
- LRBs are precisely the \mathcal{R} -trivial bands.
- \mathcal{I} -trivial implies \mathcal{R} -trivial.
- A typical example is E_n , the monoid of maps $f: [n] \rightarrow [n]$ such that $xf \leq x$ (with action on the right).
- $|E_n| = n!$.
- Each \mathcal{R} -trivial monoid of size n embeds in E_n .
- Berg, N. Bergeron, Bhargava and Saliola have computed primitive idempotents for \mathcal{R} -trivial monoids.
- We have computed quivers, Cartan invariants and semigroup theoretic models of the projective indecomposables.
- Our results are independent of Berg *et al.*

\mathcal{R} -trivial Monoids

- A monoid M is \mathcal{R} -trivial if $aM = bM \implies aM = bM$.
- LRBs are precisely the \mathcal{R} -trivial bands.
- \mathcal{J} -trivial implies \mathcal{R} -trivial.
- A typical example is E_n , the monoid of maps $f: [n] \rightarrow [n]$ such that $xf \leq x$ (with action on the right).
- $|E_n| = n!$.
- Each \mathcal{R} -trivial monoid of size n embeds in E_n .
- Berg, N. Bergeron, Bhargava and Saliola have computed primitive idempotents for \mathcal{R} -trivial monoids.
- We have computed quivers, Cartan invariants and semigroup theoretic models of the projective indecomposables.
- Our results are independent of Berg *et al.*

\mathcal{R} -trivial Monoids

- A monoid M is \mathcal{R} -trivial if $aM = bM \implies aM = bM$.
- LRBs are precisely the \mathcal{R} -trivial bands.
- \mathcal{J} -trivial implies \mathcal{R} -trivial.
- A typical example is E_n , the monoid of maps $f: [n] \rightarrow [n]$ such that $xf \leq x$ (with action on the right).
- $|E_n| = n!$.
- Each \mathcal{R} -trivial monoid of size n embeds in E_n .
- Berg, N. Bergeron, Bhargava and Saliola have computed primitive idempotents for \mathcal{R} -trivial monoids.
- We have computed quivers, Cartan invariants and semigroup theoretic models of the projective indecomposables.
- Our results are independent of Berg *et al*.

\mathcal{R} -trivial Monoids

- A monoid M is \mathcal{R} -trivial if $aM = bM \implies aM = bM$.
- LRBs are precisely the \mathcal{R} -trivial bands.
- \mathcal{J} -trivial implies \mathcal{R} -trivial.
- A typical example is E_n , the monoid of maps $f: [n] \rightarrow [n]$ such that $xf \leq x$ (with action on the right).
- $|E_n| = n!$.
- Each \mathcal{R} -trivial monoid of size n embeds in E_n .
- Berg, N. Bergeron, Bhargava and Saliola have computed primitive idempotents for \mathcal{R} -trivial monoids.
- We have computed quivers, Cartan invariants and semigroup theoretic models of the projective indecomposables.
- Our results are independent of Berg *et al*.

Finite Monoids

- Let M be a finite monoid.
- $E(M)$ is the set of **idempotents** of M .

Theorem (Standard Facts)

For $e, f \in E(M)$, the following are equivalent:

- 1 $Me \cong Mf$ as left M -sets
- 2 $eM \cong fM$ as right M -sets
- 3 $\exists a, b \in M$ such that $ab = e$ and $ba = f$
- 4 $MeM = MfM$

- If the above conditions hold, e, f are said to be **conjugate**.

Finite Monoids

- Let M be a finite monoid.
- $E(M)$ is the set of **idempotents** of M .

Theorem (Standard Facts)

For $e, f \in E(M)$, the following are equivalent:

- 1 $Me \cong Mf$ as left M -sets
- 2 $eM \cong fM$ as right M -sets
- 3 $\exists a, b \in M$ such that $ab = e$ and $ba = f$
- 4 $MeM = MfM$

- If the above conditions hold, e, f are said to be conjugate.

Finite Monoids

- Let M be a finite monoid.
- $E(M)$ is the set of **idempotents** of M .

Theorem (Standard Facts)

For $e, f \in E(M)$, the following are equivalent:

- 1 $Me \cong Mf$ as left M -sets
- 2 $eM \cong fM$ as right M -sets
- 3 $\exists a, b \in M$ such that $ab = e$ and $ba = f$
- 4 $MeM = MfM$

- If the above conditions hold, e, f are said to be conjugate.

Finite Monoids

- Let M be a finite monoid.
- $E(M)$ is the set of **idempotents** of M .

Theorem (Standard Facts)

For $e, f \in E(M)$, the following are equivalent:

- 1 $Me \cong Mf$ as left M -sets
- 2 $eM \cong fM$ as right M -sets
- 3 $\exists a, b \in M$ such that $ab = e$ and $ba = f$
- 4 $MeM = MfM$

- If the above conditions hold, e, f are said to be conjugate.

Finite Monoids

- Let M be a finite monoid.
- $E(M)$ is the set of **idempotents** of M .

Theorem (Standard Facts)

For $e, f \in E(M)$, the following are equivalent:

- 1 $Me \cong Mf$ as left M -sets
- 2 $eM \cong fM$ as right M -sets
- 3 $\exists a, b \in M$ such that $ab = e$ and $ba = f$
- 4 $MeM = MfM$

- If the above conditions hold, e, f are said to be conjugate.

Finite Monoids

- Let M be a finite monoid.
- $E(M)$ is the set of **idempotents** of M .

Theorem (Standard Facts)

For $e, f \in E(M)$, the following are equivalent:

- 1 $Me \cong Mf$ as left M -sets
- 2 $eM \cong fM$ as right M -sets
- 3 $\exists a, b \in M$ such that $ab = e$ and $ba = f$
- 4 $MeM = MfM$

- If the above conditions hold, e, f are said to be **conjugate**.

Finite Monoids

- Let M be a finite monoid.
- $E(M)$ is the set of **idempotents** of M .

Theorem (Standard Facts)

For $e, f \in E(M)$, the following are equivalent:

- 1 $Me \cong Mf$ as left M -sets
- 2 $eM \cong fM$ as right M -sets
- 3 $\exists a, b \in M$ such that $ab = e$ and $ba = f$
- 4 $MeM = MfM$

- If the above conditions hold, e, f are said to be **conjugate**.

Finite Monoids

- Let M be a finite monoid.
- $E(M)$ is the set of **idempotents** of M .

Theorem (Standard Facts)

For $e, f \in E(M)$, the following are equivalent:

- 1 $Me \cong Mf$ as left M -sets
- 2 $eM \cong fM$ as right M -sets
- 3 $\exists a, b \in M$ such that $ab = e$ and $ba = f$
- 4 $MeM = MfM$

- If the above conditions hold, e, f are said to be **conjugate**.

Maximal Subgroups

- If $e \in E(M)$, then $\text{End}_M(Me) = eMe$.
- Let G_e be the group of units of eMe .
- $G_e = \text{Aut}_M(Me)$ is called the **maximal subgroup** at e .
- Notice e, f conjugate implies $G_e \cong G_f$.
- Much of finite semigroup theory is about using maximal subgroups to understand the whole semigroup.
- For instance, the maximal subgroups control the irreducible representations of the semigroup.

Maximal Subgroups

- If $e \in E(M)$, then $\text{End}_M(Me) = eMe$.
- Let G_e be the group of units of eMe .
- $G_e = \text{Aut}_M(Me)$ is called the **maximal subgroup** at e .
- Notice e, f conjugate implies $G_e \cong G_f$.
- Much of finite semigroup theory is about using maximal subgroups to understand the whole semigroup.
- For instance, the maximal subgroups control the irreducible representations of the semigroup.

Maximal Subgroups

- If $e \in E(M)$, then $\text{End}_M(Me) = eMe$.
- Let G_e be the group of units of eMe .
- $G_e = \text{Aut}_M(Me)$ is called the **maximal subgroup** at e .
- Notice e, f conjugate implies $G_e \cong G_f$.
- Much of finite semigroup theory is about using maximal subgroups to understand the whole semigroup.
- For instance, the maximal subgroups control the irreducible representations of the semigroup.

Maximal Subgroups

- If $e \in E(M)$, then $\text{End}_M(Me) = eMe$.
- Let G_e be the group of units of eMe .
- $G_e = \text{Aut}_M(Me)$ is called the **maximal subgroup** at e .
- Notice e, f conjugate implies $G_e \cong G_f$.
- Much of finite semigroup theory is about using maximal subgroups to understand the whole semigroup.
- For instance, the maximal subgroups control the irreducible representations of the semigroup.

Maximal Subgroups

- If $e \in E(M)$, then $\text{End}_M(Me) = eMe$.
- Let G_e be the group of units of eMe .
- $G_e = \text{Aut}_M(Me)$ is called the **maximal subgroup** at e .
- Notice e, f conjugate implies $G_e \cong G_f$.
- Much of finite semigroup theory is about using maximal subgroups to understand the whole semigroup.
- For instance, the maximal subgroups control the irreducible representations of the semigroup.

Maximal Subgroups

- If $e \in E(M)$, then $\text{End}_M(Me) = eMe$.
- Let G_e be the group of units of eMe .
- $G_e = \text{Aut}_M(Me)$ is called the **maximal subgroup** at e .
- Notice e, f conjugate implies $G_e \cong G_f$.
- Much of finite semigroup theory is about using maximal subgroups to understand the whole semigroup.
- For instance, the maximal subgroups control the irreducible representations of the semigroup.

Rectangular Monoids

- We say M is a **rectangular monoid** if each conjugacy class of idempotents is a subsemigroup.
- For example, all groups are examples of rectangular monoids.
- All semigroups discussed in this talk are rectangular monoids.
- The class of rectangular monoids was introduced by Schützenberger and his school under the name **DO**.

Theorem (Almeida, Margolis, BS, Volkov)

The algebra kM of a finite monoid M is basic if and only if:

- *M is a rectangular monoid*
- *each maximal subgroup of M is abelian.*

Rectangular Monoids

- We say M is a **rectangular monoid** if each conjugacy class of idempotents is a subsemigroup.
- For example, all groups are examples of rectangular monoids.
- All semigroups discussed in this talk are rectangular monoids.
- The class of rectangular monoids was introduced by Schützenberger and his school under the name **DO**.

Theorem (Almeida, Margolis, BS, Volkov)

The algebra kM of a finite monoid M is basic if and only if:

- *M is a rectangular monoid*
- *each maximal subgroup of M is abelian.*

Rectangular Monoids

- We say M is a **rectangular monoid** if each conjugacy class of idempotents is a subsemigroup.
- For example, all groups are examples of rectangular monoids.
- All semigroups discussed in this talk are rectangular monoids.
- The class of rectangular monoids was introduced by Schützenberger and his school under the name **DO**.

Theorem (Almeida, Margolis, BS, Volkov)

The algebra kM of a finite monoid M is basic if and only if:

- 1 *M is a rectangular monoid*
- 2 *each maximal subgroup of M is abelian.*

Rectangular Monoids

- We say M is a **rectangular monoid** if each conjugacy class of idempotents is a subsemigroup.
- For example, all groups are examples of rectangular monoids.
- All semigroups discussed in this talk are rectangular monoids.
- The class of rectangular monoids was introduced by Schützenberger and his school under the name **DO**.

Theorem (Almeida, Margolis, BS, Volkov)

The algebra kM of a finite monoid M is basic if and only if:

- 1 *M is a rectangular monoid*
- 2 *each maximal subgroup of M is abelian.*

Rectangular Monoids

- We say M is a **rectangular monoid** if each conjugacy class of idempotents is a subsemigroup.
- For example, all groups are examples of rectangular monoids.
- All semigroups discussed in this talk are rectangular monoids.
- The class of rectangular monoids was introduced by Schützenberger and his school under the name **DO**.

Theorem (Almeida, Margolis, BS, Volkov)

The algebra kM of a finite monoid M is basic if and only if:

- 1 *M is a rectangular monoid*
- 2 *each maximal subgroup of M is abelian.*

Rectangular Monoids

- We say M is a **rectangular monoid** if each conjugacy class of idempotents is a subsemigroup.
- For example, all groups are examples of rectangular monoids.
- All semigroups discussed in this talk are rectangular monoids.
- The class of rectangular monoids was introduced by Schützenberger and his school under the name **DO**.

Theorem (Almeida, Margolis, BS, Volkov)

The algebra kM of a finite monoid M is basic if and only if:

- 1 *M is a rectangular monoid*
- 2 *each maximal subgroup of M is abelian.*

Rectangular Monoids

- We say M is a **rectangular monoid** if each conjugacy class of idempotents is a subsemigroup.
- For example, all groups are examples of rectangular monoids.
- All semigroups discussed in this talk are rectangular monoids.
- The class of rectangular monoids was introduced by Schützenberger and his school under the name **DO**.

Theorem (Almeida, Margolis, BS, Volkov)

The algebra kM of a finite monoid M is basic if and only if:

- 1 M is a rectangular monoid
- 2 each maximal subgroup of M is abelian.

An Example

- Our goal is to “compute” the quiver of a rectangular monoid.
- Let us give an example to explain what this means.
- Let G be a group and X a $G \times G$ -set.
- We view X as having commuting left and right actions by G .
- Let $M = G \cup X \cup \{0\}$ where $X^2 = 0$.
- M is a rectangular monoid.
- Computing the quiver of M turns out to be equivalent to decomposing the $G \times G$ -module kX into irreducibles.

An Example

- Our goal is to “compute” the quiver of a rectangular monoid.
- Let us give an example to explain what this means.
- Let G be a group and X a $G \times G$ -set.
- We view X as having commuting left and right actions by G .
- Let $M = G \cup X \cup \{0\}$ where $X^2 = 0$.
- M is a rectangular monoid.
- Computing the quiver of M turns out to be equivalent to decomposing the $G \times G$ -module kX into irreducibles.

An Example

- Our goal is to “compute” the quiver of a rectangular monoid.
- Let us give an example to explain what this means.
- Let G be a group and X a $G \times G$ -set.
- We view X as having commuting left and right actions by G .
- Let $M = G \cup X \cup \{0\}$ where $X^2 = 0$.
- M is a rectangular monoid.
- Computing the quiver of M turns out to be equivalent to decomposing the $G \times G$ -module kX into irreducibles.

An Example

- Our goal is to “compute” the quiver of a rectangular monoid.
- Let us give an example to explain what this means.
- Let G be a group and X a $G \times G$ -set.
- We view X as having commuting left and right actions by G .
- Let $M = G \cup X \cup \{0\}$ where $X^2 = 0$.
- M is a rectangular monoid.
- Computing the quiver of M turns out to be equivalent to decomposing the $G \times G$ -module kX into irreducibles.

An Example

- Our goal is to “compute” the quiver of a rectangular monoid.
- Let us give an example to explain what this means.
- Let G be a group and X a $G \times G$ -set.
- We view X as having commuting left and right actions by G .
- Let $M = G \cup X \cup \{0\}$ where $X^2 = 0$.
- M is a rectangular monoid.
- Computing the quiver of M turns out to be equivalent to decomposing the $G \times G$ -module kX into irreducibles.

An Example

- Our goal is to “compute” the quiver of a rectangular monoid.
- Let us give an example to explain what this means.
- Let G be a group and X a $G \times G$ -set.
- We view X as having commuting left and right actions by G .
- Let $M = G \cup X \cup \{0\}$ where $X^2 = 0$.
- M is a rectangular monoid.
- Computing the quiver of M turns out to be equivalent to decomposing the $G \times G$ -module kX into irreducibles.

An Example

- Our goal is to “compute” the quiver of a rectangular monoid.
- Let us give an example to explain what this means.
- Let G be a group and X a $G \times G$ -set.
- We view X as having commuting left and right actions by G .
- Let $M = G \cup X \cup \{0\}$ where $X^2 = 0$.
- M is a rectangular monoid.
- Computing the quiver of M turns out to be equivalent to decomposing the $G \times G$ -module kX into irreducibles.

An Example Continued

- Let V_1, \dots, V_s be the simple G -modules.
- Each is a simple M -module by letting $X \cup \{0\}$ act as 0.
- There is one additional simple M -module: the trivial module.
- The vertex of $Q(kM)$ corresponding to the trivial module is isolated.
- The number of arrows from V_i to V_j is the multiplicity of $V_j \otimes_k V_i^*$ as a composition factor of the $G \times G$ -module kX .
- Since every simple $G \times G$ -module is of this form, $Q(kM)$ encodes the composition factors of kX .
- So our goal is to compute the quivers of rectangular monoids modulo the character theory of finite groups.

An Example Continued

- Let V_1, \dots, V_s be the simple G -modules.
- Each is a simple M -module by letting $X \cup \{0\}$ act as 0.
- There is one additional simple M -module: the trivial module.
- The vertex of $Q(kM)$ corresponding to the trivial module is isolated.
- The number of arrows from V_i to V_j is the multiplicity of $V_j \otimes_k V_i^*$ as a composition factor of the $G \times G$ -module kX .
- Since every simple $G \times G$ -module is of this form, $Q(kM)$ encodes the composition factors of kX .
- So our goal is to compute the quivers of rectangular monoids modulo the character theory of finite groups.

An Example Continued

- Let V_1, \dots, V_s be the simple G -modules.
- Each is a simple M -module by letting $X \cup \{0\}$ act as 0.
- There is one additional simple M -module: the trivial module.
- The vertex of $Q(kM)$ corresponding to the trivial module is isolated.
- The number of arrows from V_i to V_j is the multiplicity of $V_j \otimes_k V_i^*$ as a composition factor of the $G \times G$ -module kX .
- Since every simple $G \times G$ -module is of this form, $Q(kM)$ encodes the composition factors of kX .
- So our goal is to compute the quivers of rectangular monoids modulo the character theory of finite groups.

An Example Continued

- Let V_1, \dots, V_s be the simple G -modules.
- Each is a simple M -module by letting $X \cup \{0\}$ act as 0.
- There is one additional simple M -module: the trivial module.
- The vertex of $Q(kM)$ corresponding to the trivial module is isolated.
- The number of arrows from V_i to V_j is the multiplicity of $V_j \otimes_k V_i^*$ as a composition factor of the $G \times G$ -module kX .
- Since every simple $G \times G$ -module is of this form, $Q(kM)$ encodes the composition factors of kX .
- So our goal is to compute the quivers of rectangular monoids modulo the character theory of finite groups.

An Example Continued

- Let V_1, \dots, V_s be the simple G -modules.
- Each is a simple M -module by letting $X \cup \{0\}$ act as 0.
- There is one additional simple M -module: the trivial module.
- The vertex of $Q(kM)$ corresponding to the trivial module is isolated.
- The number of arrows from V_i to V_j is the multiplicity of $V_j \otimes_k V_i^*$ as a composition factor of the $G \times G$ -module kX .
- Since every simple $G \times G$ -module is of this form, $Q(kM)$ encodes the composition factors of kX .
- So our goal is to compute the quivers of rectangular monoids modulo the character theory of finite groups.

An Example Continued

- Let V_1, \dots, V_s be the simple G -modules.
- Each is a simple M -module by letting $X \cup \{0\}$ act as 0.
- There is one additional simple M -module: the trivial module.
- The vertex of $Q(kM)$ corresponding to the trivial module is isolated.
- The number of arrows from V_i to V_j is the multiplicity of $V_j \otimes_k V_i^*$ as a composition factor of the $G \times G$ -module kX .
- Since every simple $G \times G$ -module is of this form, $Q(kM)$ encodes the composition factors of kX .
- So our goal is to compute the quivers of rectangular monoids modulo the character theory of finite groups.

An Example Continued

- Let V_1, \dots, V_s be the simple G -modules.
- Each is a simple M -module by letting $X \cup \{0\}$ act as 0.
- There is one additional simple M -module: the trivial module.
- The vertex of $Q(kM)$ corresponding to the trivial module is isolated.
- The number of arrows from V_i to V_j is the multiplicity of $V_j \otimes_k V_i^*$ as a composition factor of the $G \times G$ -module kX .
- Since every simple $G \times G$ -module is of this form, $Q(kM)$ encodes the composition factors of kX .
- So our goal is to compute the quivers of rectangular monoids modulo the character theory of finite groups.

The Simple Modules

- Let M be rectangular monoid.
- Let $\Lambda(M)$ be the set of conjugacy classes of idempotents.
- For each $X \in \Lambda(M)$, fix a maximal subgroup G_X .
- There is a surjective homomorphism $\rho_X: kM \rightarrow kG_X$.
- Each simple M -module is lifted via ρ_X from a simple G_X -module for some X .
- So

$$\text{Irr}(kM) = \coprod_{X \in \Lambda(M)} \text{Irr}(kG_X).$$

- In most of our examples, e.g., bands and \mathcal{R} -trivial monoids, the maximal subgroups are all trivial.

The Simple Modules

- Let M be rectangular monoid.
- Let $\Lambda(M)$ be the set of conjugacy classes of idempotents.
- For each $X \in \Lambda(M)$, fix a maximal subgroup G_X .
- There is a surjective homomorphism $\rho_X: kM \rightarrow kG_X$.
- Each simple M -module is lifted via ρ_X from a simple G_X -module for some X .
- So

$$\text{Irr}(kM) = \coprod_{X \in \Lambda(M)} \text{Irr}(kG_X).$$

- In most of our examples, e.g., bands and \mathcal{R} -trivial monoids, the maximal subgroups are all trivial.

The Simple Modules

- Let M be rectangular monoid.
- Let $\Lambda(M)$ be the set of conjugacy classes of idempotents.
- For each $X \in \Lambda(M)$, fix a maximal subgroup G_X .
- There is a surjective homomorphism $\rho_X: kM \rightarrow kG_X$.
- Each simple M -module is lifted via ρ_X from a simple G_X -module for some X .
- So

$$\text{Irr}(kM) = \coprod_{X \in \Lambda(M)} \text{Irr}(kG_X).$$

- In most of our examples, e.g., bands and \mathcal{R} -trivial monoids, the maximal subgroups are all trivial.

The Simple Modules

- Let M be rectangular monoid.
- Let $\Lambda(M)$ be the set of conjugacy classes of idempotents.
- For each $X \in \Lambda(M)$, fix a maximal subgroup G_X .
- There is a surjective homomorphism $\rho_X: kM \rightarrow kG_X$.
- Each simple M -module is lifted via ρ_X from a simple G_X -module for some X .
- So

$$\text{Irr}(kM) = \coprod_{X \in \Lambda(M)} \text{Irr}(kG_X).$$

- In most of our examples, e.g., bands and \mathcal{R} -trivial monoids, the maximal subgroups are all trivial.

The Simple Modules

- Let M be rectangular monoid.
- Let $\Lambda(M)$ be the set of conjugacy classes of idempotents.
- For each $X \in \Lambda(M)$, fix a maximal subgroup G_X .
- There is a surjective homomorphism $\rho_X: kM \rightarrow kG_X$.
- Each simple M -module is lifted via ρ_X from a simple G_X -module for some X .
- So

$$\text{Irr}(kM) = \coprod_{X \in \Lambda(M)} \text{Irr}(kG_X).$$

- In most of our examples, e.g., bands and \mathcal{R} -trivial monoids, the maximal subgroups are all trivial.

The Simple Modules

- Let M be rectangular monoid.
- Let $\Lambda(M)$ be the set of conjugacy classes of idempotents.
- For each $X \in \Lambda(M)$, fix a maximal subgroup G_X .
- There is a surjective homomorphism $\rho_X: kM \rightarrow kG_X$.
- Each simple M -module is lifted via ρ_X from a simple G_X -module for some X .
- So

$$\text{Irr}(kM) = \coprod_{X \in \Lambda(M)} \text{Irr}(kG_X).$$

- In most of our examples, e.g., bands and \mathcal{R} -trivial monoids, the maximal subgroups are all trivial.

The Simple Modules

- Let M be rectangular monoid.
- Let $\Lambda(M)$ be the set of conjugacy classes of idempotents.
- For each $X \in \Lambda(M)$, fix a maximal subgroup G_X .
- There is a surjective homomorphism $\rho_X: kM \rightarrow kG_X$.
- Each simple M -module is lifted via ρ_X from a simple G_X -module for some X .
- So

$$\text{Irr}(kM) = \coprod_{X \in \Lambda(M)} \text{Irr}(kG_X).$$

- In most of our examples, e.g., bands and \mathcal{R} -trivial monoids, the maximal subgroups are all trivial.

The Quiver of kM

- The vertex set of $Q(kM)$ is $\coprod_{X \in \Lambda(M)} \text{Irr}(kG_X)$.
- For each pair $(X, Y) \in \Lambda(M)^2$, we explicitly construct a certain $G_Y \times G_X$ -module $V_{X,Y}$.
- Suppose $U \in \text{Irr}(kG_X)$ and $V \in \text{Irr}(kG_Y)$.
- The number of edges $U \rightarrow V$ is the multiplicity of the simple $G_Y \times G_X$ -module $V \otimes_k U^*$ as a composition factor in $V_{X,Y}$.
- Thus computing $Q(kM)$ boils down to decomposing the $G_Y \times G_X$ -modules $V_{X,Y}$.
- When the maximal subgroups of M are trivial, the vertex set is $\Lambda(M)$.
- There are $\dim V_{X,Y}$ edges from X to Y .

The Quiver of kM

- The vertex set of $Q(kM)$ is $\coprod_{X \in \Lambda(M)} \text{Irr}(kG_X)$.
- For each pair $(X, Y) \in \Lambda(M)^2$, we explicitly construct a certain $G_Y \times G_X$ -module $V_{X,Y}$.
- Suppose $U \in \text{Irr}(kG_X)$ and $V \in \text{Irr}(kG_Y)$.
- The number of edges $U \rightarrow V$ is the multiplicity of the simple $G_Y \times G_X$ -module $V \otimes_k U^*$ as a composition factor in $V_{X,Y}$.
- Thus computing $Q(kM)$ boils down to decomposing the $G_Y \times G_X$ -modules $V_{X,Y}$.
- When the maximal subgroups of M are trivial, the vertex set is $\Lambda(M)$.
- There are $\dim V_{X,Y}$ edges from X to Y .

The Quiver of kM

- The vertex set of $Q(kM)$ is $\coprod_{X \in \Lambda(M)} \text{Irr}(kG_X)$.
- For each pair $(X, Y) \in \Lambda(M)^2$, we explicitly construct a certain $G_Y \times G_X$ -module $V_{X,Y}$.
- Suppose $U \in \text{Irr}(kG_X)$ and $V \in \text{Irr}(kG_Y)$.
- The number of edges $U \rightarrow V$ is the multiplicity of the simple $G_Y \times G_X$ -module $V \otimes_k U^*$ as a composition factor in $V_{X,Y}$.
- Thus computing $Q(kM)$ boils down to decomposing the $G_Y \times G_X$ -modules $V_{X,Y}$.
- When the maximal subgroups of M are trivial, the vertex set is $\Lambda(M)$.
- There are $\dim V_{X,Y}$ edges from X to Y .

The Quiver of kM

- The vertex set of $Q(kM)$ is $\coprod_{X \in \Lambda(M)} \text{Irr}(kG_X)$.
- For each pair $(X, Y) \in \Lambda(M)^2$, we explicitly construct a certain $G_Y \times G_X$ -module $V_{X,Y}$.
- Suppose $U \in \text{Irr}(kG_X)$ and $V \in \text{Irr}(kG_Y)$.
- The number of edges $U \rightarrow V$ is the multiplicity of the simple $G_Y \times G_X$ -module $V \otimes_k U^*$ as a composition factor in $V_{X,Y}$.
- Thus computing $Q(kM)$ boils down to decomposing the $G_Y \times G_X$ -modules $V_{X,Y}$.
- When the maximal subgroups of M are trivial, the vertex set is $\Lambda(M)$.
- There are $\dim V_{X,Y}$ edges from X to Y .

The Quiver of kM

- The vertex set of $Q(kM)$ is $\coprod_{X \in \Lambda(M)} \text{Irr}(kG_X)$.
- For each pair $(X, Y) \in \Lambda(M)^2$, we explicitly construct a certain $G_Y \times G_X$ -module $V_{X,Y}$.
- Suppose $U \in \text{Irr}(kG_X)$ and $V \in \text{Irr}(kG_Y)$.
- The number of edges $U \rightarrow V$ is the multiplicity of the simple $G_Y \times G_X$ -module $V \otimes_k U^*$ as a composition factor in $V_{X,Y}$.
- Thus computing $Q(kM)$ boils down to decomposing the $G_Y \times G_X$ -modules $V_{X,Y}$.
- When the maximal subgroups of M are trivial, the vertex set is $\Lambda(M)$.
- There are $\dim V_{X,Y}$ edges from X to Y .

The Quiver of kM

- The vertex set of $Q(kM)$ is $\coprod_{X \in \Lambda(M)} \text{Irr}(kG_X)$.
- For each pair $(X, Y) \in \Lambda(M)^2$, we explicitly construct a certain $G_Y \times G_X$ -module $V_{X,Y}$.
- Suppose $U \in \text{Irr}(kG_X)$ and $V \in \text{Irr}(kG_Y)$.
- The number of edges $U \rightarrow V$ is the multiplicity of the simple $G_Y \times G_X$ -module $V \otimes_k U^*$ as a composition factor in $V_{X,Y}$.
- Thus computing $Q(kM)$ boils down to decomposing the $G_Y \times G_X$ -modules $V_{X,Y}$.
- When the maximal subgroups of M are trivial, the vertex set is $\Lambda(M)$.
- There are $\dim V_{X,Y}$ edges from X to Y .

The Quiver of kM

- The vertex set of $Q(kM)$ is $\coprod_{X \in \Lambda(M)} \text{Irr}(kG_X)$.
- For each pair $(X, Y) \in \Lambda(M)^2$, we explicitly construct a certain $G_Y \times G_X$ -module $V_{X,Y}$.
- Suppose $U \in \text{Irr}(kG_X)$ and $V \in \text{Irr}(kG_Y)$.
- The number of edges $U \rightarrow V$ is the multiplicity of the simple $G_Y \times G_X$ -module $V \otimes_k U^*$ as a composition factor in $V_{X,Y}$.
- Thus computing $Q(kM)$ boils down to decomposing the $G_Y \times G_X$ -modules $V_{X,Y}$.
- When the maximal subgroups of M are trivial, the vertex set is $\Lambda(M)$.
- There are $\dim V_{X,Y}$ edges from X to Y .

Hochschild Cohomology

- Let me give an idea of our techniques.
- Recall the number of edges from U to V is $\dim \operatorname{Ext}_{kM}^1(U, V)$.
- There is an alternative formulation in terms of primitive idempotents.
- We use the description of Ext in terms of Hochschild cohomology.
- Namely, $\operatorname{Hom}_k(U, V)$ is an M -bimodule in a natural way.
- $\operatorname{Ext}_{kM}^n(U, V) \cong H^n(M, \operatorname{Hom}_k(U, V))$.
- We only need $n = 1$, so we recall the description in this case.

Hochschild Cohomology

- Let me give an idea of our techniques.
- Recall the number of edges from U to V is $\dim \operatorname{Ext}_{kM}^1(U, V)$.
- There is an alternative formulation in terms of primitive idempotents.
- We use the description of Ext in terms of Hochschild cohomology.
- Namely, $\operatorname{Hom}_k(U, V)$ is an M -bimodule in a natural way.
- $\operatorname{Ext}_{kM}^n(U, V) \cong H^n(M, \operatorname{Hom}_k(U, V))$.
- We only need $n = 1$, so we recall the description in this case.

Hochschild Cohomology

- Let me give an idea of our techniques.
- Recall the number of edges from U to V is $\dim \operatorname{Ext}_{kM}^1(U, V)$.
- There is an alternative formulation in terms of primitive idempotents.
- We use the description of Ext in terms of **Hochschild cohomology**.
- Namely, $\operatorname{Hom}_k(U, V)$ is an M -bimodule in a natural way.
- $\operatorname{Ext}_{kM}^n(U, V) \cong H^n(M, \operatorname{Hom}_k(U, V))$.
- We only need $n = 1$, so we recall the description in this case.

Hochschild Cohomology

- Let me give an idea of our techniques.
- Recall the number of edges from U to V is $\dim \operatorname{Ext}_{kM}^1(U, V)$.
- There is an alternative formulation in terms of primitive idempotents.
- We use the description of Ext in terms of **Hochschild cohomology**.
- Namely, $\operatorname{Hom}_k(U, V)$ is an M -bimodule in a natural way.
- $\operatorname{Ext}_{kM}^n(U, V) \cong H^n(M, \operatorname{Hom}_k(U, V))$.
- We only need $n = 1$, so we recall the description in this case.

Hochschild Cohomology

- Let me give an idea of our techniques.
- Recall the number of edges from U to V is $\dim \operatorname{Ext}_{kM}^1(U, V)$.
- There is an alternative formulation in terms of primitive idempotents.
- We use the description of Ext in terms of **Hochschild cohomology**.
- Namely, $\operatorname{Hom}_k(U, V)$ is an M -bimodule in a natural way.
- $\operatorname{Ext}_{kM}^n(U, V) \cong H^n(M, \operatorname{Hom}_k(U, V))$.
- We only need $n = 1$, so we recall the description in this case.

Hochschild Cohomology

- Let me give an idea of our techniques.
- Recall the number of edges from U to V is $\dim \operatorname{Ext}_{kM}^1(U, V)$.
- There is an alternative formulation in terms of primitive idempotents.
- We use the description of Ext in terms of **Hochschild cohomology**.
- Namely, $\operatorname{Hom}_k(U, V)$ is an M -bimodule in a natural way.
- $\operatorname{Ext}_{kM}^n(U, V) \cong H^n(M, \operatorname{Hom}_k(U, V))$.
- We only need $n = 1$, so we recall the description in this case.

Hochschild Cohomology

- Let me give an idea of our techniques.
- Recall the number of edges from U to V is $\dim \operatorname{Ext}_{kM}^1(U, V)$.
- There is an alternative formulation in terms of primitive idempotents.
- We use the description of Ext in terms of **Hochschild cohomology**.
- Namely, $\operatorname{Hom}_k(U, V)$ is an M -bimodule in a natural way.
- $\operatorname{Ext}_{kM}^n(U, V) \cong H^n(M, \operatorname{Hom}_k(U, V))$.
- We only need $n = 1$, so we recall the description in this case.

Derivations

- Let A be an M -bimodule.
- A **derivation** of M in A is a mapping $d: M \rightarrow A$ satisfying

$$d(mn) = md(n) + d(m)n.$$

- A derivation is **inner** if there exists $a \in A$ with $d(m) = ma - am$.
- $H^1(M, A)$ is the space of outer derivations, i.e., derivations modulo inner derivations.

Derivations

- Let A be an M -bimodule.
- A **derivation** of M in A is a mapping $d: M \rightarrow A$ satisfying

$$d(mn) = md(n) + d(m)n.$$

- A derivation is **inner** if there exists $a \in A$ with $d(m) = ma - am$.
- $H^1(M, A)$ is the space of outer derivations, i.e., derivations modulo inner derivations.

Derivations

- Let A be an M -bimodule.
- A **derivation** of M in A is a mapping $d: M \rightarrow A$ satisfying

$$d(mn) = md(n) + d(m)n.$$

- A derivation is **inner** if there exists $a \in A$ with $d(m) = ma - am$.
- $H^1(M, A)$ is the space of outer derivations, i.e., derivations modulo inner derivations.

Derivations

- Let A be an M -bimodule.
- A **derivation** of M in A is a mapping $d: M \rightarrow A$ satisfying

$$d(mn) = md(n) + d(m)n.$$

- A derivation is **inner** if there exists $a \in A$ with $d(m) = ma - am$.
- $H^1(M, A)$ is the space of outer derivations, i.e., derivations modulo inner derivations.

Representability

- If $U \in \text{Irr}(G_X)$ and $V \in \text{Irr}(G_Y)$, then $\text{Hom}_k(U, V)$ is in fact a $G_Y \times G_X$ -module.
- More generally, any $G_Y \times G_X$ -module is naturally an M -bimodule.
- For each $X, Y \in \Lambda(M)$, we show $H^1(M, A)$ is **representable** as a functor from $G_Y \times G_X$ -modules to k -vector spaces.
- That is, there is a $G_Y \times G_X$ -module $V_{X,Y}$ such that $H^1(M, A) \cong \text{Hom}_{k[G_Y \times G_X]}(V_{X,Y}, A)$.
- $V_{X,Y}$ is explicitly constructed.
- The construction depends on the relative positions of the ideals MXM and MYM in M .

Representability

- If $U \in \text{Irr}(G_X)$ and $V \in \text{Irr}(G_Y)$, then $\text{Hom}_k(U, V)$ is in fact a $G_Y \times G_X$ -module.
- More generally, any $G_Y \times G_X$ -module is naturally an M -bimodule.
- For each $X, Y \in \Lambda(M)$, we show $H^1(M, A)$ is **representable** as a functor from $G_Y \times G_X$ -modules to k -vector spaces.
- That is, there is a $G_Y \times G_X$ -module $V_{X,Y}$ such that $H^1(M, A) \cong \text{Hom}_{k[G_Y \times G_X]}(V_{X,Y}, A)$.
- $V_{X,Y}$ is explicitly constructed.
- The construction depends on the relative positions of the ideals MXM and MYM in M .

Representability

- If $U \in \text{Irr}(G_X)$ and $V \in \text{Irr}(G_Y)$, then $\text{Hom}_k(U, V)$ is in fact a $G_Y \times G_X$ -module.
- More generally, any $G_Y \times G_X$ -module is naturally an M -bimodule.
- For each $X, Y \in \Lambda(M)$, we show $H^1(M, A)$ is **representable** as a functor from $G_Y \times G_X$ -modules to k -vector spaces.
- That is, there is a $G_Y \times G_X$ -module $V_{X,Y}$ such that $H^1(M, A) \cong \text{Hom}_{k[G_Y \times G_X]}(V_{X,Y}, A)$.
- $V_{X,Y}$ is explicitly constructed.
- The construction depends on the relative positions of the ideals MXM and MYM in M .

Representability

- If $U \in \text{Irr}(G_X)$ and $V \in \text{Irr}(G_Y)$, then $\text{Hom}_k(U, V)$ is in fact a $G_Y \times G_X$ -module.
- More generally, any $G_Y \times G_X$ -module is naturally an M -bimodule.
- For each $X, Y \in \Lambda(M)$, we show $H^1(M, A)$ is **representable** as a functor from $G_Y \times G_X$ -modules to k -vector spaces.
- That is, there is a $G_Y \times G_X$ -module $V_{X,Y}$ such that $H^1(M, A) \cong \text{Hom}_{k[G_Y \times G_X]}(V_{X,Y}, A)$.
- $V_{X,Y}$ is explicitly constructed.
- The construction depends on the relative positions of the ideals MXM and MYM in M .

Representability

- If $U \in \text{Irr}(G_X)$ and $V \in \text{Irr}(G_Y)$, then $\text{Hom}_k(U, V)$ is in fact a $G_Y \times G_X$ -module.
- More generally, any $G_Y \times G_X$ -module is naturally an M -bimodule.
- For each $X, Y \in \Lambda(M)$, we show $H^1(M, A)$ is **representable** as a functor from $G_Y \times G_X$ -modules to k -vector spaces.
- That is, there is a $G_Y \times G_X$ -module $V_{X,Y}$ such that $H^1(M, A) \cong \text{Hom}_{k[G_Y \times G_X]}(V_{X,Y}, A)$.
- $V_{X,Y}$ is explicitly constructed.
- The construction depends on the relative positions of the ideals MXM and MYM in M .

Representability

- If $U \in \text{Irr}(G_X)$ and $V \in \text{Irr}(G_Y)$, then $\text{Hom}_k(U, V)$ is in fact a $G_Y \times G_X$ -module.
- More generally, any $G_Y \times G_X$ -module is naturally an M -bimodule.
- For each $X, Y \in \Lambda(M)$, we show $H^1(M, A)$ is **representable** as a functor from $G_Y \times G_X$ -modules to k -vector spaces.
- That is, there is a $G_Y \times G_X$ -module $V_{X,Y}$ such that $H^1(M, A) \cong \text{Hom}_{k[G_Y \times G_X]}(V_{X,Y}, A)$.
- $V_{X,Y}$ is explicitly constructed.
- The construction depends on the relative positions of the ideals MXM and MYM in M .

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - ① $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - ② $x \equiv x'$ whenever there exists $e \in e_Y Me_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - ① $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - ② $x \equiv x'$ whenever there exists $e \in e_Y M e_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - ① $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - ② $x \equiv x'$ whenever there exists $e \in e_Y M e_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - ① $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - ② $x \equiv x'$ whenever there exists $e \in e_Y M e_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - 1 $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - 2 $x \equiv x'$ whenever there exists $e \in e_Y Me_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - 1 $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - 2 $x \equiv x'$ whenever there exists $e \in e_Y M e_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - 1 $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - 2 $x \equiv x'$ whenever there exists $e \in e_Y M e_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - 1 $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - 2 $x \equiv x'$ whenever there exists $e \in e_Y M e_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - 1 $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - 2 $x \equiv x'$ whenever there exists $e \in e_Y M e_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

The Case of Bands

- Suppose M is a band (i.e., all elements of M are idempotent).
- The vertex set of $Q(kM)$ is $\Lambda(kM)$.
- If MXM, MYM are incomparable or $X = Y$, there are no edges $X \rightarrow Y$.
- If $MXM \subsetneq MYM$, then we may suppose $e_X = e_Y e_X e_Y$.
- Let $L_X = \{m \in M \mid Mm = Me_X\}$.
- Let \equiv be the smallest equivalence relation on $e_Y L_X$ with:
 - 1 $e_Y m x \equiv e_Y m e_Y x$ for $x \in L_X$ and $MmM \supseteq MXM$;
 - 2 $x \equiv x'$ whenever there exists $e \in e_Y M e_Y \setminus \{e_Y\}$ with $MXM \subsetneq MeM$ and $ex = x, ex' = x'$.
- Then the number of edges $X \rightarrow Y$ is $|e_Y L_X / \equiv| - 1$.
- If $MYM \subsetneq MXM$, then the number of edges $X \rightarrow Y$ equals the number of edges $Y \rightarrow X$ in $Q(kM^{op})$.

Quiver of the Free Band

- There is a free band $FB(A)$ on any set A .
- J. A. Green and D. Rees solved the word problem for $FB(A)$.
- They proved A finite implies $FB(A)$ is finite.
- The quiver $Q(kFB(A))$ for A finite is as follows.
- The vertex set is 2^A .
- If $Y \subsetneq X$, then there are $|X \setminus Y| - 1$ edges $X \rightarrow Y$.
- If $Y \supsetneq X$, then there are $|Y \setminus X| - 1$ edges from $X \rightarrow Y$.
- There are no other edges.

Quiver of the Free Band

- There is a free band $FB(A)$ on any set A .
- J. A. Green and D. Rees solved the word problem for $FB(A)$.
- They proved A finite implies $FB(A)$ is finite.
- The quiver $Q(kFB(A))$ for A finite is as follows.
- The vertex set is 2^A .
- If $Y \subsetneq X$, then there are $|X \setminus Y| - 1$ edges $X \rightarrow Y$.
- If $Y \supsetneq X$, then there are $|Y \setminus X| - 1$ edges from $X \rightarrow Y$.
- There are no other edges.

Quiver of the Free Band

- There is a free band $FB(A)$ on any set A .
- J. A. Green and D. Rees solved the word problem for $FB(A)$.
- They proved A finite implies $FB(A)$ is finite.
- The quiver $Q(kFB(A))$ for A finite is as follows.
- The vertex set is 2^A .
- If $Y \subsetneq X$, then there are $|X \setminus Y| - 1$ edges $X \rightarrow Y$.
- If $Y \supsetneq X$, then there are $|Y \setminus X| - 1$ edges from $X \rightarrow Y$.
- There are no other edges.

Quiver of the Free Band

- There is a free band $FB(A)$ on any set A .
- J. A. Green and D. Rees solved the word problem for $FB(A)$.
- They proved A finite implies $FB(A)$ is finite.
- The quiver $Q(kFB(A))$ for A finite is as follows.
- The vertex set is 2^A .
- If $Y \subsetneq X$, then there are $|X \setminus Y| - 1$ edges $X \rightarrow Y$.
- If $Y \supsetneq X$, then there are $|Y \setminus X| - 1$ edges from $X \rightarrow Y$.
- There are no other edges.

Quiver of the Free Band

- There is a free band $FB(A)$ on any set A .
- J. A. Green and D. Rees solved the word problem for $FB(A)$.
- They proved A finite implies $FB(A)$ is finite.
- The quiver $Q(kFB(A))$ for A finite is as follows.
- The vertex set is 2^A .
- If $Y \subsetneq X$, then there are $|X \setminus Y| - 1$ edges $X \rightarrow Y$.
- If $Y \supsetneq X$, then there are $|Y \setminus X| - 1$ edges from $X \rightarrow Y$.
- There are no other edges.

Quiver of the Free Band

- There is a free band $FB(A)$ on any set A .
- J. A. Green and D. Rees solved the word problem for $FB(A)$.
- They proved A finite implies $FB(A)$ is finite.
- The quiver $Q(kFB(A))$ for A finite is as follows.
- The vertex set is 2^A .
- If $Y \subsetneq X$, then there are $|X \setminus Y| - 1$ edges $X \rightarrow Y$.
- If $Y \supsetneq X$, then there are $|Y \setminus X| - 1$ edges from $X \rightarrow Y$.
- There are no other edges.

Quiver of the Free Band

- There is a free band $FB(A)$ on any set A .
- J. A. Green and D. Rees solved the word problem for $FB(A)$.
- They proved A finite implies $FB(A)$ is finite.
- The quiver $Q(kFB(A))$ for A finite is as follows.
- The vertex set is 2^A .
- If $Y \subsetneq X$, then there are $|X \setminus Y| - 1$ edges $X \rightarrow Y$.
- If $Y \supsetneq X$, then there are $|Y \setminus X| - 1$ edges from $X \rightarrow Y$.
- There are no other edges.

Quiver of the Free Band

- There is a free band $FB(A)$ on any set A .
- J. A. Green and D. Rees solved the word problem for $FB(A)$.
- They proved A finite implies $FB(A)$ is finite.
- The quiver $Q(kFB(A))$ for A finite is as follows.
- The vertex set is 2^A .
- If $Y \subsetneq X$, then there are $|X \setminus Y| - 1$ edges $X \rightarrow Y$.
- If $Y \supsetneq X$, then there are $|Y \setminus X| - 1$ edges from $X \rightarrow Y$.
- There are no other edges.

The End

THANK YOU FOR YOUR
ATTENTION!