

Synchronizing groups: a survey

João Araújo, Michael Brough, Peter Cameron, Ian Gent,
Cristy Kazanidis, Tom Kelsey, Peter Neumann, Colva
Roney-Dougal, Nik Ruskuc, Jan Saxl, Csaba Schneider,
Pablo Spiga and **Benjamin Steinberg**



AMS Special Session on Algorithmic Problems of Group Theory
and Their Complexity
January 10, 2013

Outline

- 1 Background
- 2 Synchronizing groups
- 3 Combinatorics
- 4 Representation theory
- 5 Beyond synchronization

Synchronizing automata

An automaton is **synchronizing** if there is an input word which “resets” the automaton to a single state.

Synchronizing automata

An automaton is **synchronizing** if there is an input word which “resets” the automaton to a single state.

Example (Černý)




Figure : Unique minimum length reset word is $b(a^3b)^2$.

The Černý Conjecture

- The obvious generalization to n states has unique minimum length reset word $b(a^{n-1}b)^{n-2}$, which has length $(n-1)^2$.

The Černý Conjecture

- The obvious generalization to n states has unique minimum length reset word $b(a^{n-1}b)^{n-2}$, which has length $(n-1)^2$.
- The following conjecture has been open for 45 years and has been the subject of over 100 papers.

The Černý Conjecture

- The obvious generalization to n states has unique minimum length reset word $b(a^{n-1}b)^{n-2}$, which has length $(n-1)^2$.
- The following conjecture has been open for 45 years and has been the subject of over 100 papers.

Conjecture (Černý)

An n -state synchronizing automaton has a reset word of length at most $(n-1)^2$.

The Černý Conjecture

- The obvious generalization to n states has unique minimum length reset word $b(a^{n-1}b)^{n-2}$, which has length $(n-1)^2$.
- The following conjecture has been open for 45 years and has been the subject of over 100 papers.

Conjecture (Černý)

An n -state synchronizing automaton has a reset word of length at most $(n-1)^2$.

- The best upper bound to date is $\frac{n^3 - n}{6}$ (Pin/Frankl '81).

Pin's Theorem

- The first nontrivial result in this area is due to Pin.

Pin's Theorem

- The first nontrivial result in this area is due to Pin.

Theorem (Pin '78)

Let \mathcal{A} have a prime number of states and assume some letter a cyclically permutes the states.

Pin's Theorem

- The first nontrivial result in this area is due to Pin.

Theorem (Pin '78)

Let \mathcal{A} have a prime number of states and assume some letter a cyclically permutes the states.

- 1 *\mathcal{A} is synchronizing iff some input does not permute the states.*

Pin's Theorem

- The first nontrivial result in this area is due to Pin.

Theorem (Pin '78)

Let \mathcal{A} have a prime number of states and assume some letter a cyclically permutes the states.

- 1 *\mathcal{A} is synchronizing iff some input does not permute the states.*
- 2 *The Černý conjecture is true for such automata.*

Pin's Theorem

- The first nontrivial result in this area is due to Pin.

Theorem (Pin '78)

Let \mathcal{A} have a prime number of states and assume some letter a cyclically permutes the states.

- 1 *\mathcal{A} is synchronizing iff some input does not permute the states.*
- 2 *The Černý conjecture is true for such automata.*

- Dubuc removed the condition on prime number of states for (2) in 1998; (1) does not generalize.

Pin's Theorem

- The first nontrivial result in this area is due to Pin.

Theorem (Pin '78)

Let \mathcal{A} have a prime number of states and assume some letter a cyclically permutes the states.

- ① *\mathcal{A} is synchronizing iff some input does not permute the states.*
- ② *The Černý conjecture is true for such automata.*

- Dubuc removed the condition on prime number of states for (2) in 1998; (1) does not generalize.
- Inspired by (1), I introduced (2005) (and indep. Araújo [unpublished]) the notion of a **synchronizing group**.

Synchronization and monoids

- An automaton with state set Ω can be viewed as a collection A of self-maps of Ω .


Synchronization and monoids

- An automaton with state set Ω can be viewed as a collection A of self-maps of Ω .
- The mapping associated to a sends a state q to the endpoint of the edge from a labeled by q .

Synchronization and monoids

- An automaton with state set Ω can be viewed as a collection A of self-maps of Ω .
- The mapping associated to a sends a state q to the endpoint of the edge from a labeled by q .
- The automaton is synchronizing iff $\langle A \rangle$ contains a constant map.

Černý's example



$$a = \begin{pmatrix} 0 & 1 & 2 & 3 \\ 1 & 2 & 3 & 0 \end{pmatrix}$$

$$b = \begin{pmatrix} 0 & 1 & 2 & 3 \\ 1 & 1 & 2 & 3 \end{pmatrix}$$

The setup

- Ω is a finite set.

The setup

- Ω is a finite set.
- $G \leq S_\Omega$ is a permutation group acting on the right.

The setup

- Ω is a finite set.
- $G \leq S_\Omega$ is a permutation group acting on the right.
- T_Ω is the monoid of all self-maps of Ω .

The setup

- Ω is a finite set.
- $G \leq S_\Omega$ is a permutation group acting on the right.
- T_Ω is the monoid of all self-maps of Ω .
- The **rank** of $f \in T_\Omega$ is $|\Omega f|$.

The setup

- Ω is a finite set.
- $G \leq S_\Omega$ is a permutation group acting on the right.
- T_Ω is the monoid of all self-maps of Ω .
- The **rank** of $f \in T_\Omega$ is $|\Omega f|$.
- So f has rank 1 iff f is a constant map.

The setup

- Ω is a finite set.
- $G \leq S_\Omega$ is a permutation group acting on the right.
- T_Ω is the monoid of all self-maps of Ω .
- The **rank** of $f \in T_\Omega$ is $|\Omega f|$.
- So f has rank 1 iff f is a constant map.
- f is **singular** if $f \in T_\Omega \setminus S_\Omega$.

Definition of synchronizing groups

Definition (BS (2005), Araújo (unpublished))

G is **synchronizing** if $\langle G, f \rangle$ contains a constant map for each singular map $f \in T_\Omega$.

Definition of synchronizing groups

Definition (BS (2005), Araújo (unpublished))

G is **synchronizing** if $\langle G, f \rangle$ contains a constant map for each singular map $f \in T_\Omega$.

- Note if $G \leq H \leq S_\Omega$ and G is synchronizing, then so is H .

Definition of synchronizing groups

Definition (BS (2005), Araújo (unpublished))

G is **synchronizing** if $\langle G, f \rangle$ contains a constant map for each singular map $f \in T_\Omega$.

- Note if $G \leq H \leq S_\Omega$ and G is synchronizing, then so is H .
- Pin's theorem says the regular action of $\mathbb{Z}/p\mathbb{Z}$ is synchronizing.

Definition of synchronizing groups

Definition (BS (2005), Araújo (unpublished))

G is **synchronizing** if $\langle G, f \rangle$ contains a constant map for each singular map $f \in T_\Omega$.

- Note if $G \leq H \leq S_\Omega$ and G is synchronizing, then so is H .
- Pin's theorem says the regular action of $\mathbb{Z}/p\mathbb{Z}$ is synchronizing.
- It is easy to see that 2-transitive and 2-homogeneous groups are synchronizing.

Definition of synchronizing groups

Definition (BS (2005), Araújo (unpublished))

G is **synchronizing** if $\langle G, f \rangle$ contains a constant map for each singular map $f \in T_\Omega$.

- Note if $G \leq H \leq S_\Omega$ and G is synchronizing, then so is H .
- Pin's theorem says the regular action of $\mathbb{Z}/p\mathbb{Z}$ is synchronizing.
- It is easy to see that 2-transitive and 2-homogeneous groups are synchronizing.
- Synchronizing groups are primitive.

Definition of synchronizing groups

Definition (BS (2005), Araújo (unpublished))

G is **synchronizing** if $\langle G, f \rangle$ contains a constant map for each singular map $f \in T_\Omega$.

- Note if $G \leq H \leq S_\Omega$ and G is synchronizing, then so is H .
- Pin's theorem says the regular action of $\mathbb{Z}/p\mathbb{Z}$ is synchronizing.
- It is easy to see that 2-transitive and 2-homogeneous groups are synchronizing.
- Synchronizing groups are primitive.
- G is **primitive** if Ω admits no nontrivial G -invariant partition.

A semigroup-free reformulation

- Araújo came up with a semigroup-free reformulation that attracted the interest of first P. Neumann and then others.

A semigroup-free reformulation

- Araújo came up with a semigroup-free reformulation that attracted the interest of first P. Neumann and then others.
- A partition π on Ω is **section-regular** if there is a section S of π such that Sg is a section for all $g \in G$.

A semigroup-free reformulation

- Araújo came up with a semigroup-free reformulation that attracted the interest of first P. Neumann and then others.
- A partition π on Ω is **section-regular** if there is a section S of π such that Sg is a section for all $g \in G$.
- If π is G -invariant, then it is section-regular where S can be any section.

A semigroup-free reformulation

- Araújo came up with a semigroup-free reformulation that attracted the interest of first P. Neumann and then others.
- A partition π on Ω is **section-regular** if there is a section S of π such that Sg is a section for all $g \in G$.
- If π is G -invariant, then it is section-regular where S can be any section.

Proposition (Araújo '06)

G is synchronizing iff Ω admits no nontrivial section-regular partition.

A semigroup-free reformulation

- Araújo came up with a semigroup-free reformulation that attracted the interest of first P. Neumann and then others.
- A partition π on Ω is **section-regular** if there is a section S of π such that Sg is a section for all $g \in G$.
- If π is G -invariant, then it is section-regular where S can be any section.

Proposition (Araújo '06)

G is synchronizing iff Ω admits no nontrivial section-regular partition.

Proof.

Section-regular partitions essentially correspond to minimal rank idempotents that can't be synchronized. □

Synchronizing groups are basic

- Let $\Omega = \Delta^n$.

Synchronizing groups are basic

- Let $\Omega = \Delta^n$.
- $S_\Delta \wr S_n = S_\Delta^n \rtimes S_n$ acts primitively on Ω .

Synchronizing groups are basic

- Let $\Omega = \Delta^n$.
- $S_\Delta \wr S_n = S_\Delta^n \rtimes S_n$ acts primitively on Ω .
- S_n permutes the coordinates and S_Δ^n acts coordinate-wise.

Synchronizing groups are basic

- Let $\Omega = \Delta^n$.
- $S_\Delta \wr S_n = S_\Delta^n \rtimes S_n$ acts primitively on Ω .
- S_n permutes the coordinates and S_Δ^n acts coordinate-wise.
- The partition $(\delta_1, \dots, \delta_n) \sim (\delta'_1, \dots, \delta'_n)$ iff $\delta_1 = \delta'_1$ is section-regular.

Synchronizing groups are basic

- Let $\Omega = \Delta^n$.
- $S_\Delta \wr S_n = S_\Delta^n \rtimes S_n$ acts primitively on Ω .
- S_n permutes the coordinates and S_Δ^n acts coordinate-wise.
- The partition $(\delta_1, \dots, \delta_n) \sim (\delta'_1, \dots, \delta'_n)$ iff $\delta_1 = \delta'_1$ is section-regular.
- The section is the set of constant vectors.

Synchronizing groups are basic

- Let $\Omega = \Delta^n$.
- $S_\Delta \wr S_n = S_\Delta^n \rtimes S_n$ acts primitively on Ω .
- S_n permutes the coordinates and S_Δ^n acts coordinate-wise.
- The partition $(\delta_1, \dots, \delta_n) \sim (\delta'_1, \dots, \delta'_n)$ iff $\delta_1 = \delta'_1$ is section-regular.
- The section is the set of constant vectors.
- Thus synchronizing groups are basic in the O'Nan-Scott classification scheme.

Synchronizing groups are basic

- Let $\Omega = \Delta^n$.
- $S_\Delta \wr S_n = S_\Delta^n \rtimes S_n$ acts primitively on Ω .
- S_n permutes the coordinates and S_Δ^n acts coordinate-wise.
- The partition $(\delta_1, \dots, \delta_n) \sim (\delta'_1, \dots, \delta'_n)$ iff $\delta_1 = \delta'_1$ is section-regular.
- The section is the set of constant vectors.
- Thus synchronizing groups are basic in the O'Nan-Scott classification scheme.
- So they are of affine, diagonal or almost simple type.

Synchronizing groups are basic

- Let $\Omega = \Delta^n$.
- $S_\Delta \wr S_n = S_\Delta^n \rtimes S_n$ acts primitively on Ω .
- S_n permutes the coordinates and S_Δ^n acts coordinate-wise.
- The partition $(\delta_1, \dots, \delta_n) \sim (\delta'_1, \dots, \delta'_n)$ iff $\delta_1 = \delta'_1$ is section-regular.
- The section is the set of constant vectors.
- Thus synchronizing groups are basic in the O'Nan-Scott classification scheme.
- So they are of affine, diagonal or almost simple type.

Theorem

$2\text{-transitive} \implies 2\text{-homogeneous} \implies \text{synchronizing} \implies$
 $\text{basic} \implies \text{primitive} \implies \text{transitive}.$

Some results of Peter Neumann

The following results are due to Peter Neumann.

- If π is a section-regular partition, then each block of π has the same size.

Some results of Peter Neumann

The following results are due to Peter Neumann.

- If π is a section-regular partition, then each block of π has the same size.
- This gives Pin's theorem that any primitive group of prime degree is synchronizing.

Some results of Peter Neumann

The following results are due to Peter Neumann.

- If π is a section-regular partition, then each block of π has the same size.
- This gives Pin's theorem that any primitive group of prime degree is synchronizing.
- If r is the size of each part and s is the number of blocks, then $r, s > 2$.

Some results of Peter Neumann

The following results are due to Peter Neumann.

- If π is a section-regular partition, then each block of π has the same size.
- This gives Pin's theorem that any primitive group of prime degree is synchronizing.
- If r is the size of each part and s is the number of blocks, then $r, s > 2$.
- Thus every primitive group of size $2p$ with p an odd prime is synchronizing.

Some results of Peter Neumann

The following results are due to Peter Neumann.

- If π is a section-regular partition, then each block of π has the same size.
- This gives Pin's theorem that any primitive group of prime degree is synchronizing.
- If r is the size of each part and s is the number of blocks, then $r, s > 2$.
- Thus every primitive group of size $2p$ with p an odd prime is synchronizing.
- Synchronizing groups have a fairly large density among primitive permutation groups using CFSG.

Connections with graph theory

- A **graph** X on Ω is a simple graph with vertex set Ω .

Connections with graph theory

- A **graph** X on Ω is a simple graph with vertex set Ω .
- X is **trivial** if it is complete or edgeless.

Connections with graph theory

- A **graph** X on Ω is a simple graph with vertex set Ω .
- X is **trivial** if it is complete or edgeless.
- X is **G -invariant** if $G \leq \text{Aut}(X)$.

Connections with graph theory

- A **graph** X on Ω is a simple graph with vertex set Ω .
- X is **trivial** if it is complete or edgeless.
- X is **G -invariant** if $G \leq \text{Aut}(X)$.
- The **clique number** $\omega(X)$ is the size of the largest complete subgraph of X .

Connections with graph theory

- A **graph** X on Ω is a simple graph with vertex set Ω .
- X is **trivial** if it is complete or edgeless.
- X is **G -invariant** if $G \leq \text{Aut}(X)$.
- The **clique number** $\omega(X)$ is the size of the largest complete subgraph of X .
- The **chromatic number** $\chi(X)$ is the minimum size of a coloring of X .

Connections with graph theory

- A **graph** X on Ω is a simple graph with vertex set Ω .
- X is **trivial** if it is complete or edgeless.
- X is **G -invariant** if $G \leq \text{Aut}(X)$.
- The **clique number** $\omega(X)$ is the size of the largest complete subgraph of X .
- The **chromatic number** $\chi(X)$ is the minimum size of a coloring of X .
- $\omega(X) \leq \chi(X)$.

Connections with graph theory

- A **graph** X on Ω is a simple graph with vertex set Ω .
- X is **trivial** if it is complete or edgeless.
- X is **G -invariant** if $G \leq \text{Aut}(X)$.
- The **clique number** $\omega(X)$ is the size of the largest complete subgraph of X .
- The **chromatic number** $\chi(X)$ is the minimum size of a coloring of X .
- $\omega(X) \leq \chi(X)$.

Theorem (Cameron)

G is synchronizing iff there are no nontrivial G -invariant graphs on Ω with $\omega(X) = \chi(X)$.

Action of S_n on k -sets

- The action of S_n on k -sets is primitive if $n \geq 2k + 1$.

Action of S_n on k -sets

- The action of S_n on k -sets is primitive if $n \geq 2k + 1$.
- For the case $k = 2$, there are exactly two S_n -invariant graphs.

Action of S_n on k -sets

- The action of S_n on k -sets is primitive if $n \geq 2k + 1$.
- For the case $k = 2$, there are exactly two S_n -invariant graphs.
- Using Cameron's theorem, one has the following result.

Action of S_n on k -sets

- The action of S_n on k -sets is primitive if $n \geq 2k + 1$.
- For the case $k = 2$, there are exactly two S_n -invariant graphs.
- Using Cameron's theorem, one has the following result.

Theorem (Neumann)

For $n \geq 5$, S_n acting on 2-sets is synchronizing iff n is odd.

Action of S_n on k -sets

- The action of S_n on k -sets is primitive if $n \geq 2k + 1$.
- For the case $k = 2$, there are exactly two S_n -invariant graphs.
- Using Cameron's theorem, one has the following result.

Theorem (Neumann)

For $n \geq 5$, S_n acting on 2-sets is synchronizing iff n is odd.

Using combinatorics like Steiner systems and the Erdős-Ko-Rado theorem leads to the following result.

Action of S_n on k -sets

- The action of S_n on k -sets is primitive if $n \geq 2k + 1$.
- For the case $k = 2$, there are exactly two S_n -invariant graphs.
- Using Cameron's theorem, one has the following result.

Theorem (Neumann)

For $n \geq 5$, S_n acting on 2-sets is synchronizing iff n is odd.

Using combinatorics like Steiner systems and the Erdős-Ko-Rado theorem leads to the following result.

Theorem

- 1 If $k \mid n$, then S_n acting on k -sets is not synchronizing.

Action of S_n on k -sets

- The action of S_n on k -sets is primitive if $n \geq 2k + 1$.
- For the case $k = 2$, there are exactly two S_n -invariant graphs.
- Using Cameron's theorem, one has the following result.

Theorem (Neumann)

For $n \geq 5$, S_n acting on 2-sets is synchronizing iff n is odd.

Using combinatorics like Steiner systems and the Erdős-Ko-Rado theorem leads to the following result.

Theorem

- 1 If $k \mid n$, then S_n acting on k -sets is not synchronizing.
- 2 If $n \geq 7$, then S_n acting on 3-sets is synchronizing iff $n \equiv 2, 4, 5 \pmod{6}$ and $n \neq 8$.

Classical groups

- By classical groups, I mean symplectic, unitary and orthogonal groups over finite fields.

Classical groups

- By classical groups, I mean symplectic, unitary and orthogonal groups over finite fields.
- I will be slightly sloppy here on technical details to avoid trivial cases.

Classical groups

- By classical groups, I mean symplectic, unitary and orthogonal groups over finite fields.
- I will be slightly sloppy here on technical details to avoid trivial cases.
- A classical group acts on a vector space V preserving a form.

Classical groups

- By classical groups, I mean symplectic, unitary and orthogonal groups over finite fields.
- I will be slightly sloppy here on technical details to avoid trivial cases.
- A classical group acts on a vector space V preserving a form.
- A subspace is **totally isotropic** if the form vanishes on it.

Classical groups

- By classical groups, I mean symplectic, unitary and orthogonal groups over finite fields.
- I will be slightly sloppy here on technical details to avoid trivial cases.
- A classical group acts on a vector space V preserving a form.
- A subspace is **totally isotropic** if the form vanishes on it.
- The points of the **polar space** Ω associated to the classical group are the 1-dimensional totally isotropic subspaces.

Classical groups

- By classical groups, I mean symplectic, unitary and orthogonal groups over finite fields.
- I will be slightly sloppy here on technical details to avoid trivial cases.
- A classical group acts on a vector space V preserving a form.
- A subspace is **totally isotropic** if the form vanishes on it.
- The points of the **polar space** Ω associated to the classical group are the 1-dimensional totally isotropic subspaces.
- Two points are **collinear** if they span a 2-dimensional totally isotropic subspace.

Classical groups

- By classical groups, I mean symplectic, unitary and orthogonal groups over finite fields.
- I will be slightly sloppy here on technical details to avoid trivial cases.
- A classical group acts on a vector space V preserving a form.
- A subspace is **totally isotropic** if the form vanishes on it.
- The points of the **polar space** Ω associated to the classical group are the 1-dimensional totally isotropic subspaces.
- Two points are **collinear** if they span a 2-dimensional totally isotropic subspace.
- The graph X of the polar space is the graph on Ω with edges the collinear points.

Synchronization and finite geometries

- The projective quotient G of the classical group acts primitively on the polar space Ω .

Synchronization and finite geometries

- The projective quotient G of the classical group acts primitively on the polar space Ω .
- A consequence of Witt's lemma is that X and \overline{X} are the only G -invariant graphs.

Synchronization and finite geometries

- The projective quotient G of the classical group acts primitively on the polar space Ω .
- A consequence of Witt's lemma is that X and \overline{X} are the only G -invariant graphs.

Theorem

A classical group acting on its polar space is synchronizing iff its polar space possesses neither:

- ① *both an ovoid and a spread; nor*
- ② *a partition into ovoids.*

Synchronization and finite geometries

- The projective quotient G of the classical group acts primitively on the polar space Ω .
- A consequence of Witt's lemma is that X and \overline{X} are the only G -invariant graphs.

Theorem

A classical group acting on its polar space is synchronizing iff its polar space possesses neither:

- ① *both an ovoid and a spread; nor*
- ② *a partition into ovoids.*

- Ovoids and spreads are notions from finite geometry that I cannot/will not define.

Synchronization and finite geometries

- The projective quotient G of the classical group acts primitively on the polar space Ω .
- A consequence of Witt's lemma is that X and \overline{X} are the only G -invariant graphs.

Theorem

A classical group acting on its polar space is synchronizing iff its polar space possesses neither:

- ① *both an ovoid and a spread; nor*
- ② *a partition into ovoids.*

- Ovoids and spreads are notions from finite geometry that I cannot/will not define.
- Not everything is known about the existence of these creatures.

Some examples

Example

- $\mathrm{PSp}(2r, q)$, $\mathrm{PSU}(2r + 1, q)$ and $\mathrm{P}\Omega^-(2r + 2, q)$ are synchronizing for all $r \geq 2$, except $\mathrm{PSp}(4, q)$ with q even.
- $\mathrm{P}\Omega(5, q)$ for $q = 3, 5, 7$ is synchronizing.

Some examples

Example

- $\mathrm{PSp}(2r, q)$, $\mathrm{PSU}(2r + 1, q)$ and $\mathrm{P}\Omega^-(2r + 2, q)$ are synchronizing for all $r \geq 2$, except $\mathrm{PSp}(4, q)$ with q even.
- $\mathrm{P}\Omega(5, q)$ for $q = 3, 5, 7$ is synchronizing.

The general situation for classical groups is still very much open.

$\mathbb{k}I$ -groups

- Let \mathbb{k} be a field of characteristic 0.

$\mathbb{k}I$ -groups

- Let \mathbb{k} be a field of characteristic 0.
- $\mathbb{k}\Omega$ has 2 obvious $\mathbb{k}G$ -submodules:

$\mathbb{k}I$ -groups

- Let \mathbb{k} be a field of characteristic 0.
- $\mathbb{k}\Omega$ has 2 obvious $\mathbb{k}G$ -submodules:
 - the trivial submodule spanned by $\sum_{\omega \in \Omega} \omega$; and

$\mathbb{k}I$ -groups

- Let \mathbb{k} be a field of characteristic 0.
- $\mathbb{k}\Omega$ has 2 obvious $\mathbb{k}G$ -submodules:
 - the trivial submodule spanned by $\sum_{\omega \in \Omega} \omega$; and
 - its orthogonal complement V .

$\mathbb{k}I$ -groups

- Let \mathbb{k} be a field of characteristic 0.
- $\mathbb{k}\Omega$ has 2 obvious $\mathbb{k}G$ -submodules:
 - the trivial submodule spanned by $\sum_{\omega \in \Omega} \omega$; and
 - its orthogonal complement V .
- Call G a $\mathbb{k}I$ -group if V is irreducible.

$\mathbb{k}I$ -groups

- Let \mathbb{k} be a field of characteristic 0.
- $\mathbb{k}\Omega$ has 2 obvious $\mathbb{k}G$ -submodules:
 - the trivial submodule spanned by $\sum_{\omega \in \Omega} \omega$; and
 - its orthogonal complement V .
- Call G a $\mathbb{k}I$ -group if V is irreducible.
- If $\mathbb{F} \leq \mathbb{k}$, then $\mathbb{k}I \implies \mathbb{F}I$.

$\mathbb{k}I$ -groups

- Let \mathbb{k} be a field of characteristic 0.
- $\mathbb{k}\Omega$ has 2 obvious $\mathbb{k}G$ -submodules:
 - the trivial submodule spanned by $\sum_{\omega \in \Omega} \omega$; and
 - its orthogonal complement V .
- Call G a $\mathbb{k}I$ -group if V is irreducible.
- If $\mathbb{F} \leq \mathbb{k}$, then $\mathbb{k}I \implies \mathbb{F}I$.

Theorem (Classical)

- 1 G is $\mathbb{C}I \iff G$ is 2-transitive.
- 2 G is $\mathbb{R}I \iff G$ is 2-homogeneous.

$\mathbb{Q}I$ -group

Theorem (Arnold, BS '06)

A $\mathbb{Q}I$ -group is synchronizing.

- Any primitive group of prime degree is $\mathbb{Q}I$, but most are not 2-homogeneous ($=\mathbb{R}I$).

$\mathbb{Q}I$ -group

Theorem (Arnold, BS '06)

A $\mathbb{Q}I$ -group is synchronizing.

- Any primitive group of prime degree is $\mathbb{Q}I$, but most are not 2-homogeneous ($=\mathbb{R}I$).
- I asked John Dixon what is known about such groups.

$\mathbb{Q}I$ -group

Theorem (Arnold, BS '06)

A $\mathbb{Q}I$ -group is synchronizing.

- Any primitive group of prime degree is $\mathbb{Q}I$, but most are not 2-homogeneous ($=\mathbb{R}I$).
- I asked John Dixon what is known about such groups.
- He observed they are 3/2-transitive.

$\mathbb{Q}I$ -group

Theorem (Arnold, BS '06)

A $\mathbb{Q}I$ -group is synchronizing.

- Any primitive group of prime degree is $\mathbb{Q}I$, but most are not 2-homogeneous ($=\mathbb{R}I$).
- I asked John Dixon what is known about such groups.
- He observed they are 3/2-transitive.
- He proved that an affine group G acting on \mathbb{F}_q^n is synchronizing iff the group generated by G and \mathbb{F}_q^* is 2-transitive on \mathbb{F}_q^n .

$\mathbb{Q}I$ -group

Theorem (Arnold, BS '06)

A $\mathbb{Q}I$ -group is synchronizing.

- Any primitive group of prime degree is $\mathbb{Q}I$, but most are not 2-homogeneous ($=\mathbb{R}I$).
- I asked John Dixon what is known about such groups.
- He observed they are 3/2-transitive.
- He proved that an affine group G acting on \mathbb{F}_q^n is synchronizing iff the group generated by G and \mathbb{F}_q^* is 2-transitive on \mathbb{F}_q^n .
- Using CFSG, it has been shown that the only remaining examples are $\text{PSL}(2, q)$ and $\text{P}\Gamma\text{L}(2, q)$ acting in degree $\frac{1}{2}q(q-1)$ with $q = 2^n \geq 8$ and $q-1$ a Mersenne prime.

Summary

So far we have the following strict hierarchy of permutation groups:

Summary

So far we have the following strict hierarchy of permutation groups:

- 1 2-transitive groups

Summary

So far we have the following strict hierarchy of permutation groups:

- 1 2-transitive groups
- 2 2-homogeneous groups

Summary

So far we have the following strict hierarchy of permutation groups:

- 1 2-transitive groups
- 2 2-homogeneous groups
- 3 QI -groups

Summary

So far we have the following strict hierarchy of permutation groups:

- 1 2-transitive groups
- 2 2-homogeneous groups
- 3 $\mathbb{Q}I$ -groups
- 4 synchronizing groups

Summary

So far we have the following strict hierarchy of permutation groups:

- 1 2-transitive groups
- 2 2-homogeneous groups
- 3 $\mathbb{Q}I$ -groups
- 4 synchronizing groups
- 5 basic groups

Summary

So far we have the following strict hierarchy of permutation groups:

- ① 2-transitive groups
- ② 2-homogeneous groups
- ③ $\mathbb{Q}I$ -groups
- ④ synchronizing groups
- ⑤ basic groups
- ⑥ primitive groups

Summary

So far we have the following strict hierarchy of permutation groups:

- 1 2-transitive groups
- 2 2-homogeneous groups
- 3 $\mathbb{Q}I$ -groups
- 4 synchronizing groups
- 5 basic groups
- 6 primitive groups
- 7 transitive groups

Summary

So far we have the following strict hierarchy of permutation groups:

- ① 2-transitive groups
- ② 2-homogeneous groups
- ③ $\mathbb{Q}I$ -groups
- ④ synchronizing groups
- ⑤ basic groups
- ⑥ primitive groups
- ⑦ transitive groups

Another class we have considered are [separating groups](#) (introduced by P. Neumann).

Neumann's separation lemma

Lemma (P. Neumann)

Let $G \leq S_\Omega$ be transitive and let $A, B \subseteq \Omega$ such that $|A| \cdot |B| < |\Omega|$. Then there exists $g \in G$ with $A \cap Bg = \emptyset$.

Neumann's separation lemma

Lemma (P. Neumann)

Let $G \leq S_\Omega$ be transitive and let $A, B \subseteq \Omega$ such that $|A| \cdot |B| < |\Omega|$. Then there exists $g \in G$ with $A \cap Bg = \emptyset$.

- In other words, the subsets A, B can be separated by G if they are not large enough.

Neumann's separation lemma

Lemma (P. Neumann)

Let $G \leq S_\Omega$ be transitive and let $A, B \subseteq \Omega$ such that $|A| \cdot |B| < |\Omega|$. Then there exists $g \in G$ with $A \cap Bg = \emptyset$.

- In other words, the subsets A, B can be separated by G if they are not large enough.
- A transitive group G is called **separating** if whenever $|A| \cdot |B| = |\Omega|$, there exists $g \in G$ with $A \cap Bg = \emptyset$.

Neumann's separation lemma

Lemma (P. Neumann)

Let $G \leq S_\Omega$ be transitive and let $A, B \subseteq \Omega$ such that $|A| \cdot |B| < |\Omega|$. Then there exists $g \in G$ with $A \cap Bg = \emptyset$.

- In other words, the subsets A, B can be separated by G if they are not large enough.
- A transitive group G is called **separating** if whenever $|A| \cdot |B| = |\Omega|$, there exists $g \in G$ with $A \cap Bg = \emptyset$.

Theorem (Neumann)

Separating groups are synchronizing.

Neumann's separation lemma

Lemma (P. Neumann)

Let $G \leq S_\Omega$ be transitive and let $A, B \subseteq \Omega$ such that $|A| \cdot |B| < |\Omega|$. Then there exists $g \in G$ with $A \cap Bg = \emptyset$.

- In other words, the subsets A, B can be separated by G if they are not large enough.
- A transitive group G is called **separating** if whenever $|A| \cdot |B| = |\Omega|$, there exists $g \in G$ with $A \cap Bg = \emptyset$.

Theorem (Neumann)

Separating groups are synchronizing.

Proof.

If π is a section-regular partition with section S and A is a block of π , then $|A| \cdot |S| = |\Omega|$ and $|A \cap Sg| = 1$ for all $g \in G$. \square

Connections with combinatorics

- The **independence number** $\alpha(X)$ of a graph X is the largest size of a coclique of X .

Connections with combinatorics

- The **independence number** $\alpha(X)$ of a graph X is the largest size of a coclique of X .

Theorem (Cameron)

G is separating iff there are no nontrivial G -invariant graphs X on Ω with $\alpha(X) \cdot \omega(X) = |\Omega|$.

Connections with combinatorics

- The **independence number** $\alpha(X)$ of a graph X is the largest size of a coclique of X .

Theorem (Cameron)

G is separating iff there are no nontrivial G -invariant graphs X on Ω with $\alpha(X) \cdot \omega(X) = |\Omega|$.

- The results for S_n acting on k -sets holds verbatim for separating groups.

Connections with combinatorics

- The **independence number** $\alpha(X)$ of a graph X is the largest size of a coclique of X .

Theorem (Cameron)

G is separating iff there are no nontrivial G -invariant graphs X on Ω with $\alpha(X) \cdot \omega(X) = |\Omega|$.

- The results for S_n acting on k -sets holds verbatim for separating groups.

Theorem

A classical group acting on its polar space is separating iff its polar space does not possess an ovoid.

Connections with combinatorics

- The **independence number** $\alpha(X)$ of a graph X is the largest size of a coclique of X .

Theorem (Cameron)

G is separating iff there are no nontrivial G -invariant graphs X on Ω with $\alpha(X) \cdot \omega(X) = |\Omega|$.

- The results for S_n acting on k -sets holds verbatim for separating groups.

Theorem

A classical group acting on its polar space is separating iff its polar space does not possess an ovoid.

- $P\Omega(5, q)$ for $q = 3, 5, 7$ is synchronizing but not separating.

Revised Summary

We have the following strict hierarchy of permutation groups:

Revised Summary

We have the following strict hierarchy of permutation groups:

- 1 2-transitive groups

Revised Summary

We have the following strict hierarchy of permutation groups:

- 1 2-transitive groups
- 2 2-homogeneous groups

Revised Summary

We have the following strict hierarchy of permutation groups:

- 1 2-transitive groups
- 2 2-homogeneous groups
- 3 $\mathbb{Q}I$ -groups

Revised Summary

We have the following strict hierarchy of permutation groups:

- ① 2-transitive groups
- ② 2-homogeneous groups
- ③ $\mathbb{Q}I$ -groups
- ④ separating groups

Revised Summary

We have the following strict hierarchy of permutation groups:

- ① 2-transitive groups
- ② 2-homogeneous groups
- ③ $\mathbb{Q}I$ -groups
- ④ separating groups
- ⑤ synchronizing groups

Revised Summary

We have the following strict hierarchy of permutation groups:

- ① 2-transitive groups
- ② 2-homogeneous groups
- ③ $\mathbb{Q}I$ -groups
- ④ separating groups
- ⑤ synchronizing groups
- ⑥ basic groups

Revised Summary

We have the following strict hierarchy of permutation groups:

- ① 2-transitive groups
- ② 2-homogeneous groups
- ③ $\mathbb{Q}I$ -groups
- ④ separating groups
- ⑤ synchronizing groups
- ⑥ basic groups
- ⑦ primitive groups

Revised Summary

We have the following strict hierarchy of permutation groups:

- ① 2-transitive groups
- ② 2-homogeneous groups
- ③ $\mathbb{Q}I$ -groups
- ④ separating groups
- ⑤ synchronizing groups
- ⑥ basic groups
- ⑦ primitive groups
- ⑧ transitive groups

The End

THANKS FOR YOUR
ATTENTION!