

# The submonoid membership problem for groups

Markus Lohrey<sup>1</sup> Benjamin Steinberg<sup>2</sup>

<sup>1</sup>Universität Leipzig

<sup>2</sup>City College of New York

[bsteinberg@ccny.cuny.edu](mailto:bsteinberg@ccny.cuny.edu)

<http://www.sci.ccny.cuny.edu/~benjamin/>

January 4, 2012

# Integer programming

- INTEGER PROGRAMMING:
  - Given  $A \in M_{mn}(\mathbb{Z})$  and  $\mathbf{b} \in \mathbb{Z}^m$ , does  $A\mathbf{x} = \mathbf{b}$  have a solution  $\mathbf{x} \in \mathbb{N}^n$ ?

# Integer programming

- INTEGER PROGRAMMING:
  - Given  $A \in M_{mn}(\mathbb{Z})$  and  $\mathbf{b} \in \mathbb{Z}^m$ , does  $A\mathbf{x} = \mathbf{b}$  have a solution  $\mathbf{x} \in \mathbb{N}^n$ ?
- Algebraically speaking, the problem is to determine whether  $\mathbf{b}$  is a non-negative linear combination of the columns of  $A$ .

# Integer programming

- INTEGER PROGRAMMING:
  - Given  $A \in M_{mn}(\mathbb{Z})$  and  $\mathbf{b} \in \mathbb{Z}^m$ , does  $A\mathbf{x} = \mathbf{b}$  have a solution  $\mathbf{x} \in \mathbb{N}^n$ ?
- Algebraically speaking, the problem is to determine whether  $\mathbf{b}$  is a non-negative linear combination of the columns of  $A$ .
- In other words, does  $\mathbf{b}$  belong to the submonoid of  $\mathbb{Z}^m$  generated by the columns of  $A$ ?

# Integer programming

- INTEGER PROGRAMMING:
  - Given  $A \in M_{mn}(\mathbb{Z})$  and  $\mathbf{b} \in \mathbb{Z}^m$ , does  $A\mathbf{x} = \mathbf{b}$  have a solution  $\mathbf{x} \in \mathbb{N}^n$ ?
- Algebraically speaking, the problem is to determine whether  $\mathbf{b}$  is a non-negative linear combination of the columns of  $A$ .
- In other words, does  $\mathbf{b}$  belong to the submonoid of  $\mathbb{Z}^m$  generated by the columns of  $A$ ?
- So integer programming is the submonoid membership problem for abelian groups.

# Integer programming

- **INTEGER PROGRAMMING:**
  - Given  $A \in M_{mn}(\mathbb{Z})$  and  $\mathbf{b} \in \mathbb{Z}^m$ , does  $A\mathbf{x} = \mathbf{b}$  have a solution  $\mathbf{x} \in \mathbb{N}^n$ ?
- Algebraically speaking, the problem is to determine whether  $\mathbf{b}$  is a non-negative linear combination of the columns of  $A$ .
- In other words, does  $\mathbf{b}$  belong to the submonoid of  $\mathbb{Z}^m$  generated by the columns of  $A$ ?
- So integer programming is the submonoid membership problem for abelian groups.
- Integer programming is well known to be NP-complete.

# Integer programming

- **INTEGER PROGRAMMING:**
  - Given  $A \in M_{mn}(\mathbb{Z})$  and  $\mathbf{b} \in \mathbb{Z}^m$ , does  $A\mathbf{x} = \mathbf{b}$  have a solution  $\mathbf{x} \in \mathbb{N}^n$ ?
- Algebraically speaking, the problem is to determine whether  $\mathbf{b}$  is a non-negative linear combination of the columns of  $A$ .
- In other words, does  $\mathbf{b}$  belong to the submonoid of  $\mathbb{Z}^m$  generated by the columns of  $A$ ?
- So integer programming is the submonoid membership problem for abelian groups.
- Integer programming is well known to be NP-complete.
- The submonoid membership problem for arbitrary groups is a non-commutative analogue of integer programming.

## Some decision problems

- Fix a group  $G$  and a finite symmetric generating set  $\Sigma$ .



## Some decision problems

- Fix a group  $G$  and a finite symmetric generating set  $\Sigma$ .
- Let  $\pi: \Sigma^* \rightarrow G$  be the canonical projection.

## Some decision problems

- Fix a group  $G$  and a finite symmetric generating set  $\Sigma$ .
- Let  $\pi: \Sigma^* \rightarrow G$  be the canonical projection.
- Consider the following algorithmic problems for  $G$ .

## Some decision problems

- Fix a group  $G$  and a finite symmetric generating set  $\Sigma$ .
- Let  $\pi: \Sigma^* \rightarrow G$  be the canonical projection.
- Consider the following algorithmic problems for  $G$ .
- THE WORD PROBLEM:
  - Given  $w \in \Sigma^*$ , does  $\pi(w) = 1$ ?

## Some decision problems

- Fix a group  $G$  and a finite symmetric generating set  $\Sigma$ .
- Let  $\pi: \Sigma^* \rightarrow G$  be the canonical projection.
- Consider the following algorithmic problems for  $G$ .
- THE WORD PROBLEM:
  - Given  $w \in \Sigma^*$ , does  $\pi(w) = 1$ ?
- THE (UNIFORM) GENERALIZED WORD PROBLEM:
  - Given  $w, w_1, \dots, w_n \in \Sigma^*$ , is  $\pi(w) \in \langle \pi(w_1), \dots, \pi(w_n) \rangle$ ?

## Some decision problems

- Fix a group  $G$  and a finite symmetric generating set  $\Sigma$ .
- Let  $\pi: \Sigma^* \rightarrow G$  be the canonical projection.
- Consider the following algorithmic problems for  $G$ .
- THE WORD PROBLEM:
  - Given  $w \in \Sigma^*$ , does  $\pi(w) = 1$ ?
- THE (UNIFORM) GENERALIZED WORD PROBLEM:
  - Given  $w, w_1, \dots, w_n \in \Sigma^*$ , is  $\pi(w) \in \langle \pi(w_1), \dots, \pi(w_n) \rangle$ ?
- THE (UNIFORM) SUBMONOID MEMBERSHIP PROBLEM:
  - Given  $w, w_1, \dots, w_n \in \Sigma^*$ , is  $\pi(w) \in \pi(\{w_1, \dots, w_n\}^*)$ ?

## Some decision problems

- Fix a group  $G$  and a finite symmetric generating set  $\Sigma$ .
- Let  $\pi: \Sigma^* \rightarrow G$  be the canonical projection.
- Consider the following algorithmic problems for  $G$ .
- **THE WORD PROBLEM:**
  - Given  $w \in \Sigma^*$ , does  $\pi(w) = 1$ ?
- **THE (UNIFORM) GENERALIZED WORD PROBLEM:**
  - Given  $w, w_1, \dots, w_n \in \Sigma^*$ , is  $\pi(w) \in \langle \pi(w_1), \dots, \pi(w_n) \rangle$ ?
- **THE (UNIFORM) SUBMONOID MEMBERSHIP PROBLEM:**
  - Given  $w, w_1, \dots, w_n \in \Sigma^*$ , is  $\pi(w) \in \pi(\{w_1, \dots, w_n\}^*)$ ?
- **THE (UNIFORM) RATIONAL SUBSET MEMBERSHIP PROBLEM:**
  - Given  $w \in \Sigma^*$  and a finite automaton  $\mathcal{A}$  over  $\Sigma$ , is  $\pi(w) \in \pi(L(\mathcal{A}))$ ?

## Some decision problems

- Fix a group  $G$  and a finite symmetric generating set  $\Sigma$ .
- Let  $\pi: \Sigma^* \rightarrow G$  be the canonical projection.
- Consider the following algorithmic problems for  $G$ .
- **THE WORD PROBLEM:**
  - Given  $w \in \Sigma^*$ , does  $\pi(w) = 1$ ?
- **THE (UNIFORM) GENERALIZED WORD PROBLEM:**
  - Given  $w, w_1, \dots, w_n \in \Sigma^*$ , is  $\pi(w) \in \langle \pi(w_1), \dots, \pi(w_n) \rangle$ ?
- **THE (UNIFORM) SUBMONOID MEMBERSHIP PROBLEM:**
  - Given  $w, w_1, \dots, w_n \in \Sigma^*$ , is  $\pi(w) \in \pi(\{w_1, \dots, w_n\}^*)$ ?
- **THE (UNIFORM) RATIONAL SUBSET MEMBERSHIP PROBLEM:**
  - Given  $w \in \Sigma^*$  and a finite automaton  $\mathcal{A}$  over  $\Sigma$ , is  $\pi(w) \in \pi(L(\mathcal{A}))$ ?
- Decidability of these problems is independent of  $\Sigma$ .

# The generalized word problem

- The above decision problems were listed in order of difficulty.



# The generalized word problem

- The above decision problems were listed in order of difficulty.
- It is natural to search for groups distinguishing these problems.

# The generalized word problem

- The above decision problems were listed in order of difficulty.
- It is natural to search for groups distinguishing these problems.
- $F_2 \times F_2$  has undecidable generalized word problem (Mihailova)

# The generalized word problem

- The above decision problems were listed in order of difficulty.
- It is natural to search for groups distinguishing these problems.
- $F_2 \times F_2$  has undecidable generalized word problem (Mihailova)
- Free solvable groups of derived length  $\geq 3$  and rank  $\geq 2$  have undecidable generalized word problem (Umirbaev).

# The generalized word problem

- The above decision problems were listed in order of difficulty.
- It is natural to search for groups distinguishing these problems.
- $F_2 \times F_2$  has undecidable generalized word problem (Mihailova)
- Free solvable groups of derived length  $\geq 3$  and rank  $\geq 2$  have undecidable generalized word problem (Umirbaev).
- Compare: all finitely generated metabelian groups have decidable generalized word problem (Romanovskii).

# The generalized word problem

- The above decision problems were listed in order of difficulty.
- It is natural to search for groups distinguishing these problems.
- $F_2 \times F_2$  has undecidable generalized word problem (Mihailova)
- Free solvable groups of derived length  $\geq 3$  and rank  $\geq 2$  have undecidable generalized word problem (Umirbaev).
- Compare: all finitely generated metabelian groups have decidable generalized word problem (Romanovskii).
- The Rips construction produces hyperbolic groups with undecidable generalized word problem.

## Intermezzo: Finite automata

- A finite automaton  $\mathcal{A}$  over an alphabet  $\Sigma$  consists of:

## Intermezzo: Finite automata

- A **finite automaton**  $\mathcal{A}$  over an alphabet  $\Sigma$  consists of:
  - a finite directed graph with edges labeled by elements of  $\Sigma$ ;

## Intermezzo: Finite automata

- A **finite automaton**  $\mathcal{A}$  over an alphabet  $\Sigma$  consists of:
  - a finite directed graph with edges labeled by elements of  $\Sigma$ ;
  - a distinguished **initial** vertex;



## Intermezzo: Finite automata

- A **finite automaton**  $\mathcal{A}$  over an alphabet  $\Sigma$  consists of:
  - a finite directed graph with edges labeled by elements of  $\Sigma$ ;
  - a distinguished **initial** vertex;
  - a set of **final** vertices.

## Intermezzo: Finite automata

- A **finite automaton**  $\mathcal{A}$  over an alphabet  $\Sigma$  consists of:
  - a finite directed graph with edges labeled by elements of  $\Sigma$ ;
  - a distinguished **initial** vertex;
  - a set of **final** vertices.
- The **language**  $L(\mathcal{A})$  of the automaton consists of all words labeling a path from the initial vertex to a final vertex.

## Intermezzo: Finite automata

- A **finite automaton**  $\mathcal{A}$  over an alphabet  $\Sigma$  consists of:
  - a finite directed graph with edges labeled by elements of  $\Sigma$ ;
  - a distinguished **initial** vertex;
  - a set of **final** vertices.
- The **language**  $L(\mathcal{A})$  of the automaton consists of all words labeling a path from the initial vertex to a final vertex.
- A language is called **rational** if it is accepted by some finite automaton.

## Intermezzo: Finite automata

- A **finite automaton**  $\mathcal{A}$  over an alphabet  $\Sigma$  consists of:
  - a finite directed graph with edges labeled by elements of  $\Sigma$ ;
  - a distinguished **initial** vertex;
  - a set of **final** vertices.
- The **language**  $L(\mathcal{A})$  of the automaton consists of all words labeling a path from the initial vertex to a final vertex.
- A language is called **rational** if it is accepted by some finite automaton.
- **Examples:**
  - The language of geodesic words in a hyperbolic group;

## Intermezzo: Finite automata

- A **finite automaton**  $\mathcal{A}$  over an alphabet  $\Sigma$  consists of:
  - a finite directed graph with edges labeled by elements of  $\Sigma$ ;
  - a distinguished **initial** vertex;
  - a set of **final** vertices.
- The **language**  $L(\mathcal{A})$  of the automaton consists of all words labeling a path from the initial vertex to a final vertex.
- A language is called **rational** if it is accepted by some finite automaton.
- **Examples:**
  - The language of geodesic words in a hyperbolic group;
  - The language of geodesic words belonging to a quasiconvex subgroup of a hyperbolic group.

# Rational subsets

- Let  $\text{Rat}(G)$  be the collection of rational subsets of  $G$ , i.e., sets of the form  $\pi(L(\mathcal{A}))$  with  $\mathcal{A}$  a finite automaton.

# Rational subsets

- Let  $\text{Rat}(G)$  be the collection of rational subsets of  $G$ , i.e., sets of the form  $\pi(L(\mathcal{A}))$  with  $\mathcal{A}$  a finite automaton.
- $\text{Rat}(G)$  is the smallest collection of subsets of  $G$  containing the finite subsets and closed under:

# Rational subsets

- Let  $\text{Rat}(G)$  be the collection of rational subsets of  $G$ , i.e., sets of the form  $\pi(L(\mathcal{A}))$  with  $\mathcal{A}$  a finite automaton.
- $\text{Rat}(G)$  is the smallest collection of subsets of  $G$  containing the finite subsets and closed under:
  - union;



# Rational subsets

- Let  $\text{Rat}(G)$  be the collection of rational subsets of  $G$ , i.e., sets of the form  $\pi(L(\mathcal{A}))$  with  $\mathcal{A}$  a finite automaton.
- $\text{Rat}(G)$  is the smallest collection of subsets of  $G$  containing the finite subsets and closed under:
  - union;
  - product;

# Rational subsets

- Let  $\text{Rat}(G)$  be the collection of **rational subsets** of  $G$ , i.e., sets of the form  $\pi(L(\mathcal{A}))$  with  $\mathcal{A}$  a finite automaton.
- $\text{Rat}(G)$  is the smallest collection of subsets of  $G$  containing the finite subsets and closed under:
  - union;
  - product;
  - generation of submonoids  $X \mapsto X^*$ .

# Rational subsets

- Let  $\text{Rat}(G)$  be the collection of rational subsets of  $G$ , i.e., sets of the form  $\pi(L(\mathcal{A}))$  with  $\mathcal{A}$  a finite automaton.
- $\text{Rat}(G)$  is the smallest collection of subsets of  $G$  containing the finite subsets and closed under:
  - union;
  - product;
  - generation of submonoids  $X \mapsto X^*$ .
- **Examples:**
  - finitely generated subgroups;

# Rational subsets


- Let  $\text{Rat}(G)$  be the collection of rational subsets of  $G$ , i.e., sets of the form  $\pi(L(\mathcal{A}))$  with  $\mathcal{A}$  a finite automaton.
- $\text{Rat}(G)$  is the smallest collection of subsets of  $G$  containing the finite subsets and closed under:
  - union;
  - product;
  - generation of submonoids  $X \mapsto X^*$ .
- **Examples:**
  - finitely generated subgroups;
  - finitely generated submonoids;

# Rational subsets

- Let  $\text{Rat}(G)$  be the collection of **rational subsets** of  $G$ , i.e., sets of the form  $\pi(L(\mathcal{A}))$  with  $\mathcal{A}$  a finite automaton.
- $\text{Rat}(G)$  is the smallest collection of subsets of  $G$  containing the finite subsets and closed under:
  - union;
  - product;
  - generation of submonoids  $X \mapsto X^*$ .
- **Examples:**
  - finitely generated subgroups;
  - finitely generated submonoids;
  - double cosets of finitely generated subgroups.

# Examples

- The automaton



recognizes the submonoid  $\{g, h\}^*$  generated by  $g, h$ .


# Examples

- The automaton



recognizes the submonoid  $\{g, h\}^*$  generated by  $g, h$ .

- The automaton



recognizes the double coset  $\langle g_1, g_2 \rangle g \langle g_1, g_2 \rangle$ .

# The theorem of Anissimov and Seifert

## Theorem (Anissimov, Seifert)

*A subgroup  $H \leq G$  belongs to  $\text{Rat}(G)$  iff  $H$  is finitely generated.*



# The theorem of Anissimov and Seifert

## Theorem (Anissimov, Seifert)

*A subgroup  $H \leq G$  belongs to  $\text{Rat}(G)$  iff  $H$  is finitely generated.*

- Rational submonoids need not be finitely generated.

# The theorem of Anissimov and Seifert

## Theorem (Anissimov, Seifert)

*A subgroup  $H \leq G$  belongs to  $\text{Rat}(G)$  iff  $H$  is finitely generated.*

- Rational submonoids need not be finitely generated.
- Rational subsets are not in general closed under complement and intersection.

# The theorem of Anissimov and Seifert

## Theorem (Anissimov, Seifert)

*A subgroup  $H \leq G$  belongs to  $\text{Rat}(G)$  iff  $H$  is finitely generated.*

- Rational submonoids need not be finitely generated.
- Rational subsets are not in general closed under complement and intersection.
- If  $\text{Rat}(G)$  is closed under intersection, then  $G$  is a Howson group.

## Why rational subsets?

- Diekert, Gutiérrez and Hagenah showed solving equations with rational constraints over free groups is PSPACE-complete.

## Why rational subsets?

- Diekert, Gutiérrez and Hagenah showed solving equations with rational constraints over free groups is PSPACE-complete.
- Diekert and Lohrey used this to solve equations and decide the positive theory for graph groups.

## Why rational subsets?

- Diekert, Gutiérrez and Hagenah showed solving equations with rational constraints over free groups is PSPACE-complete.
- Diekert and Lohrey used this to solve equations and decide the positive theory for graph groups.
- Dahmani and Guirardel solved equations over hyperbolic groups with special rational constraints.

## Why rational subsets?

- Diekert, Gutiérrez and Hagenah showed solving equations with rational constraints over free groups is PSPACE-complete.
- Diekert and Lohrey used this to solve equations and decide the positive theory for graph groups.
- Dahmani and Guirardel solved equations over hyperbolic groups with special rational constraints.
- Dahmani and Groves use rational subsets in their solution to the isomorphism problem for toral relatively hyperbolic groups.

## Why rational subsets?

- Diekert, Gutiérrez and Hagenah showed solving equations with rational constraints over free groups is PSPACE-complete.
- Diekert and Lohrey used this to solve equations and decide the positive theory for graph groups.
- Dahmani and Guirardel solved equations over hyperbolic groups with special rational constraints.
- Dahmani and Groves use rational subsets in their solution to the isomorphism problem for toral relatively hyperbolic groups.
- The order of  $g$  is finite if and only if  $g^{-1} \in g^*$ , so decidability of submonoid membership gives decidability of order.



# History

Theorem (Benois (1969))

*Rational subset membership is decidable for free groups.*

# History

## Theorem (Benois (1969))

*Rational subset membership is decidable for free groups.*

- The proof uses an automata theoretic analogue of Stallings folding.

# History

## Theorem (Benois (1969))

*Rational subset membership is decidable for free groups.*

- The proof uses an automata theoretic analogue of Stallings folding.

## Theorem (Eilenberg, Schützenberger (1969))

*Rational subset membership in an abelian group is decidable.*

# History

## Theorem (Benois (1969))

*Rational subset membership is decidable for free groups.*

- The proof uses an automata theoretic analogue of Stallings folding.

## Theorem (Eilenberg, Schützenberger (1969))

*Rational subset membership in an abelian group is decidable.*

- It reduces to INTEGER PROGRAMMING.

## Recent history

- Decidability of rational subset membership is a virtual property (Grunschlag 1999).

## Recent history

- Decidability of rational subset membership is a virtual property (Grunschlag 1999).
- For every  $c \geq 2$ , there is an  $r \gg 1$  so that the free nilpotent group of class  $c$  and rank  $r$  has undecidable rational subset membership (Roman'kov 1999).

## Recent history

- Decidability of rational subset membership is a virtual property (Grunschlag 1999).
- For every  $c \geq 2$ , there is an  $r \gg 1$  so that the free nilpotent group of class  $c$  and rank  $r$  has undecidable rational subset membership (Roman'kov 1999).
- The decidability of rational subset membership passes through free products (Nedbaj 2000).

## Recent history

- Decidability of rational subset membership is a virtual property (Grunschlag 1999).
- For every  $c \geq 2$ , there is an  $r \gg 1$  so that the free nilpotent group of class  $c$  and rank  $r$  has undecidable rational subset membership (Roman'kov 1999).
- The decidability of rational subset membership passes through free products (Nedbaj 2000).
- Independently, with Kambites and Silva, we proved:



## Recent history

- Decidability of rational subset membership is a virtual property (Grunschlag 1999).
- For every  $c \geq 2$ , there is an  $r \gg 1$  so that the free nilpotent group of class  $c$  and rank  $r$  has undecidable rational subset membership (Roman'kov 1999).
- The decidability of rational subset membership passes through free products (Nedbaj 2000).
- Independently, with Kambites and Silva, we proved:

### Theorem (Kambites, Silva, BS (2007))

*Decidability of rational subset membership is preserved by free products with amalgamation and HNN-extensions with finite edge groups.*

## A general decidability result

- Let  $\mathcal{C}$  be the smallest class of groups containing the trivial group and closed under:

## A general decidability result

- Let  $\mathcal{C}$  be the smallest class of groups containing the trivial group and closed under:
  - Taking finitely generated subgroups;

## A general decidability result

- Let  $\mathcal{C}$  be the smallest class of groups containing the trivial group and closed under:
  - Taking finitely generated subgroups;
  - Taking finite index overgroups;

## A general decidability result

- Let  $\mathcal{C}$  be the smallest class of groups containing the trivial group and closed under:
  - Taking finitely generated subgroups;
  - Taking finite index overgroups;
  - Free products with amalgamation and HNN extensions with finite edge groups;

## A general decidability result

- Let  $\mathcal{C}$  be the smallest class of groups containing the trivial group and closed under:
  - Taking finitely generated subgroups;
  - Taking finite index overgroups;
  - Free products with amalgamation and HNN extensions with finite edge groups;
  - Direct product with  $\mathbb{Z}$ .

## A general decidability result

- Let  $\mathcal{C}$  be the smallest class of groups containing the trivial group and closed under:
  - Taking finitely generated subgroups;
  - Taking finite index overgroups;
  - Free products with amalgamation and HNN extensions with finite edge groups;
  - Direct product with  $\mathbb{Z}$ .
- The following theorem encompasses all groups known to have decidable rational subset membership.

## A general decidability result

- Let  $\mathcal{C}$  be the smallest class of groups containing the trivial group and closed under:
  - Taking finitely generated subgroups;
  - Taking finite index overgroups;
  - Free products with amalgamation and HNN extensions with finite edge groups;
  - Direct product with  $\mathbb{Z}$ .
- The following theorem encompasses all groups known to have decidable rational subset membership.

### Theorem (Lohrey, BS (2008))

*Every group in the class  $\mathcal{C}$  has decidable rational subset membership problem.*



# Graph groups: the generalized word problem

- For  $\Gamma$  a graph, the associated **graph group** is


$$\mathcal{G}(\Gamma) = \langle V(\Gamma) \mid [v, w] : (v, w) \in E(\Gamma) \rangle.$$

# Graph groups: the generalized word problem

- For  $\Gamma$  a graph, the associated **graph group** is

$$\mathcal{G}(\Gamma) = \langle V(\Gamma) \mid [v, w] : (v, w) \in E(\Gamma) \rangle.$$

- Let




# Graph groups: the generalized word problem

- For  $\Gamma$  a graph, the associated **graph group** is

$$\mathcal{G}(\Gamma) = \langle V(\Gamma) \mid [v, w] : (v, w) \in E(\Gamma) \rangle.$$

- Let




- Then  $\mathcal{G}(C_4) = F_2 \times F_2$  and so this group has undecidable generalized word problem.

# Graph groups: the generalized word problem

- For  $\Gamma$  a graph, the associated **graph group** is

$$\mathcal{G}(\Gamma) = \langle V(\Gamma) \mid [v, w] : (v, w) \in E(\Gamma) \rangle.$$

- Let




- Then  $\mathcal{G}(C_4) = F_2 \times F_2$  and so this group has undecidable generalized word problem.
- A graph is **chordal** if it has no induced cycle of length  $\geq 4$ .

# Graph groups: the generalized word problem

- For  $\Gamma$  a graph, the associated **graph group** is

$$\mathcal{G}(\Gamma) = \langle V(\Gamma) \mid [v, w] : (v, w) \in E(\Gamma) \rangle.$$

- Let




- Then  $\mathcal{G}(C_4) = F_2 \times F_2$  and so this group has undecidable generalized word problem.
- A graph is **chordal** if it has no induced cycle of length  $\geq 4$ .


**Theorem (Kapovich, Myasnikov, Weidmann (2005))**

*The generalized word problem is decidable for chordal graph groups.*

# Graph groups: the rational subset problem

Let  $P_4 = \bullet \text{ --- } \bullet \text{ --- } \bullet \text{ --- } \bullet$  and  $C_4 =$  


# Graph groups: the rational subset problem

Let  $P_4 = \bullet \text{ --- } \bullet \text{ --- } \bullet \text{ --- } \bullet$  and  $C_4 =$  

Theorem (Lohrey, BS (2008))

*Let  $\Gamma$  be a graph. Then the following are equivalent:*

# Graph groups: the rational subset problem

Let  $P_4 = \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet$  and  $C_4 =$  


Theorem (Lohrey, BS (2008))

Let  $\Gamma$  be a graph. Then the following are equivalent:

1. rational subset membership is decidable for  $\mathcal{G}(\Gamma)$ ;



# Graph groups: the rational subset problem


Let  $P_4 = \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet$  and  $C_4 =$  

Theorem (Lohrey, BS (2008))

Let  $\Gamma$  be a graph. Then the following are equivalent:

1. rational subset membership is decidable for  $\mathcal{G}(\Gamma)$ ;
2. submonoid membership is decidable for  $\mathcal{G}(\Gamma)$ ;

# Graph groups: the rational subset problem


Let  $P_4 = \bullet \text{ --- } \bullet \text{ --- } \bullet \text{ --- } \bullet$  and  $C_4 =$  

Theorem (Lohrey, BS (2008))

Let  $\Gamma$  be a graph. Then the following are equivalent:

1. rational subset membership is decidable for  $\mathcal{G}(\Gamma)$ ;
2. submonoid membership is decidable for  $\mathcal{G}(\Gamma)$ ;
3.  $\Gamma$  contains neither an induced  $C_4$  nor  $P_4$ .

# Graph groups: the rational subset problem

Let  $P_4 = \bullet \text{ --- } \bullet \text{ --- } \bullet \text{ --- } \bullet$  and  $C_4 =$  

Theorem (Lohrey, BS (2008))

Let  $\Gamma$  be a graph. Then the following are equivalent:

1. rational subset membership is decidable for  $\mathcal{G}(\Gamma)$ ;
2. submonoid membership is decidable for  $\mathcal{G}(\Gamma)$ ;
3.  $\Gamma$  contains neither an induced  $C_4$  nor  $P_4$ .

Since  $P_4$  is chordal, we obtain the first example of a group with decidable generalized word problem but undecidable submonoid membership problem.

# The direct product of two free monoids

## Theorem (Lohrey, BS)

*Any group containing a direct product of two free monoids has undecidable rational subset membership problem.*

# The direct product of two free monoids

## Theorem (Lohrey, BS)

*Any group containing a direct product of two free monoids has undecidable rational subset membership problem.*

- This is a simple encoding of the Post correspondence problem.

## Submonoids vs. rational subsets

- The submonoid and rational subset membership problems are equivalent for graph groups.

## Submonoids vs. rational subsets

- The submonoid and rational subset membership problems are equivalent for graph groups.
- We have no examples of a group with decidable submonoid membership but undecidable rational subset membership.

## Submonoids vs. rational subsets

- The submonoid and rational subset membership problems are equivalent for graph groups.
- We have no examples of a group with decidable submonoid membership but undecidable rational subset membership.
- In fact, we have the following result:



## Submonoids vs. rational subsets

- The submonoid and rational subset membership problems are equivalent for graph groups.
- We have no examples of a group with decidable submonoid membership but undecidable rational subset membership.
- In fact, we have the following result:

### Theorem (Lohrey, BS (2010))

*The submonoid and rational subset membership problems are equivalent for groups with two or more ends.*

## Submonoids vs. rational subsets

- The submonoid and rational subset membership problems are equivalent for graph groups.
- We have no examples of a group with decidable submonoid membership but undecidable rational subset membership.
- In fact, we have the following result:

### Theorem (Lohrey, BS (2010))

*The submonoid and rational subset membership problems are equivalent for groups with two or more ends.*

- Recall: a group has 2 or more ends iff it splits over a finite subgroup.

# Metabelian groups

- A group  $G$  is **metabelian** if  $[G, G]$  is abelian.

# Metabelian groups

- A group  $G$  is **metabelian** if  $[G, G]$  is abelian.
- If  $G$  is finitely generated, there is an exact sequence

$$1 \longrightarrow [G, G] \longrightarrow G \longrightarrow G/[G, G] \longrightarrow 1$$

with  $[G, G]$  a finitely generated  $G/[G, G]$ -module.

# Metabelian groups

- A group  $G$  is **metabelian** if  $[G, G]$  is abelian.
- If  $G$  is finitely generated, there is an exact sequence

$$1 \longrightarrow [G, G] \longrightarrow G \longrightarrow G/[G, G] \longrightarrow 1$$

with  $[G, G]$  a finitely generated  $G/[G, G]$ -module.

- The generalized word problem reduces to the membership problem for finitely generated submodules of modules over group rings of abelian groups.

# Metabelian groups

- A group  $G$  is **metabelian** if  $[G, G]$  is abelian.
- If  $G$  is finitely generated, there is an exact sequence

$$1 \longrightarrow [G, G] \longrightarrow G \longrightarrow G/[G, G] \longrightarrow 1$$

with  $[G, G]$  a finitely generated  $G/[G, G]$ -module.

- The generalized word problem reduces to the membership problem for finitely generated submodules of modules over group rings of abelian groups.
- This latter membership problem is decidable.

# Subsemimodule membership problem

- A  $G$ -module is an abelian group with an action of  $G$  by automorphisms.

# Subsemimodule membership problem

- A  $G$ -module is an abelian group with an action of  $G$  by automorphisms.
- A  $G$ -semimodule is a commutative monoid with an action of  $G$  by automorphisms.



# Subsemimodule membership problem

- A  $G$ -module is an abelian group with an action of  $G$  by automorphisms.
- A  $G$ -semimodule is a commutative monoid with an action of  $G$  by automorphisms.
- THE SUBSEMIMODULE MEMBERSHIP PROBLEM:
  - Given a finitely generated  $G$ -module  $M$ , a finite subset  $F$  of  $M$  and an element  $m \in M$ , decide whether  $m \in \mathbb{N}G \cdot F$ .

# Subsemimodule membership problem

- A  $G$ -module is an abelian group with an action of  $G$  by automorphisms.
- A  $G$ -semimodule is a commutative monoid with an action of  $G$  by automorphisms.
- THE SUBSEMIMODULE MEMBERSHIP PROBLEM:
  - Given a finitely generated  $G$ -module  $M$ , a finite subset  $F$  of  $M$  and an element  $m \in M$ , decide whether  $m \in \mathbb{N}G \cdot F$ .
- When  $G$  is trivial, the subsemimodule membership problem is precisely INTEGER PROGRAMMING.

# Subsemimodule membership problem

- A  $G$ -module is an abelian group with an action of  $G$  by automorphisms.
- A  $G$ -semimodule is a commutative monoid with an action of  $G$  by automorphisms.
- THE SUBSEMIMODULE MEMBERSHIP PROBLEM:
  - Given a finitely generated  $G$ -module  $M$ , a finite subset  $F$  of  $M$  and an element  $m \in M$ , decide whether  $m \in \mathbb{N}G \cdot F$ .
- When  $G$  is trivial, the subsemimodule membership problem is precisely INTEGER PROGRAMMING.
- Recall: INTEGER PROGRAMMING is NP-complete.

## Subsemimodule membership problem II

- Let  $M$  be a finitely generated  $G$ -module.

## Subsemimodule membership problem II

- Let  $M$  be a finitely generated  $G$ -module.
- Suppose  $N$  is a finitely generated  $G$ -subsemimodule with undecidable membership problem.

## Subsemimodule membership problem II

- Let  $M$  be a finitely generated  $G$ -module.
- Suppose  $N$  is a finitely generated  $G$ -subsemimodule with undecidable membership problem.
- Then  $N \rtimes G$  is a finitely generated submonoid of  $M \rtimes G$  with undecidable membership problem.

## Subsemimodule membership problem II

- Let  $M$  be a finitely generated  $G$ -module.
- Suppose  $N$  is a finitely generated  $G$ -subsemimodule with undecidable membership problem.
- Then  $N \rtimes G$  is a finitely generated submonoid of  $M \rtimes G$  with undecidable membership problem.

### Theorem (Lohrey, BS (2011))

*There is a finitely generated free  $\mathbb{Z} \times \mathbb{Z}$ -module containing a finitely generated subsemimodule with undecidable membership problem.*

## Subsemimodule membership problem II

- Let  $M$  be a finitely generated  $G$ -module.
- Suppose  $N$  is a finitely generated  $G$ -subsemimodule with undecidable membership problem.
- Then  $N \rtimes G$  is a finitely generated submonoid of  $M \rtimes G$  with undecidable membership problem.

### Theorem (Lohrey, BS (2011))

*There is a finitely generated free  $\mathbb{Z} \times \mathbb{Z}$ -module containing a finitely generated subsemimodule with undecidable membership problem.*

- The proof encodes an undecidable problem on planar tilings.



## The wreath product $\mathbb{Z} \wr (\mathbb{Z} \times \mathbb{Z})$

- If  $G$  is a group, then  $\mathbb{Z} \wr G = \mathbb{Z}G \rtimes G$ .

## The wreath product $\mathbb{Z} \wr (\mathbb{Z} \times \mathbb{Z})$

- If  $G$  is a group, then  $\mathbb{Z} \wr G = \mathbb{Z}G \rtimes G$ .
- Set  $G = \mathbb{Z} \times \mathbb{Z}$  and let  $r$  be large enough so that  $\mathbb{Z}G^r$  contains a finitely generated subsemimodule with undecidable membership.

## The wreath product $\mathbb{Z} \wr (\mathbb{Z} \times \mathbb{Z})$

- If  $G$  is a group, then  $\mathbb{Z} \wr G = \mathbb{Z}G \rtimes G$ .
- Set  $G = \mathbb{Z} \times \mathbb{Z}$  and let  $r$  be large enough so that  $\mathbb{Z}G^r$  contains a finitely generated subsemimodule with undecidable membership.
- Let  $H = r\mathbb{Z} \times \mathbb{Z}$ ; so  $[G : H] = r$ .

## The wreath product $\mathbb{Z} \wr (\mathbb{Z} \times \mathbb{Z})$

- If  $G$  is a group, then  $\mathbb{Z} \wr G = \mathbb{Z}G \rtimes G$ .
- Set  $G = \mathbb{Z} \times \mathbb{Z}$  and let  $r$  be large enough so that  $\mathbb{Z}G^r$  contains a finitely generated subsemimodule with undecidable membership.
- Let  $H = r\mathbb{Z} \times \mathbb{Z}$ ; so  $[G : H] = r$ .
- Then  $\mathbb{Z}G$  is a free  $\mathbb{Z}H$ -module of rank  $r$  and so contains a finitely generated  $H$ -subsemimodule with undecidable membership.

## The wreath product $\mathbb{Z} \wr (\mathbb{Z} \times \mathbb{Z})$

- If  $G$  is a group, then  $\mathbb{Z} \wr G = \mathbb{Z}G \rtimes G$ .
- Set  $G = \mathbb{Z} \times \mathbb{Z}$  and let  $r$  be large enough so that  $\mathbb{Z}G^r$  contains a finitely generated subsemimodule with undecidable membership.
- Let  $H = r\mathbb{Z} \times \mathbb{Z}$ ; so  $[G : H] = r$ .
- Then  $\mathbb{Z}G$  is a free  $\mathbb{Z}H$ -module of rank  $r$  and so contains a finitely generated  $H$ -subsemimodule with undecidable membership.

### Theorem (Lohrey, BS (2011))

*The wreath product  $\mathbb{Z} \wr (\mathbb{Z} \times \mathbb{Z})$  has undecidable submonoid membership problem.*

# The free metabelian group

- Let  $M_2$  be a free metabelian group on  $\{x, y\}$ .

# The free metabelian group

- Let  $M_2$  be a free metabelian group on  $\{x, y\}$ .
- The word problem for  $M_2$  has a geometric description:

# The free metabelian group

- Let  $M_2$  be a free metabelian group on  $\{x, y\}$ .
- The word problem for  $M_2$  has a geometric description:
  - A word  $w$  is 1 in  $M_2$  iff it labels a loop in the Cayley graph  $\Gamma$  of  $\mathbb{Z} \times \mathbb{Z}$  that is trivial in homology.



## The free metabelian group

- Let  $M_2$  be a free metabelian group on  $\{x, y\}$ .
- The word problem for  $M_2$  has a geometric description:
  - A word  $w$  is 1 in  $M_2$  iff it labels a loop in the Cayley graph  $\Gamma$  of  $\mathbb{Z} \times \mathbb{Z}$  that is trivial in homology.
- $[M_2, M_2]$  can be identified with  $H_1(\Gamma)$  as a  $\mathbb{Z} \times \mathbb{Z}$ -module and hence is free of rank 1 (generated by  $[x, y]$ ).

## The free metabelian group

- Let  $M_2$  be a free metabelian group on  $\{x, y\}$ .
- The word problem for  $M_2$  has a geometric description:
  - A word  $w$  is 1 in  $M_2$  iff it labels a loop in the Cayley graph  $\Gamma$  of  $\mathbb{Z} \times \mathbb{Z}$  that is trivial in homology.
- $[M_2, M_2]$  can be identified with  $H_1(\Gamma)$  as a  $\mathbb{Z} \times \mathbb{Z}$ -module and hence is free of rank 1 (generated by  $[x, y]$ ).
- Putting  $G = \mathbb{Z} \times \mathbb{Z}$  we have an exact sequence

$$1 \longrightarrow \mathbb{Z}G \longrightarrow M_2 \longrightarrow G \longrightarrow 1$$

## The free metabelian group

- Let  $M_2$  be a free metabelian group on  $\{x, y\}$ .
- The word problem for  $M_2$  has a geometric description:
  - A word  $w$  is 1 in  $M_2$  iff it labels a loop in the Cayley graph  $\Gamma$  of  $\mathbb{Z} \times \mathbb{Z}$  that is trivial in homology.
- $[M_2, M_2]$  can be identified with  $H_1(\Gamma)$  as a  $\mathbb{Z} \times \mathbb{Z}$ -module and hence is free of rank 1 (generated by  $[x, y]$ ).
- Putting  $G = \mathbb{Z} \times \mathbb{Z}$  we have an exact sequence

$$1 \longrightarrow \mathbb{Z}G \longrightarrow M_2 \longrightarrow G \longrightarrow 1$$

- If  $H = \langle x^m, y \rangle$ , then  $[H, H]$  is a rank 1 direct summand (as an  $m\mathbb{Z} \times \mathbb{Z}$ -module) in  $\mathbb{Z}G$ .

## The free metabelian group

- Let  $M_2$  be a free metabelian group on  $\{x, y\}$ .
- The word problem for  $M_2$  has a geometric description:
  - A word  $w$  is 1 in  $M_2$  iff it labels a loop in the Cayley graph  $\Gamma$  of  $\mathbb{Z} \times \mathbb{Z}$  that is trivial in homology.
- $[M_2, M_2]$  can be identified with  $H_1(\Gamma)$  as a  $\mathbb{Z} \times \mathbb{Z}$ -module and hence is free of rank 1 (generated by  $[x, y]$ ).
- Putting  $G = \mathbb{Z} \times \mathbb{Z}$  we have an exact sequence

$$1 \longrightarrow \mathbb{Z}G \longrightarrow M_2 \longrightarrow G \longrightarrow 1$$

- If  $H = \langle x^m, y \rangle$ , then  $[H, H]$  is a rank 1 direct summand (as an  $m\mathbb{Z} \times \mathbb{Z}$ -module) in  $\mathbb{Z}G$ .
- Choosing  $m$  large enough, we can encode our subsemimodule with undecidable membership as a finitely generated submonoid of  $M_2$ .

# The free metabelian group II

Theorem (Lohrey, BS (2011))

*The free metabelian group of rank 2 has undecidable submonoid membership problem.*

# The free metabelian group II

## Theorem (Lohrey, BS (2011))

*The free metabelian group of rank 2 has undecidable submonoid membership problem.*

Using similar ideas, we proved:

# The free metabelian group II

## Theorem (Lohrey, BS (2011))

*The free metabelian group of rank 2 has undecidable submonoid membership problem.*

Using similar ideas, we proved:

## Theorem (Lohrey, BS (2011))

*The 2-dimensional lamplighter group  $\mathbb{Z}/2\mathbb{Z} \wr (\mathbb{Z} \times \mathbb{Z})$  has undecidable rational subset membership problem.*

# Open questions

## Question

*Does there exist a group with decidable submonoid membership and undecidable rational subset membership?*



# Open questions

## Question

*Does there exist a group with decidable submonoid membership and undecidable rational subset membership?*

This question is equivalent to the following one.

# Open questions

## Question

*Does there exist a group with decidable submonoid membership and undecidable rational subset membership?*

This question is equivalent to the following one.

## Question

*Is decidability of submonoid membership preserved by free products?*

# Open questions

## Question

*Does there exist a group with decidable submonoid membership and undecidable rational subset membership?*

This question is equivalent to the following one.

## Question

*Is decidability of submonoid membership preserved by free products?*

## Question

*Does  $\mathbb{Z} \wr \mathbb{Z}$  have decidable submonoid membership problem?*

# Open questions

## Question

*Does there exist a group with decidable submonoid membership and undecidable rational subset membership?*

This question is equivalent to the following one.

## Question

*Is decidability of submonoid membership preserved by free products?*

## Question

*Does  $\mathbb{Z} \wr \mathbb{Z}$  have decidable submonoid membership problem?*

## Question

*Is submonoid membership decidable for nilpotent groups?*

The end

THANK YOU FOR YOUR  
ATTENTION!