

3.3 Partial Derivatives_Guide

Monday, January 12, 2026

10:40 PM




3.3 Partial
Derivative...

3.3 Partial Derivatives

Motivation: Suppose we have a surface given by the two-variable function $z = f(x, y)$ and a point P (with coordinates of (a, b, c)) on the surface determined by f .

Suppose we want to know the rate at which the z -coordinate changes at P when we are moving across the surface parallel to the x -axis. This means that we would be moving along the curve C_1 toward P in the picture above. At this point P we can imagine a tangent line T_1 being constructed.

The slope of T_1 would represent the rate of change of the z -coordinate at P when moving parallel to the x -axis.



Suppose we want to know the rate at which the z -coordinate changes at P when we are moving across the surface parallel to the y -axis. This means that we would be moving along the curve C_2 toward P in the picture above. At this point P we can imagine a tangent line T_2 being constructed.

The slope of T_2 would represent the rate of change of the z -coordinate at P when moving parallel to the y -axis.

In Calculus I we found slopes of tangent lines to a curve $y = f(x)$ by calculating the derivative $f'(x)$ and evaluating it at the appropriate point.

In Calculus III for a function $z = f(x, y)$ we also need a way to find a derivative, but in a way that allows us to account for the direction that we are moving along the surface. This leads to the idea of what we call partial derivatives.

Note: The information that we are seeking above can be key when one wishes to better understand how quickly a surface is changing at a point.

Definition: If f is a function of two variables, its partial derivatives are the functions f_x and f_y defined

$$\text{by: } f_x(x, y) = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h} \quad \text{and} \quad f_y(x, y) = \lim_{h \rightarrow 0} \frac{f(x, y+h) - f(x, y)}{h}$$

$\uparrow$ y is fixed $\uparrow$ x is fixed

Note: $f_x(x, y)$ represents the rate of change of z with respect to x when y is fixed, and $f_y(x, y)$ is the rate of change of z with respect to y when x is fixed.

Note: In the context of the motivation, we would use f_x to find the slope of the line tangent to the brown curve at P , and we would use f_y to find the slope of the line tangent to the green curve at P .

Notation: There are many notations for partial derivatives. We go over 7 possibilities. If $z = f(x, y)$, we write

$$\begin{aligned} f_x(x, y) &= \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} f(x, y) = \frac{\partial z}{\partial x} = f_1 = D_1 f = D_x f \\ f_y(x, y) &= \frac{\partial f}{\partial y} = \frac{\partial}{\partial y} f(x, y) = \frac{\partial z}{\partial y} = f_2 = D_2 f = D_y f \end{aligned}$$

Typically use

Example 1: If $f(x, y) = 2xy^3 + 6xy - x^7$, find $f_x(x, y)$ and $f_y(x, y)$.

To find f_x we imagine y is a constant.

$$f_x(x, y) = 2y^3 + 6y - 7x^6$$

To find f_y we imagine x is a constant

$$f_y(x, y) = 6xy^2 + 6x$$


Formulas to find
slopes of
tangent lines

Example 2: Consider the surface given by $f(x, y) = 4 - x^2 - 2y^2$. Let C be the curve of intersection between this surface and the plane $x = 2$. Find the slope of the line tangent to C at the point $(2, 1, -2)$ on the surface.

$$\text{We compute } f_y(x, y) = 0 - 0 - 4y = -4y$$

$$\text{So } f_y(2, 1) = -4(1) = \boxed{-4}$$

Slope



Example 3: Find $\frac{\partial z}{\partial x}$ if z is defined implicitly as a function of x and y by $x^3 + y^3 + z^3 + 6xyz = 1$.

$$\frac{\partial}{\partial x} [x^3 + y^3 + z^3 + 6xy \cdot z] = \frac{\partial}{\partial x} [1]$$

$$3x^2 + 0 + 3z^2 \cdot \frac{\partial z}{\partial x} + (6xy \cdot \frac{\partial z}{\partial x} + 6yz) = 0$$

$$(3z^2 + 6xy) \frac{\partial z}{\partial x} = -3x^2 - 6yz$$

$$\frac{\partial z}{\partial x} = \frac{-3x^2 - 6yz}{(3z^2 + 6xy)}$$

Note: There is a fairly subtle point that could be made about the above example. In particular, we assumed the equation defines z implicitly as a function of x and y . What guarantees that the above equation defines z implicitly as a function of x and y ? In order to fully answer this question one needs the implicit function theorem (the proof of which is beyond the scope of this course).

Definition: If f is a function of two variables, then f_x and f_y are functions of two variables. The functions $(f_x)_x$, $(f_x)_y$, $(f_y)_x$, and $(f_y)_y$ are called the second partial derivatives of f . If $z = f(x, y)$, we use the following notation:

$$\begin{aligned}(f_x)_x &= f_{xx} = f_{11} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 z}{\partial x^2} \\(f_x)_y &= f_{xy} = f_{12} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial^2 z}{\partial y \partial x} \\(f_y)_x &= f_{yx} = f_{21} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) = \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 z}{\partial x \partial y} \\(f_y)_y &= f_{yy} = f_{22} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial y} \right) = \frac{\partial^2 f}{\partial y^2} = \frac{\partial^2 z}{\partial y^2}\end{aligned}$$

$\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} [f] \right)$

Example 4: Find all four second partial derivatives of $f(x, y) = x^2 y^2 - y \sin(x^2)$.

$$f_x = 2xy^2 - y \cos(x^2) \cdot 2x$$

$$f_y = 2x^2 y - \sin(x^2)$$

$$f_{xx} = 2y^2 - [y \cos(x^2) \cdot 2 - 2xy \sin(x^2) \cdot 2x]$$

$$f_{xy} = 4xy - \cos(x^2) \cdot 2x$$

$$f_{yy} = 2x^2$$

$$f_{yx} = 4xy - \cos(x^2) \cdot 2x$$

SAME!

Note: Interestingly, in the above example f_{xy} and f_{yx} are the same! The next theorem addresses this.

Note: As one would probably expect, the second partials f_{xx} and f_{yy} tell us something about the concavity of the curve of intersection between the surface given by $f(x, y)$ and a plane of the form $y = a$ or $x = b$. We will see some more applications of these second partials later in chapter 4.

Note: It is more difficult to perceive what the second partials f_{xy} and f_{yx} tell us. What f_{xy} does communicate is how the slope of the tangent line with respect to x changes as the value of y varies (similarly for f_{yx}). To see an animation of what we mean by this one can visit the following website:

https://web.archive.org/web/20180215172659/http://www.math.harvard.edu/archive/21a_fall_08/exhibits/fxy/index.html

Theorem: Suppose f is defined on a disk D that contains the point (a, b) . If the functions f_{xy} and f_{yx} are both continuous on D , then $f_{xy}(a, b) = f_{yx}(a, b)$.

Proof: Beyond the scope of this course

Note: In the most recent example, f_{xy} and f_{yx} are continuous on all of $\mathbb{R}^2$, and we saw that they were indeed equal.

We can have partial derivatives of order 3 and higher.

Example 5: If $f(x, y, z) = \sin(3x + yz)$, then find f_{xy} .

$$f_x = \cos(3x + yz) \cdot (3) + \int h(x, z) dx + k(y, z)$$

$$f_{xx} = -3 \sin(3x + yz) \cdot (3) + h(x, z)$$

$$\begin{aligned} f_{xxy} &= -9 \cos(3x + yz)(z) \\ &= -9z \cos(3x + yz) \end{aligned}$$