

5.1 Vector Fields_Guide

Monday, January 12, 2026 10:44 PM



5.1 Vector
Fields_Gu...

5.1 Vector Fields


Up to this point in MTH 202 we have studied functions of the form $\mathbf{r}: \mathbb{R} \rightarrow \mathbb{R}^n$ and $f: \mathbb{R}^n \rightarrow \mathbb{R}$. In this final chapter of MTH 202 we turn our focus to the most general situation. Namely we will study functions of the form $\mathbf{F}: \mathbb{R}^m \rightarrow \mathbb{R}^n$.

Definition:

1. Let D be a set in $\mathbb{R}^2$. A vector field on $\mathbb{R}^2$ is a function $\mathbf{F}$ that assigns to each point (x, y) in D a two-dimensional vector $\mathbf{F}(x, y)$.
2. Let E be a set in $\mathbb{R}^3$. A vector field on $\mathbb{R}^3$ is a function $\mathbf{F}$ that assigns to each point (x, y, z) in E a three-dimensional vector $\mathbf{F}(x, y, z)$.

Example 1: Use a picture to illustrate the vector field $\mathbf{F}(x, y) = \langle -y, x \rangle$.

(x, y)	$\langle -y, x \rangle$
• $(0, 0)$	$\langle 0, 0 \rangle = \vec{0}$
• $(1, 0)$	$\langle 0, 1 \rangle$
• $(\frac{1}{2}, \frac{1}{2})$	$\langle -\frac{1}{2}, \frac{1}{2} \rangle$



Remark: Vector fields have a wide variety of applications. For example, a vector field on $\mathbb{R}^2$ can be used to model the amount of force exerted by an electric charge on some charge located at the point (x, y) on a two dimensional plate.


A vector field on $\mathbb{R}^3$ can be used to model the force of the wind at some point (x, y, z) in space.

A vector field on $\mathbb{R}^3$ can be used to model the current of the ocean at some point (x, y, z) in the ocean.

A vector field on $\mathbb{R}^3$ can be used to model the force due to gravity at some point (x, y, z) in space.

A vector field in $\mathbb{R}^3$ can be used to model the velocity of a moving fluid at some point (x, y, z) in space.

Example 2: The gravitational force from an object of mass M centered at the origin acting on a small object of mass m at $\mathbf{x} = \langle x, y, z \rangle$ is $\mathbf{F}(x, y, z) = -\frac{mMG}{\|\mathbf{x}\|^3} \mathbf{x}$ where G is the universal gravitational constant. Suppose a large three dimensional object has center of mass at the origin and has mass equal to 100,000 kg. Suppose a small object of mass 5 kg is located at the terminal point of the position vector $\mathbf{x} = \langle 10, 5, 10 \rangle$ (you may assume distance is in meters). What is the gravitational force acting on the small object (due to the large object)?

$$\begin{aligned} \vec{F}(10, 5, 10) &= -\frac{(5)(100,000)G}{(\sqrt{10^2 + 5^2 + 10^2})^3} \cdot \vec{x} \\ &= -\frac{500,000 G}{(15)^3} \langle 10, 5, 10 \rangle \\ &= \left\langle \frac{5,000,000 G}{15^3}, \frac{-2,500,000 G}{(15)^3}, \frac{-5,000,000 G}{(15)^3} \right\rangle \end{aligned}$$


$G = 6.67 \times 10^{-11}$

Definition: If f is a scalar function of two or three variables, $\vec{\nabla} f$ is called the gradient vector field of f .

Definition: A vector field $\mathbf{F}$ is called a conservative vector field if it is the gradient of some scalar function f . Mathematically, this means there must exist a scalar function f so that $\mathbf{F} = \vec{\nabla} f$. In this situation f is called a potential function for $\mathbf{F}$.

Example 3: Prove that $\mathbf{F}(x, y) = \langle x, y \rangle$ is a conservative vector field.

Consider $f(x, y) = \frac{x^2}{2} + \frac{y^2}{2}$.

Note $\nabla f = \vec{F}$

So $\vec{F}$ is conservative.

Question: Is $f(x, y) = \frac{1}{2}x^2 + \frac{1}{2}y^2$ the only scalar function that would have worked in the above proof?

Note: For now, the only technique we have for proving a vector field is conservative is to guess and check until we successfully find a potential function for the vector field. We will develop more sophisticated techniques as in section 5.3.

Note: Right now, we do not have ANY technique for proving that a vector field is not conservative. We will develop techniques for proving a vector field is not conservative in section 5.3.

Example 4: Consider $f(x, y, z) = xye^z$.

a) Find the gradient vector field of f .

$$\nabla f(x, y, z) = \langle ye^z, xe^z, xye^z \rangle$$

b) Is the vector field you found in (a) conservative? Explain.

Yes! It has a potential function of $f(x, y, z)$.

Note: Later in chapter 5 we will discover that conservative vector fields have some very nice properties.