

1. Determine the equation of the line tangent to the curve $f(x) = x^2 e^x$ at $x = 1$.
 2. Determine all values of x where $f(x) = (2x^2 + 4x + 5)(6 - 3x)$ has tangent lines parallel to the line $\frac{1}{2}y - \frac{7}{2}x = 1$.
 3. Determine the equation of the line tangent to the curve $f(x) = \frac{x}{x^3 - 4x}$ at $x = 1$.
 4. Determine all values of x where $f(x) = \frac{x+2}{x+1}$ has tangent lines perpendicular to the line $3x + 6y = 2$.
-
5. Find conditions on a , b , c , & d so that the graph of the polynomial $f(x) = ax^3 + bx^2 + cx + d$ has...
 - a) exactly two horizontal tangents.
 - b) exactly one horizontal tangent.
 - c) no horizontal tangents.
 6. Find k if the curve $y = x^2 + k$ is tangent to the line $y = 2x$.
 7. Find all values of a such that the curves $y = \frac{a}{x-1}$ and $y = x^2 - 2x + 1$ intersect at right angles.
 8. The segment of any tangent line to the graph of $y = \frac{1}{x}$ that is cut off by the coordinate axes is bisected by the point of tangency.
 - a) Show this works for when $x = 1$.
 - b) Show this works for when $x = a$.
-
9. Suppose that the function f is differentiable everywhere and that $F(x) = xf(x)$.
 - a) Express a formula for $F''(x)$ in terms of x and derivatives of f .
 - b) Express a formula for $F'''(x)$ in terms of x and derivatives of f .
 - c) For $n \geq 2$, conjecture of formula for $F^{(n)}(x)$.
 10. Consider the polynomial functions $f(x) = a_1x + a_0$, $g(x) = a_2x^2 + a_1x + a_0$, and $h(x) = a_3x^3 + a_2x^2 + a_1x + a_0$, where $a_0, a_1, a_2, \& a_3$ represent real numbers.
 - a) Find $f''(x)$, $g''(x)$, and $h^{(4)}(x)$.
 - b) If n represents the degree of the polynomial, then what is the $(n+1)$ -st derivative of any polynomial? Why?

11. Newton's Law of Universal Gravitation states that the magnitude F of the force exerted by a point with mass M on a point with mass m is $F = \frac{GmM}{r^2}$ where G is a constant and r is the distance between the bodies. Assuming that the points are moving, find a formula for the instantaneous rate of change of F with respect to r .

12. Show that if $f'(x)$ exists for each x in (a, b) , then $f(x)$ is continuous on (a, b) .

13. The following graph depicts the position of two racers.

- Which racer has a faster average speed? When was each racer running faster than the other (approximately)? Which racer was running fastest at the end of the race?
- On a separate plot from this picture, sketch the graphs of the velocities of these racers.
- Suppose that the distance (from the starting point) of the red racer is given by the function $p_r(t)$ and the distance of the blue racer is given by $p_b(t)$. Suppose that $p_r(t) = 10t$ & $p_b(t) = t^2$. Use this new information (and your knowledge of derivatives) to check whether your answers in parts (a) and (b) were accurate.

