HW 20 Solution

The table below gives the energy use of five gas ranges for seven menu days. (The units are equivalent kilowatt-hours; 0.238kWh=1ft³ of natural gas.) The energy use calculate is an average over several cooks. Your goal is to describe the variation in the table below in terms of the effects of different ranges and different menu days.

```
mydata <- read.csv("https://raw.githubusercontent.com/ProfSuzy/Stat61/main/gasranges.csv",
                   header=T)
mydata
##
     Day Range1 Range2 Range3 Range4 Range5
## 1
           8.25
                  8.26
                         6.55
                                 8.21
                                        6.69
       1
## 2
       2
           5.12
                  4.81
                         3.87
                                4.81
                                        3.99
## 3
       3
           5.32
                  4.37
                         3.76
                                4.67
                                        4.37
## 4
           8.00
                         5.38
       4
                  6.50
                                6.51
                                        5.60
           6.97
                  6.26
                         5.03
                                6.40
## 5
       5
                                        5.60
## 6
       6
           7.65
                  5.84
                         5.23
                                6.24
                                        5.73
       7
           7.86
                         5.87
                  7.31
                                6.64
                                        6.03
##Put data table into proper form (each unit on one row)
kWh = as.vector(unlist(mydata[,2:6]))
menu_day=rep(1:7, 5)
range = c(rep("Range1", 7), rep("Range2", 7), rep("Range3", 7),
          rep("Range4", 7), rep("Range5", 7))
## Double check that the lengths of these vectors match
length(kWh); length(menu_day); length(range)
## [1] 35
## [1] 35
## [1] 35
## Create the data frame to analyze
unit data energy <- data.frame(kWh, menu day, range)
unit_data_energy
##
       kWh menu_day range
## 1 8.25
                  1 Range1
## 2 5.12
                  2 Range1
## 3
     5.32
                  3 Range1
## 4 8.00
                  4 Range1
## 5 6.97
                  5 Range1
## 6 7.65
                  6 Range1
## 7
     7.86
                  7 Range1
## 8 8.26
                  1 Range2
## 9 4.81
                  2 Range2
                  3 Range2
## 10 4.37
## 11 6.50
                  4 Range2
## 12 6.26
                  5 Range2
## 13 5.84
                  6 Range2
```

```
## 14 7.31
                  7 Range2
## 15 6.55
                  1 Range3
## 16 3.87
                  2 Range3
## 17 3.76
                  3 Range3
                  4 Range3
## 18 5.38
## 19 5.03
                  5 Range3
## 20 5.23
                  6 Range3
## 21 5.87
                  7 Range3
## 22 8.21
                  1 Range4
## 23 4.81
                  2 Range4
## 24 4.67
                  3 Range4
## 25 6.51
                  4 Range4
## 26 6.40
                  5 Range4
                  6 Range4
## 27 6.24
## 28 6.64
                  7 Range4
## 29 6.69
                  1 Range5
## 30 3.99
                  2 Range5
## 31 4.37
                  3 Range5
## 32 5.60
                  4 Range5
## 33 5.60
                  5 Range5
## 34 5.73
                  6 Range5
## 35 6.03
                  7 Range5
```

Problem 1

- (a) Find and interpret the grand mean.
- (b) Find and interpret the differential effect of a menu day, for each of the seven days.
- (c) Find and interpret the differential effect of the range, for each of the five ranges.

Problem 2

Report the estimated additive model for energy use based on this data. Complete the following table of the differences of the observed energy use and the predicted energy use from your model above. Do these residuals indicate a possible interaction effect between the ranges and the menu days?

Menu day	Range 1	Range 2	Range 3	Range 4	Range 5
1					
2					
3					
4					
5					
6					
7					

#if you use R code for problem 2 put it here

Problem 3

Complete the following 2-way ANOVA table for this data "by hand" (just show your work). Then calculate the F-test statistic and corresponding p-value for a test of the significance of an interaction effect between the ranges and the menu days.

Source	df	SS	MS	F
Menu day				
Range				
Interaction				
Error				
Total				

Not solvable without data

We can however fit a 2-way ANOVA table without an interaction effect with the following table:

Source	df	SS	MS	F
Menu day				
Range				
Error				
Total				

More generally, the ANOVA table for a 2-way model without interaction effects and with one observation per combination of factor level is:

Source	df	SS	MS	F
FactorA	(I-1)	$SSA = J \sum_{i} (\bar{Y}_{i.} - \bar{Y}_{})^{2}$	$MSA = \frac{SSA}{I-1}$	$F = \frac{MSA}{MSE}$
FactorB	(J-1)	$SSB = I \sum_{i} (\bar{Y}_{.j} - \bar{Y}_{})^2$	$MSB = \frac{SSB}{J-1}$	$F = \frac{MSB}{MSE}$
Error	(I-1)(J-1)	$SSE = \sum_{i} \sum_{j} (Y_{ij} - \bar{Y}_{i.} - \bar{Y}_{i.} - \bar{Y}_{i.})^{2}$	$MSE = \frac{SSE}{(I-1)(J-1)}$	
Total	(IJ-1)	$Y_{.j} - Y_{.})^{2}$ $SS_{Tot} = \sum_{i} \sum_{j} (Y_{ij} - \bar{Y}_{.})^{2}$		

Here, since K=1 the two-way ANOVA model becomes

$$Y_{ij} = \mu + \alpha_i + \beta_j + \epsilon_{ij}.$$