

difference in incidence of cognitive decline between patients undergoing on-pump and off-pump procedures was found 3 months after CABG, which almost disappeared 1 year after operation [6]. Finally, it is conceivable that a higher incidence of self-reported cognitive failures can be found in older CABG patients with more comorbidity.

In conclusion, we did not find an increase in self-reported cognitive failures in CABG patients 1 year after operation. Moreover, there was no difference in reported cognitive failures between patients undergoing on-pump and off-pump CABG. In contrast, we found a higher rate of self-reported cognitive failures in healthy control subjects than in CABG patients. The present findings suggest that CABG does not result in a substantial proportion of patients with subjectively experienced cognitive decline 1 year after operation, irrespective of the type of surgical technique (on-pump versus off-pump).

References

1. Roach GW, Kanchuger M, Mangano CM, et al. Adverse cerebral outcomes after coronary bypass surgery. *N Engl J Med* 1996;335:1857-63.
2. Van Dijk D, Keizer AMA, Diephuis JC, Durand C, Vos LJ, Hijman R. Neurocognitive dysfunction after coronary artery bypass surgery: a systematic review. *J Thorac Cardiovasc Surg* 2000;120:632-9.
3. Newman MF, Kirchner JL, Phillips-Bute B, et al. Longitudinal assessment of neurocognitive function after coronary artery bypass surgery. *N Engl J Med* 2001;344:395-402.
4. Newman MF, Grocott HP, Mathew JP, et al. Report of the substudy assessing the impact of neurocognitive function on quality of life 5 years after cardiac surgery. *Stroke* 2001;32:2874-81.
5. Diegeler A, Hirsch R, Schneider F, et al. Neuromonitoring, and neurocognitive outcome in off-pump versus conventional coronary bypass operation. *Ann Thorac Surg* 2000;69:1162-6.
6. Van Dijk D, Jansen EWL, Hijman R, et al. Cognitive outcome after off-pump, and on-pump coronary artery bypass graft surgery. *JAMA* 2002;287:1405-12.
7. Newman S, Klinger L, Venn G, Smith P, Harrison M, Treasure T. Subjective reports of cognition in relation to assessed cognitive performance following coronary artery bypass surgery. *J Psychosom Res* 1989;33:227-33.
8. Vingerhoets G, Soete GD, Jannes C. Subjective complaints versus neuropsychological test performance after cardiopulmonary bypass. *J Psychosom Res* 1995;39:843-53.
9. Khatri P, Babyak M, Clancy C, et al. Perception of cognitive function in older adults following coronary artery bypass surgery. *Health Psychiatry* 1999;18:301-6.
10. Rabbitt P. 'Lost, and found'. Some logical and methodological limitations of self-report questionnaires as tools to study cognitive ageing. *Br J Psychiatry* 1990;81:1-16.
11. Van Dijk D, Nierich AP, Eefting FD, et al. The Octopus Study. Rationale and design of two randomized trials on medical effectiveness, safety, and cost-effectiveness of bypass surgery on the beating heart. *Control Clin Trials* 2000;21:595-609.
12. Ponds R WHM, Rozendaal N, Jolles J. The Cognitive Failure Questionnaire: factor structure, effects of age, sex, and education and the relation with cognitive performance and psychosocial variables. Neuropsych Publishers: Maastricht, the Netherlands, 1999.
13. Merckelbach H, Muris P, Nijman H, Jong PJD. Self-reported cognitive failures and neurotic symptomatology. *Pers Individ Differ* 1996;20:715-24.
14. Broadbent DE, Cooper PF, FitzGerald P, Parkes KR. The Cognitive Failures Questionnaire (CFQ) and its correlates. *Br J Psychiatry* 1982;21:1-16.
15. Rasmussen LS, Larsen K, Houx PJ, et al. The assessment of postoperative function. *Act Anaesth Scand* 2001;45:275-89.

INVITED COMMENTARY

"Measure all that is measurable, and make those things measurable, which have hitherto not been measured."

Galileo Galilei

For those investigators trying to assess neurobehavioral outcomes following cardiac surgery, the traditional use of stroke rates to determine outcome has become obsolete unless the number of patients in the study is quite large. This is because stroke, as assessed by neurological examination, is rare in most modern clinical series. Thus it has been necessary to use more intricate testing to measure subtle neurobehavioral outcomes, and this study is no exception. The authors found no differences in quality of life as assessed, at one year postoperatively, by Cognitive Failures Questionnaire (CFQ) and the so-called Worry Index. In addition, there were no differences between patients operated on using cardiopulmonary bypass or an off-pump technique in this small patient sample. The neurobehavioral outcome of this particular data set has been reported in several different journals using several different measuring techniques. This patient group is relatively young and has few co-morbidities, and thus good results should be expected; for this, the authors should be congratulated.

Of interest, a group of "healthy controls" had more cognitive deficits, as measured by the CFQ, than the surgical group. The authors present numerous reasons why this may have been the case but, suffice it to say, the group contained more females and were more highly educated, and thus may have been more aware of their own personal health status and more likely to report cognitive difficulties. Anecdotal reports from patients regarding cognitive failures are fortunately quite rare in our specialty. Thus measuring tools, which rely on patient self-assessment, are the least accurate outcome measure. A more accurate instrument has been developed to assess cognitive performance, which is usually referred to as neuropsychological tests. When batteries of these tests are given pre- and postoperatively to patients having cardiac operations, a substantial number demonstrate postop deficits, particularly early after operation. It has been the goal of cardiac surgical teams to reduce these deficits by altering intraoperative anesthetic and surgical techniques to ameliorate brain dysfunction early after operation and to limit the size of ischemic lesions in the brain induced by particulate embolism.

In our own institution, the neuropsychological deficit

rate has been reduced from >30% at one month following surgery to <20% in our current series. The results of this reported study, and many in the literature, should make surgeons feel good about current techniques for performing cardiac surgery in reference to postop CNS function. It is hoped that continued investigation in this area will permit advanced surgical techniques to be used on high-risk groups of patients who are elderly and have extensive peripheral vessel atheroscle-

rosis, so that improved neurobehavioral results will be obtained.

John W. Hammon, MD

*Department of Cardiothoracic Surgery
Wake Forest University School of Medicine
Medical Center Boulevard
Winston-Salem, NC 27157
e-mail: jhammon@wfubmc.edu.*

Notice From the American Board of Thoracic Surgery

The 2003 Part I (written) examination will be held at the Radisson Hotel O'Hare, Rosemont, Chicago, IL, on November 23, 2003. The closing date for registration is August 1, 2003. Those wishing to be considered for examination must request an application because it is not automatically sent.

To be admissible to the Part II (oral) examination, a candidate must have successfully completed the Part I (written) examination.

A candidate applying for admission to the certifying examination must fulfill all the requirements of the Board in force at the time the application is received.

Please address all communications to the American Board of Thoracic Surgery, One Rotary Center, Suite 803, Evanston, IL 60201; telephone: (847) 475-1520; fax: (847) 475-6240; e-mail: abts_evanston@msn.com.