

growth factors for inducing angiogenesis. In this study, both the angiogenic growth factors were higher after BMCI treatment than after the injection of PBS alone. We assumed that the production of multiple angiogenic growth factors and the endothelial differentiation from these implanted bone marrow cells accelerated the development of microvessels, resulting in the enhanced regional blood flow. We also considered that the enhanced cardiac function after BMCI treatment mainly contributed to the increased blood flow in the LV anterior wall. Although previous studies found that cardiomyocytes were differentiated from bone marrow stem cells [19, 20], this differentiation would be insufficient to affect any increase in cardiac function.

According to past reports, impaired cardiac function can be improved by the implantation of bone marrow cells cultured with 5-azacytidine, but not by that of fresh bone marrow cells, although the implantation of fresh bone marrow cells induces angiogenesis [21]. These conflicting results could be related to differences in the experimental design or implanted cell number.

In conclusion, the implantation of autologous bone marrow cells is an effective method of inducing therapeutic angiogenesis in ischemic heart disease. BMCI is a simple treatment and easy to perform in clinical trials compared with other methods such as the direct injection or the gene transfer of angiogenic cytokines [22, 23].

This work was supported by the Research Fund for the Development of New Medical Treatments of the Ministry of Education and by the Research Grant for Cardiovascular Disease.

References

- Schultz A, Lavie L, Hochberg I, et al. Interindividual heterogeneity in the hypoxic regulation of VEGF. *Circulation* 1999;100:547-52.
- Lee SH, Wolf PL, Escudero R, Deutsch R, Jamieson SW, Thistlethwaite PA. Early expression of angiogenesis factors in acute myocardial ischemia and infarction. *N Engl J Med* 2000;342:626-33.
- Brizzi MF, Formato L, Bonamini R. The molecular mechanisms of angiogenesis: a new approach to cardiovascular disease. *Ital Heart J* 2001;2:81-92.
- Asahara T, Murohara T, Sullivan A, et al. Isolation of putative progenitor endothelial cells for angiogenesis. *Science* 1997;275:964-7.
- Asahara T, Masuda H, Takahashi T, et al. Bone marrow origin of endothelial progenitor cells responsible for postnatal vasculogenesis in physiological and pathological neovascularization. *Circ Res* 1999;85:221-8.
- Hamano K, Li TS, Kobayashi T, Kobayashi S, Matsuzaki M, Esato K. Angiogenesis induced by the implantation of self-bone marrow cells: a new material for therapeutic angiogenesis. *Cell Transplant* 2000;9:439-43.
- Kobayashi T, Hamano K, Li TS, et al. Enhancement of angiogenesis by the implantation of self bone marrow cells in a rat ischemic heart model. *J Surg Res* 2000;89:189-95.
- Ikenaga S, Hamano K, Nishida M, et al. Autologous bone marrow implantation induced angiogenesis and improved deteriorated exercise capacity in a rat ischemic hindlimb model. *J Surg Res* 2001;96:277-83.
- Hamano K, Li TS, Kobayashi T, et al. The induction of angiogenesis by the implantation of autologous bone marrow cells: a novel and simple therapeutic method. *Surgery* 2001;130:44-54.
- Yaoita H, Sakabe A, Maehara K, Maruyama Y. Different effects of carvedilol, metoprolol, and propranolol on left ventricular remodeling after coronary stenosis or after permanent coronary occlusion in rats. *Circulation* 2002;105:975-80.
- Schwarz ER, Pollick C, Meehan WP, Kloner RA. Evaluation of cardiac structures and function in small experimental animals: transthoracic, transesophageal, and intraventricular echocardiography to assess contractile function in rat heart. *Basic Res Cardiol* 1998;93:477-86.
- Litwin SE, Katz SE, Morgan JP, Douglas PS. Serial echocardiographic assessment of left ventricular geometry and function after large myocardial infarction in the rat. *Circulation* 1994;89:345-54.
- Litwin SE, Raya TE, Anderson PG, Litwin CM, Bressler R, Goldman S. Induction of myocardial hypertrophy after coronary ligation in rats decreases ventricular dilatation and improves systolic function. *Circulation* 1991;84:1819-27.
- Scheinowitz M, Kotlyar A, Zimand S, et al. Basic fibroblast growth factor induces myocardial hypertrophy following acute infarction in rats. *Exper Physiol* 1998;83:585-93.
- Takeshita S, Isshiki T, Ochiai M, et al. Endothelium-dependent relaxation of collateral microvessels after intramuscular gene transfer of vascular endothelial growth factor in a rat model of hindlimb ischemia. *Circulation* 1998;98:1261-3.
- Kowallik P, Schulz R, Guth BD, et al. Measurement of regional myocardial blood flow with multiple colored microspheres. *Circulation* 1991;83:974-82.
- Pfeffer MA, Pfeffer JM, Fishbein MC, et al. Myocardial infarct size and ventricular function in rats. *Circ Res* 1979;44:503-12.
- Pfeffer JM, Pfeffer MA, Fletcher PJ, Braunwald E. Progressive ventricular remodeling in rat with myocardial infarction. *Am J Physiol* 1991;260:H1406-14.
- Orlic D, Kajstura J, Chimenti S, et al. Bone marrow cells regenerate infarcted myocardium. *Nature* 2001;410:701-5.
- Jackson KA, Majka SM, Wang H, et al. Regeneration of ischemic cardiac muscle and vascular endothelium by adult stem cells. *J Clin Invest* 2001;107:1395-402.
- Tomita S, Li RK, Weisel RD, et al. Autologous transplantation of bone marrow cells improves damaged heart function. *Circulation* 1999;100(Suppl II):247-56.
- Losordo DW, Vale PR, Symes JF, et al. Gene therapy for myocardial angiogenesis: initial clinical results with direct myocardial injection of phVEGF165 as sole therapy for myocardial ischemia. *Circulation* 1998;98:2800-4.
- Sellke FW, Laham RJ, Edelman ER, Perlman JD, Simons M. Therapeutic angiogenesis with basic fibroblast growth factor: technique and early results. *Ann Thorac Surg* 1998;65:1540-4.

INVITED COMMENTARY

The study by Nishida and colleagues complements other studies published by this group and that of other groups which have examined the effects of implantation of bone marrow cells into the ischemic myocardium or ischemic

hind limb of rats. The distinguishing aspect of the present study is that a novel method was used to create ischemic but not infarcted tissue. The authors used a thin copper wire as a guide to place a ligature around the left

anterior descending coronary artery. The authors contend that this produced a stenotic but open vessel.

Most prior studies in rodent models of ischemia have simply ligated the left anterior descending coronary artery with a suture and the authors have acknowledged the limitation of having an acute MI versus chronically ischemic tissue. Larger animal models of chronic myocardial ischemia used to investigate angiogenesis, improved myocardial performance, or basic physiology of the collateralized myocardium have significant advantages. The two more popular of these methods are the ameroid occluder model and the repetitive embolization model of chronic ischemia, often adapted in the pig, dog, or pony. The ameroid occluder is a hydrophilic casein plastic (ameroid) encased into a stainless steel or hard plastic ring. As the center portion collects water, it occludes the vessel, usually over 10-20 days. The repetitive embolization model uses an injection of microspheres or similar small particles into the coronary artery. Both models have advantages and disadvantages. However, the use of large animal models is very expensive and time consuming.

If the authors have, in fact, developed a new rodent model of chronic myocardial ischemia, this could have a major impact on the progress of future work. However, the authors have not directly assessed whether the artery remains open after the ligature is placed or on the state of the artery after 60 days when flow measurements are made. Whereas, the anterior wall of the hearts was hypokinetic and not dyskinetic according to the authors, my impression is that many or most of these arteries had

occluded within days after placement of the ligature. Further studies will be required to verify the model.

The use of bone marrow, skeletal myocytes, and other transplanted cells to improve vascular density and myocardial perfusion and function is still somewhat novel in practice but not novel in concept. In deed, Philippe Menasche has even transplanted skeletal myoblasts into the hearts of patients. This is a long jump from rat research. However, to date, the results have been modest and many obstacles must be overcome before cell-based therapy will be in widespread clinical practice. Despite these issues, the authors have nicely demonstrated increased end-systolic dimension and improved fractional shortening, increased vascular density, blood flow, and left ventricular thickness in rats having received the bone marrow implantation. They should be congratulated on their work. However, before dramatic conclusions can be drawn from animal experiments, they need to be repeated in a large animal model of chronic myocardial ischemia. Better yet, positive results need to be demonstrated in patients before cell-based therapy is expanded to patient care.

Frank W. Sellke, MD

*Division of Cardiothoracic Surgery
Beth Israel Deaconess Medical Center
Harvard Medical School
110 Francis St, No. LMOB Suite 2A
Boston, MA 02215
e-mail: fsellke@caregroup.harvard.edu*