

Forecasting U.S. GDP, Price Level, and Unemployment Statistics with ARIMA Models

Alexander Yu

Spring 2020

Abstract

In this project, Box-Jenkins ARIMA (Autoregressive Integrated Moving Average) Models are used to construct purely statistical forecasts for four key macroeconomic indicators in the U.S. economy: Real gross domestic product growth (rGDP growth), consumer price index (CPI), and unemployment rate. Data for all series were obtained from the Federal Reserve Bank of St. Louis.

1 Economics Overview

The four variables observed in this project are considered some of the most important figures for assessing the health of the U.S. macroeconomy as well as national economic development as a whole. In the following section, a brief overview of each variable will be provided.

1.1 Real Gross Domestic Product Growth (rGDP Growth)

In macroeconomics, gross domestic product (GDP) represents the market value of all final goods and services produced by an economy during a period of time. It is a statistic that accounts for consumer purchases, government spending, new investment, and net exports (exports minus imports). However, due to the impact of changing price levels over time, the statistic that best represents actual economic output is real GDP (rGDP), which calculates yearly output using the prices of a selected base year. In the case of our data, real GDP growth is expressed as the percent change from the preceding period. Data is reported quarterly each year in January, April, July, and October.

1.2 Consumer Price Index (CPI)

The Consumer Price Index measures the average price for goods and services purchased by consumers, including food, clothing, shelter, transportation, service fees, and sales taxes. The index can be used to determine changes in price levels over time, such as periods of inflation and deflation. For example,

a significant increase in CPI over a short period of time may be indicative of a period of inflation while a significant decrease in CPI may signal a period of deflation. In our project, the index used is the Consumer Price Index for All Urban Consumers, a seasonally-adjusted index published each month that tracks the price of goods and services purchased by urban consumers relative to prices within a baseline time period (1982 – 1984 = 100). The index accounts for the spending of nearly 88 percent of the U.S. population, including wage workers, salaried workers, self-employed workers, retirees, the unemployed, and people not in the labor force.

1.3 Unemployment Rate

Derived from data collected by the Bureau of Labor Statistics through its Current Population Survey (Household Survey), the unemployment rate represents the percentage of the labor force that unemployed after adjusting for seasonal. The definition of unemployment used for this series refers to people 16 years of age and older who are not employed, available to work, and actively looking for a job within the last four weeks.

2 Statistical Overview

In forecasting, ARIMA models can be used to produce short-term predictions for a univariate (single variable) time series. As a forecasting method, the ARIMA(p, d, q) model combines three different processes: an autoregressive component (p AR terms), differencing (I, integration of order d), and a moving average component (q MA terms). Thus, in order to understand how ARIMA model can be used, it is first necessary to understand the properties of each component as well as the Box-Jenkins ARIMA modeling approach.

2.1 Autoregressive Models (AR)

In an ARIMA model the autoregressive component is derived from an autoregressive model, which can be used to forecast \hat{y}_t , the predicted y value at time t .¹ In other words, an autoregressive model uses the weighted lagged (past) values of y to predict future values of y . It is a regression against itself, hence the term autoregression.² This can be represented as:

$$\begin{aligned}\hat{y}_t &= \mu + \phi_1 y_{t-1} + \dots + \phi_p y_{t-p} + \epsilon_t \\ &= \mu + \left(\sum_{i=1}^p \phi_i y_{t-i} \right) + \epsilon_t\end{aligned}$$

where μ is a constant, ϕ_i are coefficients (weights) for lagged values of y_i , and ϵ_t is error due to white noise.

¹Penn State University

²Hyndman and Athanasopoulos, FPP2

2.2 Moving Average Models (MA)

ARIMA models also incorporate elements of moving average models, which use past *forecast errors* ($y_i - \hat{y}_i$) rather than past *values* of the variable being forecasted.³ As an equation, this can be represented as:

$$\begin{aligned}\hat{y}_t &= \mu + \theta_1 \epsilon_{t-1} + \dots + \theta_q \epsilon_{t-q} + \epsilon_t \\ &= \mu + \left(\sum_{i=1}^q \theta_i \epsilon_{t-i} \right) + \epsilon_t\end{aligned}$$

where μ is a constant, θ_i are coefficients for past forecast errors ϵ_{t-q} , and ϵ_t is error due to white noise.

2.3 Autocorrelation and Partial Autocorrelation

In determining the correct parameters of a $\text{ARIMA}(p, d, q)$ model, autocorrelation and partial autocorrelation plots are commonly used to describe the relationships between an observation and its lagged (past) values. In the following section, a brief description of each type of plot will be provided

- **Autocorrelation:** In time series analysis, autocorrelation summarizes the relationship between each observation and its lagged values. Expressed as a graph, this summarizes whether each observation can be effectively explained by previous values. Autocorrelation plots are commonly used for determining stationary as well as the correct order of difference.
- **Partial Autocorrelation:** Similar to autocorrelation, partial autocorrelation summarizes the relationship between each observation and its lagged values *while accounting for the autocorrelation of lower lags*. (For example, a lag 3 partial autocorrelation excludes the effects of lag 2 and lag 1 autocorrelations.) When partial autocorrelation plots are used in conjuncture with autocorrelation plots, they can be used to determine the correct order of AR and MA terms in a model, a process that will be explained in a later section.


2.4 Differencing (Integration of Order d)

Another key component of ARIMA models is the uses of differencing, a process that transforms a nonstationary time series into a stationary one by subtracting lagged (past) values from each observation.

In statistics, a time series is said to be stationary when its statistical properties of each observation are not correlated with past observations or dependent on time. In other words, a stationary time series is a series that has both constant mean and variance over time where each observed value cannot be effectively explained by previous values. As such, the autocorrelation between

³Hyndman and Athanasopoulos

Figure 1: The ACF plots of a non-stationary (left) and stationary series (right)⁵



each observation and lagged observations is statistically insignificant within a certain level of confidence. Hence, the graph of the autocorrelations between each observation and its lags should also display a pattern that is statistically insignificant.

Since the statistical properties of a stationary time series are assumed to remain the same in the future, it is easier to predict compared to a nonstationary series.⁶ Later, to obtain predictions for the original time series, these stationary series can be untransformed by reversing the processes used in differencing.

In nonseasonal ARIMA models, the variable d represents the number of first differences used to render a series into a stationary one. Let first difference be defined as such:

$$y_t - y_{t-1}$$

Commonly, the number of first differences used in an ARIMA model will be either 0, 1, or 2.

- When the time series in question is already stationary, no first differences are needed and $d = 0$.
- When first differences are required, $d = 1$ or 2. It is also important to note that $d = 2$ refers to the first difference of the first difference, or $(y_t - y_{t-1}) - (y_{t-1} - y_{t-2})$, not $y_t - y_{t-2}$.

Generally, the order of differencing in an ARIMA model rarely exceeds $d = 2$ due to the possibility overdifferencing, which induces negative autocorrelation in the time series.⁷

⁶Nau

⁷Feldman, Non-stationary models: ARIMA. Transformations

2.5 ARIMA Models

An ARIMA model combines the properties of AR and MA models while applying first-differences when necessary. Hence, a nonseasonal ARIMA(p, d, q) model can be expressed as the following equation⁸:

$$\begin{aligned}\Delta \hat{y}_t &= \mu + \phi_1 \Delta y_{t-1} + \dots + \phi_p \Delta y_{t-p} + \theta_1 \epsilon_t + \dots + \theta_q \epsilon_{t-q} + \epsilon_t \\ &= \mu + \left(\sum_{i=1}^p \phi_i \Delta y_{t-i} \right) + \left(\sum_{i=1}^q \theta_i \epsilon_{t-i} \right) + \epsilon_t\end{aligned}$$

where μ is a constant, ϕ are the weights for the AR component, Δy are terms in the series after d first differences (lagged values), ϵ_t is error due to white noise, θ are the weights for the MA component, and ϵ_{t-q} are past forecast errors of lagged values (lagged errors).

At this point, it's also worth noting that pure AR(p) and MA(q) models are special cases of ARIMA(p, d, q) models where

$$\begin{aligned}\text{AR}(p) &\leftrightarrow \text{ARIMA}(p, 0, 0) \\ &\text{and} \\ \text{MA}(q) &\leftrightarrow \text{ARIMA}(0, 0, q)\end{aligned}$$

It is also important to note that any differencing performed during the forecast must be reversed after the forecast is complete since the original time series is undifferenced.⁹ However, most statistical software and packages (like the Forecast package in R) perform this task by default.

⁸National Institute of Standards and Technology

⁹Nau

2.6 The Box-Jenkins Method

In this project, the method used select and determine ARIMA models for forecasting is the Box-Jenkins method. First proposed by statisticians George Box and Gwilym Jenkins in 1970, the Box-Jenkins method outlines a process for identifying parameters, estimating coefficients, and checking the appropriateness of ARIMA models in time-series analysis.¹⁰ In greater detail, the three stages of the Box-Jenkins process are as follows:

1. **Identification:** This step can be further separated into two processes:

(a) **Determining the number of differences (d) required to stationize the time series:**

In constructing an ARIMA model, it is first necessary to determine the lowest order of differencing needed to render the time series stationary. As such, the correct order of differencing is the lowest number of first-differences needed to transform a time series into a series with a constant variance, well-defined mean, and rapidly decaying ACF plot.¹¹

In general, there are three rules to follow when determining the correct order of differencing:¹²

- **Rule 1: If the autocorrelations of a series remain positive and statistically significant up to higher lags, the series may need a higher order of differencing.** Since differencing induces negative autocorrelations within a time series, additional first differences reduce the magnitude of highly positive autocorrelations to within statistically insignificant bounds, thus rendering the series stationary.
- **Rule 2: If the autocorrelations of a series are statistically insignificant at lower lags and remain insignificant up to higher lags, the series does not need additional differencing.** By the definition of stationary, a series with a constant mean, variance, and statistically insignificant autocorrelations is said to be stationary.
- **Rule 3: If the autocorrelations of a series are negative and statistically significant at lower lags, the series may be overdifferenced.** In the case of this, it is important to check whether the time series in question has undergone any prior transformations before differencing and whether such transformations are necessary for rendering the series stationary

Alternatively, unit root tests (hypothesis tests of stationarity can be used to determine whether a time series in question is either stationary or not stationary at a given confidence level. If the series

¹⁰Brownlee, A Gentle Introduction to the Box-Jenkins Method for Time Series Forecasting

¹¹Nau

¹²Nau

is not stationary, and first difference is taken and the unit root test repeated. This process is repeated until the unit root test determines that the series is stationary.¹³ In this project, the built-in Kwiatkowski-Phillips-Schmit-Shin (KPSS) unit root test from the Forecast package is used to test for stationarity alongside traditional methods involving autocorrelation plots.

(b) **Determining the number of AR (p) and MA (q) terms**

After differencing, the times series should appear approximately stationary but may appear slightly overdifferenced or underdifferenced. To recognize and address this issue, the autocorrelation function (ACF) and partial autocorrelation (PACF) graphs of the differenced series can be used to determine whether the series is over/underdifferenced as well as the appropriate number of AR (p) and MA (q) terms to add.

Generally, there are two rules for determining the appropriate number of AR and MA terms to use in an ARIMA model, both of which require the use of a ACF or PACF plot:¹⁴

- **Rule 1: If the PACF of the stationarized series becomes statistically insignificant at lag n and the lag 1 autocorrelation is positive, the series is said to be slightly underdifferenced.** The lag n where the PACF becomes insignificant should be set as the number of AR terms ($p = n$).
- **Rule 2: If the ACF of the stationarized series becomes statistically insignificant at lag m and the lag 1 autocorrelation is negative, the series is said to be slightly overdifferenced.** The lag m where the ACF becomes insignificant should be set as the number of MA terms ($q = m$).

2. **Estimation:** Estimate all AR and MA coefficients, ϕ and θ . In this project, the Forecast package in R automatically approximates the coefficients using maximum likelihood estimation and sums of squares.¹⁵
3. **Validation/Model Checking:** After the ARIMA model is constructed, it is important to check two diagnostics:¹⁶

- **Overfitting:** When an model overfits the training data (the imported data), the model captures trends that can be attributed to random noise. In other words, an overfit model may predict data within the training sample too well and may perform poorly with out of sample data.
- **Distribution of Residuals:** A well-constructed model should have a distribution of residuals that resembles white noise with a symmetrical variance and mean of zero. This ensures that the forecasts

¹³Hyndman and Athanasopoulos


¹⁴Nau

¹⁵Robert Hyndman, <https://www.rdocumentation.org/packages/forecast/versions/8.12>

¹⁶Brownlee

produced by the forecast may not suggest a skew or bias. To check for this, one can observe the behavior of the ACF and PACF plots of the residuals as a time series.

Figure 2: The Box-Jenkins method summarized as a flowchart



3 Possible Models for Forecasting


3.1 Data Overview

As discussed in Section 1, this project focuses primarily on forecasting four economic statistics: the real GDP growth rate, consumer price index, and unemployment rate.

3.2 Real GDP Growth


The initial assessment of U.S. real GDP growth data was done with a visual assessment of the undifferenced time plot of the series as the undifferenced plot displays an inconsistent variance (changes between values are not constant). From the first graph in Figure 3, it is clear that the rGDP series is not a stationary series and needs to undergo differencing. The nonstationary of this series is confirmed by running the KPSS test of stationarity, which returns a test statistic of 0.7339, a value that exceeds the critical value of 0.347 for a significance level of $\alpha = 0.10$. This shows that the null hypothesis of stationarity can be rejected and first differences are needed. However, after performing a single first-difference, the series still exhibits inconsistent variance, as the variance of the series is significantly higher before the year 1983 than after that year. As such, the condition of stationary cannot be met when looking at the series in its entirety.

Figure 3: The time plot of the full rGDP growth series



To address this issue, the data from before 1983 has been omitted from the forecast altogether. Figure 4 shows both the undifferenced and differenced series post 1983. Upon visual inspection, it is clear that the variance of the differenced series remains relatively more consistent during this abbreviated time frame than during the full time frame.

Figure 4: The time plot of the rGDP growth series after 1983



For forecasting rGDP growth, two candidate models have been considered, $ARIMA(0,1,1)$ and $ARIMA(1,1,1)$. The former was produced using the `auto.arima` function from the `Forecast` package while the latter was chosen manually after considering the ACF and PACF plot of the differenced series in Figure 5.

Figure 5: The ACF and PACF plot of the differenced series





Figure 6: rGDP growth rate forecast with $ARIMA(0,1,1)$ and $ARIMA(1,1,1)$



In Figure 6 and all subsequent forecast graphs (Figures 11 and 15), the 85, 90, and 95 percent confidence intervals are represented by the shaded blue bands surrounding the black mean forecast line. Out-of-sample data released after September 2019 is represented by small black dots.

Table 1: ARIMA(0,1,1) parameter estimates

Coefficient	Value	Standard Error
MA1	-0.6519	0.0822

Table 2: ARIMA(1,1,1) parameter estimates

Coefficient	Value	Standard Error
AR1	0.2187	0.1745
MA1	-0.8297	0.1336

Using the two candidate models above, the following table shows the mean forecast values for rGDP growth (values on the black line in Figure 6) up to the Q2 of 2020.

Table 3: ARIMA(0,1,1) forecasts

Period	Mean Forecast
2019 Q3	2.383027
2019 Q4	2.383027
2020 Q1	2.383027
2020 Q2	2.383027

Table 4: ARIMA(1,1,1) forecasts

Period	Mean Forecast
2019 Q3	2.337689
2019 Q4	2.411554
2020 Q1	2.427711
2020 Q2	2.431245

Checking the test diagnostics, the candidate model ARIMA(0,1,1) appears to be a better model for forecasting the rGDP growth rate. With a lower AIC of (Akaike Information Criterion), the ARIMA(0,1,1) achieves a balance between maintaining goodness-of-fit and reducing overfitting.¹⁷

Table 5: ARIMA(0,1,1) model diagnostics


σ^2	Log Likelihood	AIC
4.796	-319.18	642.37

Table 6: ARIMA(1,1,1) model diagnostics

σ^2	Log Likelihood	AIC
4.716	-318.6	643.2

Checking the residuals of the chosen candidate model, it appears that the residuals are stationary and exhibit constant variance. Therefore this model can be used for forecasting.

Figure 7: ARIMA(0,1,1) residuals




¹⁷Sachin Date, The Akaike Information Criterion (<https://towardsdatascience.com/the-akaike-information-criterion-c20c8fd832f2>)

3.3 Consumer Price Index


The initial assessment of the U.S. Consumer Price Index data was also done with a visual assessment of the undifferenced time plot of the series (Figure 8, top). Similar to the rGDP series, the CPI series is not a stationary series and needs to undergo differencing since the undifferenced series clearly displays an upward trend. The KPSS test of stationarity confirms the nonstationarity of the CPI series, as the test statistic returned is 12.3099, a value that far exceeds the critical value of 0.347 for a level of significance of $\alpha = 0.10$. As such, the null hypothesis of stationarity can be rejected and first differences are needed. Running the KPSS tests a second time after applying one first difference yields a test statistic of 3.5488, a value that still exceeds the critical value. Running the KPSS test a third time after applying a total of 2 first differences yields a value of 0.0058, a value that lets us reject the null hypothesis at $\alpha = 0.01$ with a critical value of 0.739.

Figure 8: The time plot of the full CPI series



However, after performing a single first-difference, the series still exhibits inconsistent variance, as the variance of the series is significantly higher after the year 1999 than before that year. As such, the condition of stationary cannot be met when looking at the series in its entirety. Omitting the observations before January 1999 addresses this issue, however a new order of differencing is required. Running the KPSS test on the abbreviated series returns the test statistic of 4.2231, a value that exceeds the above critical value of $\alpha = 0.10$. After applying one single difference, the test statistic returned is 0.0838, a value that lets us reject the null hypothesis of stationarity at $\alpha = 0.01$. As such, the proper order of differencing for the abbreviated CPI series is $d = 1$. The undifferenced and differenced abbreviated CPI series is shown in Figure 9.

Figure 9: The time plot of the CPI series after 1999



An important observation to note about the post-1999 series is that there is a significant negative spike in the differenced series between August and December 2008. This spike can likely be attributed the effects of the Great Recession, which occurred between late 2007 and early 2009. High unemployment, declines in commodity prices, and various economic factors put unusually high pressure on the U.S. economy, thus leading to this period of deflation.¹⁸ However, as the variance of the rest of the differenced series is relatively stable and consistent, the forecast may proceed.

¹⁸Investopedia, Were There Any Periods of Major Deflation in U.S. History? (<https://www.investopedia.com/ask/answers/040715/were-there-any-periods-major-deflation-us-history.asp>)

For forecasting CPI, two candidate models have been considered, $\text{ARIMA}(3,1,2)$ and $\text{ARIMA}(3,1,3)$. The former was produced using the `auto.arima` function from the `Forecast` package while the latter was chosen manually after considering the ACF and PACF plot of the differenced series in Figure 10. Note that the candidate model produced by `auto.arima` includes a drift term since the function recognizes that the mean of the series is increasing over time.

Figure 10: The ACF and PACF plot of the differenced series




Figure 11: CPI forecast with $\text{ARIMA}(3,1,2)$ and $\text{ARIMA}(3,1,3)$




Table 7: ARIMA(3,1,2) parameter estimates

Coefficient	Value	Standard Error
AR1	-0.1496	0.2225
AR2	-0.5027	0.1656
AR3	0.1836	0.1218
MA1	0.6647	0.2124
MA2	0.6477	0.1590
Drift	0.3690	0.0528

Table 8: ARIMA(3,1,3) parameter estimates

Coefficient	Value	Standard Error
AR1	0.7571	0.2723
AR2	0.0254	0.3662
AR3	0.2173	0.1593
MA1	-0.2309	0.2696
MA2	-0.3669	0.2251
MA3	-0.3863	0.1363

Using the two candidate models above, the following table shows the mean forecast values for CPI up to the June of 2020.

Table 9: ARIMA(3,1,2) forecasts

Period	Mean Forecast
2019 October	256.6048
2019 November	257.0836
2019 December	257.4405
2020 January	257.7337
2020 February	258.1402
2020 March	258.5395
2020 April	258.8711
2020 May	259.2374
2020 June	259.6311

Table 10: ARIMA(3,1,3) forecasts

Period	Mean Forecast
2019 October	256.5921
2019 November	257.0186
2019 December	257.3743
2020 January	257.7053
2020 February	258.0576
2020 March	258.4101
2020 April	258.7578
2020 May	259.1065
2020 June	259.4559

With regards to test diagnostics, the candidate model ARIMA(3,1,2) appears to be a better model for forecasting the CPI since it has a lower AIC.

Table 11: ARIMA(3,1,2) model diagnostics


σ^2	Log Likelihood	AIC
0.2869	-194.17	402.34

Table 12: ARIMA(3,1,3) model diagnostics

σ^2	Log Likelihood	AIC
0.2844	-197.39	408.79

Checking the time plot of the residuals of the ARIMA(3,1,2) model in Figure 12, it appears that the residuals are stationary. With the exception of several outliers in late 2008, the residuals exhibit a relatively consistent variance and this model can be used for forecasting.


Figure 12: ARIMA(3,1,2) residuals



3.4 Unemployment Rate

The assessment of the U.S. unemployment rate data was also done with a visual assessment of the undifferenced time plot of the series. Similar to the previous series, the unemployment rate is not a stationary series and needs to undergo differencing since the undifferenced series displays multiple peaks and troughs. The KPSS test of stationarity confirms the nonstationarity of the unemployment rate series, as the test statistic returned is 1.3527, a value that far exceeds the critical value of 0.347 for a level of significance of $\alpha = 0.10$. As such, the null hypothesis of stationarity can be rejected and first differences are needed. However, running the KPSS test after applying a first difference yields a value of 0.0991, a value that lets us reject the null hypothesis at $\alpha = 0.01$ with a critical value of 0.739.

Figure 13: The time plot of the full unemployment rate series



Note the fluctuations in differenced series in the period immediately before 1950. While these spikes are significant, since the variance of the rest of the differenced series is relatively stable and consistent, the forecast may still proceed.

For forecasting CPI, two candidate models have been considered, $\text{ARIMA}(3,1,2)$ and $\text{ARIMA}(3,1,3)$. The former was produced using the `auto.arima` function from the `Forecast` package while the latter was chosen manually after considering the ACF and PACF plot of the differenced series in Figure 14. Note that the candidate model produced by `auto.arima` includes a drift term since the function recognizes that the mean of the series is increasing over time.

Figure 14: The ACF and PACF plot of the differenced series




Figure 15: Unemployment rate forecast with $\text{ARIMA}(2,1,2)$ and $\text{ARIMA}(4,1,1)$




Table 13: ARIMA(2,1,2) parameter estimates

Coefficient	Value	Standard Error
AR1	1.6531	0.0394
AR2	-0.7774	0.0444
MA1	-1.6356	0.0393
MA2	0.8542	0.0472

Table 14: ARIMA(4,1,1) parameter estimates

Coefficient	Value	Standard Error
AR1	0.4096	0.1461
AR2	0.2173	0.0370
AR3	0.0699	0.0517
AR4	0.0343	0.0464
MA1	-0.3954	0.1426

Using the two candidate models above, the following table shows the monthly mean forecast values for the unemployment rate up to the June of 2020.

Table 15: ARIMA(2,1,2) forecasts

Period	Mean Forecast
2019 October	3.476366
2019 November	3.451822
2019 December	3.429623
2020 January	3.412006
2020 February	3.400142
2020 March	3.394225
2020 April	3.393666
2020 May	3.397344
2020 June	3.403857

Table 16: ARIMA(4,1,1) forecasts

Period	Mean Forecast
2019 October	3.503494
2019 November	3.461471
2019 December	3.431033
2020 January	3.402826
2020 February	3.381842
2020 March	3.363551
2020 April	3.348484
2020 May	3.335906
2020 June	3.325482

With regards to test diagnostics, the candidate model ARIMA(4,1,1) appears to be a better model for forecasting the CPI since it has an AIC closer to zero.

Table 17: ARIMA(2,1,2) model diagnostics


σ^2	Log Likelihood	AIC
0.03746	193.65	-377.31

Table 18: ARIMA(4,1,1) model diagnostics

σ^2	Log Likelihood	AIC
0.03787	187.16	-362.3

Checking the time plot of the residuals of the ARIMA(4,1,1) model in Figure 16, the residuals appear to be stationary and the model can be used for forecasting.

Figure 16: ARIMA(4,1,1) residuals



4 Conclusion and Summary

In this project, the ARIMA models used to produce near-term forecasts produced results were only effective to a limited extent. For all three series, actual statistics released by the St. Louis Fed remained relatively in-line with the mean forecast between the start of the forecast in October 2019 to February 2020. All actual statistics released within this time frame fell within the 85 percent confidence intervals for all three datasets. Additionally, some of our models performed very well such as our two models for forecasting the unemployment rate, which both produced mean forecasts in line with out-of-sample data until to February 2020. Admittedly, several of the confidence intervals produced by the models is quite large, such as those produced for the rGDP growth rate series (with the narrowest being -0.78 to +5.46 for October 2019). Similar trends can be seen in CPI. As such, it is not surprising that much of the actual statistics fall within even the narrowest of intervals produced by these models, especially during times of relative economic stability.

However, when compared to data released after February 2020, the forecasts performed considerably worse, even though ARIMA models tend to produce wider confidence intervals for longer term forecasts. This can be primarily attributed to the largely unexpected coronavirus epidemic that has crippled the United State economy starting from late February and early March 2020. Due to COVID-19, all three economic statistics have begun to exhibit recessionary behavior, with CPI dropping to 255.90 in April from 259.05 in February and Q1 rGDP growth projected to be -4.8 percent. In terms of the confidence intervals produced by models in our project, the CPI data barely lies within April's forecasted 95 percent confidence interval (254.7326 to 263.0097) and GDP lies outside of the Q1 rGDP projection (-2.266998 to 7.122419). However, the most striking difference between actual data and our forecasts come from April's unemployment statistics, which exceed even widest of confidence intervals. At 14.7 percent, April's unemployment rate lies far outside the upper bound of the corresponding 95 percent confidence interval produced by our forecast (2.0226006 to 4.764732). Even when looking at a corresponding 99.99 percent confidence interval for the same period (0.1925654 to 6.479246), the actual data lies well beyond its upper bound. As only 0.01 percent of observations are expected to lie outside this confidence interval, April's unemployment data is undoubtedly significant, both statistically and economically.

Yet, the most important observation gained from comparing the effectiveness of our model during times of stability and its effectiveness during times of economic distress is the following: forecasts produced by rudimentary models are 'ballpark estimates' at best, and utterly useless at worst. After all, a model that merely satisfies statistical conditions is not enough to predict the future with absolute certainty, especially the future contains unforeseeable contingencies and unpredictable acts of god.

5 Additional Acknowledgements and Information

Without the support of my mentor, Mr. Carey Kopeikin, this project would not have been possible. Thank you for your guidance in statistics and research over the past year!

Additionally the technical elements of project, such as R scripts and LaTeX source code, can be found in the following GitHub repository.

<https://github.com/ProfuTofu/CRH-2020-Spring-Capstone>

Bibliography

- Contreras, J., R. Espinola, F.J. Nogales, and A.J. Conejo. "ARIMA Models to Predict Next-Day Electricity Prices." *IEEE Transactions on Power Systems* 18, no. 3 (August 2003): 1014–20. <https://doi.org/10.1109/TPWRS.2002.804943>.
- Date, Sachin. "The Akaike Information Criterion." Medium, March 10, 2020. <https://towardsdatascience.com/the-akaike-information-criterion-c20c8fd832f2>.
- Feldman, Raya E. "Non-Stationary Models: ARIMA. Transformations." University of California, Santa Barbara. Accessed May 20, 2020. <http://feldman.faculty.pstat.ucsb.edu/174-03/lectures/111.pdf>.
- Hyndman, Rob J., and George Athanasopoulos. "Forecasting: Principles and Practice." Accessed May 20, 2020. <https://otexts.com/fpp2/>.
- Investopedia. "Were There Any Periods of Major Deflation in U.S. History?" Investopedia. Accessed May 20, 2020. <https://www.investopedia.com/ask/answers/040715/were-there-any-periods-major-deflation-us-history.asp>.
- Jason Brownlee. "A Gentle Introduction to the Box-Jenkins Method for Time Series Forecasting," January 13, 2017. <https://machinelearningmastery.com/gentle-introduction-box-jenkins-method-time-series-forecasting/>.
- National Institute of Standards and Technology. "Box-Jenkins Models." Accessed May 20, 2020. <https://www.itl.nist.gov/div898/handbook/pmc/section4/pmc445.htm>.
- Nau, Robert. "Statistical Forecasting: Introduction to ARIMA Models." Accessed May 20, 2020. <https://people.duke.edu/~rnau/411arim.htm>.
- Penn State Eberly College of Science. "ARIMA Models." PennState: Statistics Online Courses. Accessed May 20, 2020. <https://online.stat.psu.edu/stat501/lesson/14/14.5/14.5.1>.
- Salam, Muhammad Abdus, Shazia Salam, and Mete Feridun. "MODELING AND FORECASTING PAKISTAN'S INFLATION BY USING TIME SERIES ARIMA MODELS," n.d., 34.
- St. Louis Fed. "Federal Reserve Economic Data." Accessed May 20, 2020. <https://fred.stlouisfed.org/>.